

IBUPROFEN- ibuprofen tablet
WALGREENS

Walgreens Ibuprofen 200mg Caplets

DRUG FACTS

Active ingredient (in each caplet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/Fever reducer

Uses

- temporarily relieves minor aches and pains due to:

headache

muscular aches

minor pain of arthritis

toothache

backache

the common cold

menstrual cramps

- temporarily reduces fever

Warnings

Allergy alert:

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood-thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

you have symptoms of heart problems or stroke:

- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling

pain gets worse or lasts more than 10 days

fever gets worse or lasts more than 3 days

redness or swelling is present in the painful area

any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely

directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222)

Directions

do not take more than directed

the smallest effective dose should be used

adults and children 12 years and older:

- take 1 caplet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 caplet, 2 caplets may be used
- do not exceed 6 caplets in 24 hours, unless directed by a doctor

children under 12 years:

- ask a doctor

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, FD&C yellow #6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide

Questions or comments? call **1-877-290-4008**

50 COUNT 612R

100 COUNT 612R

Drug Facts (continued)

Lot No.:
Exp Date:

IS DISTRIBUTED BY: WALGREEN CO.
DO NOT WRITE OR REELECT IN 11 STATES

Active ingredient (in each capsule)	Purpose
100 mg of ampicillin sodium	Antibiotic
250 mg of ampicillin sodium	Antibiotic
500 mg of ampicillin sodium	Antibiotic
1000 mg of ampicillin sodium	Antibiotic

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

- temporarily relieves minor aches and pains due to:
 - backache ● headache ● menstrual cramps
 - minor pain of arthritis ● muscular aches
 - the common cold ● toothache
- temporarily reduces fever

- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke

Symptoms may include:

- asthma (wheezing)
- hives
- rash
- stomach, cold and sick
- itchy eyes, itching throat, skin and neck
- medical help right away.

Someone's breathing worsening. This product contains an NSAID, which may cause severe stomach bleeding. The danger is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood-thinning (anticoagulant) or steroid drug
- take other drugs containing aspirin, ibuprofen, naproxen, or other NSAIDs (aspirin, ibuprofen, naproxen, or others)
- use 3 or more alcoholic drinks every day while taking this product
- drink milk for a longer time than directed

- you are taking a diuretic
- Ask a doctor or pharmacist before use if you are
 - taking aspirin for heart attack or stroke, because aspirin may decrease the level of aspirin
 - under a doctor's care for any serious condition
 - taking any other drug

When using this product

- Use with food or milk. Stomach upset occurs
- Don't use an alkali tablet if:
 - you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have blood or black stools
 - have stomach pain that does not get better
 - you have symptoms of heart and kidney trouble
 - feel pain
 - trouble breathing

Heart attack and stroke warning: MSADs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning signs to you
- you have a history of stomach problems, such as heartburn

- weakness in one part or side of the body
- slurred speech ● big swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area

■ any new symptoms appear

<ul style="list-style-type: none"> ■ do not take more than five doses ■ the smallest effective dose should be used 	adults and children 12 years and over: <ul style="list-style-type: none"> ■ take 1 capsule every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 capsule, 2 capsules may be used ■ do not exceed 6 capsules in 24 hours, unless directed by a doctor
	children under 12 years: <ul style="list-style-type: none"> ■ ask a doctor

Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)

Compare to Motrin® IB
Caplets active ingredient††

NDC 0363-9612-01

IBUPROFEN TABLETS USP, 200 mg / PAIN RELIEVER / FEVER REDUCER (NSAID)

CAPLETS
100 CAPLETS**
(***CAPSULE-SHAPED TABLETS)
SEE NEW WARNINGS INFORMATION

Varnish/Print Omit Area

E827

Varnish

Omrit

Omitempty

Drug Facts (continued)

Inactive ingredients cellulose, silicon dioxide, croscarmellose sodium, FD & C yellow #6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, polybutyrate starch, talc, titanium dioxide.

Questions or comments? Call 1-877-290-4008

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED PLOT SEAL OVER THE MONTH OF THE BOTTOM LEFT CORNER IS BROKEN OR MISSING.

Drug Facts (continued)

Warnings ■ If you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer, do not take it after heart surgery.

Ask a doctor before use if: ■ you have problems or serious side effects from taking pain relievers, ever or now ■ you are taking blood thinners ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver problems, kidney disease, asthma, or had a stroke ■ you are taking a diuretic ■ ask a doctor if ibuprofen will be used if you are ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug

When using this product: ■ take with food or milk if stomach upset occurs ■ stay up and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ heartburn ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke ■ chest pain ■ trouble breathing ■ pain or tenderness on the left side of your stomach ■ heartburn ■ you are taking blood thinners ■ you are taking aspirin ■ you are taking other drugs ■ you are taking more than 10 days ■ liver gets worse or feels more than 3 days ■ redness or swelling is present in the genital area ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ adults and children 12 years and over: ● take 1 caplet every 4 to 6 hours while symptoms persist ● if pain or fever does not respond to 1 caplet, 2 caplets may be used ● do not exceed 6 caplets in 24 hours, unless directed by a doctor ■ children under 12 years: ● ask a doctor

Other information ■ read all warnings and directions before use ■ store at 20-25°C (68-77°F)

Inactive ingredients croscellose, silica, dextrose, croscellose sodium, DCC, yellow #6 aluminum lake, magnesium stearate, monostelline or linoleic, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide

Questions or comments? call 1-877-234-4448

Adhesive Area

Waldgreen's

Compare to Motrin® IB Caplets active ingredient††

NDC 0383-961-2-26

Ibuprofen 200

IBUPROFEN TABLETS USP, 200 mg / PAIN RELIEVER / FEVER REDUCER (NSAID)

CAPLETS

225 CAPLETS** ACTUAL SIZE
(**CAPLETS=SWAGED TABLETS)

SEE NEW WARNINGS INFORMATION

Drug Facts

Active ingredient (in each caplet): Ibuprofen, 200 mg (NSAID)††

Purpose: Pain reliever/fever reducer

Warnings: Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ asthma (wheezing) ■ hives ■ shock ■ skin redness ■ facial swelling ■ reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ take other drugs (especially blood thinners) or steroid drug ■ take a blood thinner (warfarin) or aspirin, or other NSAID (aspirin, ibuprofen, naproxen, or others) ■ take 3 or more alcohol drinks every day while using this product ■ take more or for a longer time than directed ■ heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Directions: Take 1 caplet with or without food, 2 to 3 times a day as needed. Do not take more than directed. Do not take for more than 10 days unless directed by a doctor.

Other information: Read all warnings and directions before use. Store at 20-25°C (68-77°F). Inactive ingredients: croscellose, silica, dextrose, croscellose sodium, DCC, yellow #6 aluminum lake, magnesium stearate, monostelline or linoleic, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide.

Questions or comments? call 1-877-234-4448

Lot No.: 1917 20481
Exp. Date:

Varnish Omit Area

Adhesive Area

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use ■ if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer ■ right before or after heart surgery

Ask a doctor before use if ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic

Ask a doctor or pharmacist before use if you are ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin ■ under a doctor's care for any serious condition ■ taking any other drug

When using this product ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
- you have stomach pain that does not get better
- you have symptoms of heart problems or stroke: ● chest pain ● trouble breathing ● weakness in one part or side of body ● slurred speech ● leg swelling
- pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present in the painful area ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions ■ **do not take more than directed** ■ the **smallest effective dose should be used**

adults and children 12 years and older:	■ take 1 caplet every 4 to 6 hours while symptoms persist ■ if pain or fever does not respond to 1 caplet, 2 caplets may be used
children under 12 years:	■ do not exceed 6 caplets in 24 hours, unless directed by a doctor ■ ask a doctor

Other information ■ read all warnings and directions before use ■ store at 20-25°C (68-77°F)

Inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, FD&C yellow #6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, tlc, titanium dioxide

Questions or comments? call 1-877-290-4008

VALUE SIZE NDC 0363-9612-50

Walgreens

Compare to the active ingredient in Motrin® IB Caplets*

Ibuprofen 200

IBUPROFEN TABLETS USP, 200 mg / PAIN RELIEVER FEVER REDUCER (NSAID)

500 CAPLETS**
(*capsule-shaped tablets)

Drug Facts

Active ingredient (in each caplet)
Ibuprofen 200 mg (NSAID)*Pain reliever/Fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - toothache
 - backache
 - the common cold
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away. Cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood-thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Drug Facts (continued under label)

*Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.
**This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., a McNeil Consumer Healthcare Division, owner of the registered trademark Motrin®.

DISTRIBUTED BY: WALGREEN CO. ITEM 7220777 W000000-0000-0
200 WILKOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
Walgreens.com ©2021 Walgreen Co.
MADE IN INDIA 612R 1021

WSORG0122-F REV-1222
Lot No.:
Exp Date:

Varnish Omit Area

IBUPROFEN

ibuprofen tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-9612
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)		IBUPROFEN	200 mg
Inactive Ingredients			
Ingredient Name			Strength

SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	orange	Score	no score
Shape	CAPSULE	Size	10mm
Flavor		Imprint Code	120
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-9612-05	50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/11/2018	
2	NDC:0363-9612-01	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/11/2018	
3	NDC:0363-9612-26	225 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/11/2018	
4	NDC:0363-9612-50	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/11/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091237	12/11/2018	

Labeler - WALGREENS (008965063)

Registrant - TIME CAP LABORATORIES, INC (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		925822975	manufacture(0363-9612)