

**ADVIL- ibuprofen tablet, coated**  
**Lil' Drug Store Products, Inc.**

-----  
**Advil® Tablets**

***Drug Facts***

**Active ingredient (in each tablet)**

Ibuprofen 200 mg (NSAID)<sup>1</sup>

---

<sup>1</sup> nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever/Fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - toothache
  - backache
  - menstrual cramps
  - the common cold
  - muscular aches
  - minor pain of arthritis
- temporarily reduces fever

**Warnings**

**Allergy alert**

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

#### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

#### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma
- you are taking a diuretic

#### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

#### **When using this product**

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

#### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### **Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

**Other information**

- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

**Inactive ingredients**

acetylated monoglycerides, colloidal silicon dioxide, corn starch, croscarmellose sodium, methylparaben, microcrystalline cellulose, pharmaceutical glaze, pharmaceutical ink, povidone, pregelatinized starch, propylparaben, sodium benzoate, sodium lauryl sulfate, stearic acid, sucrose, synthetic iron oxide, titanium dioxide, white wax

**Questions or comments?**

call toll free **1-800-88-ADVIL**

**Repackaged and distributed with permission of Pfizer Consumer Healthcare by:**  
Lil' Drug Store Products, Inc., 1201 Continental Place NE, Cedar Rapids, IA 52402

**PRINCIPAL DISPLAY PANEL - 200 mg Tablet Carton**

**Advil®**

Ibuprofen Tablets, 200 mg

Pain Reliever / Fever Reducer (NSAID)

*Tablets*

**4**

**COATED**

**TABLETS**

**Lil'**

**Drug Store®**

# Advil®

Ibuprofen Tablets, 200 mg  
Pain Reliever / Fever Reducer (NSAID)



Tablets

4 COATED  
TABLETS



Ibuprofen Tablets, 200 mg  
Pain Reliever / Fever Reducer  
(NSAID)  
4 COATED  
TABLETS  
Advil®  
Ibuprofen Tablets, 200 mg  
Pain Reliever / Fever Reducer  
(NSAID)

READ AND KEEP CARTON FOR COMPLETE  
WARNINGS AND INFORMATION

### Drug Facts

#### Active ingredient (in each tablet)

Ibuprofen 200 mg (NSAID)\* ..... Pain reliever/Fever reducer  
\*nonsteroidal anti-inflammatory drug

#### Purpose

#### Uses

- temporarily relieves minor aches and pains due to:
  - headache
  - menstrual cramps
  - minor pain of arthritis
  - temporarily reduces fever
  - toothache
  - the common cold
  - backache
  - muscular aches

#### Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
  - shock
  - facial swelling
  - skin reddening
  - asthma (wheezing)
  - rash
  - blisters
- If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed
- have had stomach ulcers or bleeding problems

#### Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

LIFT HERE for More Drug Facts

Ibuprofen Tablets, 200 mg  
Pain Reliever / Fever Reducer  
(NSAID)  
4 COATED  
TABLETS  
Advil®  
Ibuprofen Tablets, 200 mg  
Pain Reliever / Fever Reducer  
(NSAID)

### Drug Facts (continued)

#### Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

#### Other information

- read all warnings and directions before use. Keep carton.
- store at 20-25°C (68-77°F) ■ avoid excessive heat above 40°C (104°F)

#### Inactive ingredients

acetylated monoglycerides, colloidal silicon dioxide, corn starch, croscarmellose sodium, methylparaben, microcrystalline cellulose, pharmaceutical glaze, pharmaceutical ink, povidone, pregelatinized starch, propylparaben, sodium benzoate, sodium lauryl sulfate, stearic acid, sucrose, synthetic iron oxide, titanium dioxide, white wax

Questions or comments? call toll free 1-800-88-ADVIL

Do Not Use if the sealed child resistant pouch is broken or torn.

For most recent product information, visit [www.Advil.com](http://www.Advil.com)



Pfizer Consumer Healthcare, Madison, NJ 07940 USA © 2011 Pfizer Inc.

4 COATED  
TABLETS  
Advil®  
Ibuprofen Tablets, 200 mg  
Pain Reliever / Fever  
(NSAID)

### Drug Facts (continued)

#### Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma
- you are taking a diuretic

#### Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

#### When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

#### Stop use and ask a doctor if


- you experience any of the following signs of stomach bleeding:
  - feel faint
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear
- vomit blood

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in

Repackaged and distributed with permission of Pfizer Consumer Healthcare by:  
 Lil' Drug Store Products, Inc., 1201 Continental Place NE, Cedar Rapids, IA 52402  
 1-877-507-6516 (M-F 8AM-4:30PM CST) www.lildrugstore.com 97002C-US-10-11



unless differently directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.



## ADVIL

ibuprofen tablet, coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:66715-9700(NDC:0573-0154)
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>Ibuprofen</b> (UNII: WK2XYI10QM) (Ibuprofen - UNII:WK2XYI10QM)	Ibuprofen	200 mg

### Inactive Ingredients

Ingredient Name	Strength
diacetylated monoglycerides (UNII: 5Z17386USF)	
silicon dioxide (UNII: ETJ7Z6XBU4)	
starch, corn (UNII: O8232NY3SJ)	
croscarmellose sodium (UNII: M28OL1HH48)	
methylparaben (UNII: A2I8C7HI9T)	
microcrystalline cellulose (UNII: OP1R32D61U)	
shellac (UNII: 46N107B71O)	
povidone, unspecified (UNII: FZ989GH94E)	
propylparaben (UNII: Z8IX2SC1OH)	
sodium benzoate (UNII: OJ245FE5EU)	
sodium lauryl sulfate (UNII: 368GB5141J)	
stearic acid (UNII: 4ELV7Z65AP)	
sucrose (UNII: C151H8M554)	
titanium dioxide (UNII: 15FIX9V2JP)	
white wax (UNII: 7G1J5DA97F)	

### Product Characteristics

<b>Color</b>	BROWN (pinkish brown)	<b>Score</b>	no score
<b>Shape</b>	ROUND (Smooth in texture w/ polish surface)	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	Advil
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-9700-1	1 in 1 CARTON	02/22/2010	
1		2 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:66715-9700-2	2 in 1 CARTON	02/22/2010	
2		2 in 1 POUCH; Type 0: Not a Combination Product		
3	NDC:66715-9700-3	3 in 1 CARTON	02/22/2010	
3		2 in 1 POUCH; Type 0: Not a Combination Product		
4	NDC:66715-9700-8	6 in 1 CARTON	02/22/2010	03/05/2014
4		2 in 1 POUCH; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA018989	02/22/2010	

**Labeler** - Lil' Drug Store Products, Inc. (093103646)

## Establishment

Name	Address	ID/FEI	Business Operations
CCB Packaging Inc.		362374464	REPACK(66715-9700) , RELABEL(66715-9700)

## Establishment

Name	Address	ID/FEI	Business Operations
Pfizer Consumer Healthcare		828831730	LABEL(66715-9700)

Revised: 4/2017

Lil' Drug Store Products, Inc.