

**FORTICEPT UDDER FORTE SANITIZING TEAT DIP- benzethonium chloride,
thymol emulsion
LIDAN, INC.**

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, [click here](#).

Forticept Udder Forte Sanitizing Teat Dip

Active Ingredients

Benzethonium Chloride 0.1%

Thymol 0.1%

Inactive Ingredients

Emollients, botanical extracts, and FD&C color...99.80%

Uses

Teat dip is an aid in reducing the spread of microorganisms which may cause mastitis.

Helps maintain udder hygiene and promote healthy skin

Caution

For external use only * Not for human use

Keep out of reach of children

Directions for use

IMPORTANT: Do not mix FORTICEPT™ UDDER FORTE Teat Dip with water, other teat dips, or any other products. When transferring from this container to any other, make sure that the other container is clean and is the proper container for dipping.

If product in dip cup becomes visibly dirty, discard contents and refill with undiluted product. Do not reuse or return any unused product to the original container.

PRIOR TO MILKING: Wash teats thoroughly right before milking with FORTICEPT™ UDDER WASH solution or pre-milking teat wash to avoid contamination of milk. Teats should be dried with single-service towels. Use proper procedures for udder washing.

POST-MILKING DIPPING: Immediately after each milking, use FORTICEPT™ UDDER FORTE Teat Dip. Submerge each teat full-length into the teat dip cup. Allow to air dry. Do not wipe. Always use fresh, full-strength FORTICEPT™ UDDER FORTE Teat Dip.

Storage and handling

Keep container closed when not in use. Do not store below 32F/0°C or above 95F/35°C for extended periods. In case of spill, flood area with water.

FIRST AID

Eye contact: In case of contact, immediately flush eyes with cool running water.

Remove contact lenses and continue flushing with plenty of water for at least 15 minutes. Get medical attention if irritation persists.

Skin contact: Wash the skin with water and soap. Wash clothing before reuse.

Ingestion: DO NOT induce vomiting unless directed to do so by medical personnel.

If large quantities of this material are swallowed, call a poison control center immediately.

Questions

For questions, please call:

+1(212)709-8133

Forticept 1L



LIDAN, INC 30 WALL STR. 8th FLOOR NEW YORK, NY 10005 USA WWW.FORTICEPT.COM

Active Ingredients

Benzethonium Chloride0.1%
Thymol0.1%

Inactive Ingredients

Emollients, botanical extracts, and FD&C color.....99.80%

Uses

Teat dip is an aid in reducing the spread of microorganisms which may cause mastitis. Helps maintain udder hygiene and promote healthy skin

Caution

For external use only * Not for human use * Keep out of reach of children

Directions for use

IMPORTANT: Do not mix FORTICEPT™ UDDER FORTE Teat Dip with water, other teat dips, or any other products. When transferring from this container to any other, make sure that the other container is clean and is the proper container for dipping.

If product in dip cup becomes visibly dirty, discard contents and refill with undiluted product. Do not reuse or return any unused product to the original container.

PRIOR TO MILKING: Wash teats thoroughly right before milking with FORTICEPT™ UDDER WASH solution or pre-milking teat wash to avoid contamination of milk. Teats should be dried with single-service towels. Use proper procedures for udder washing.

POST-MILKING DIPPING: Immediately after each milking, use FORTICEPT™ UDDER FORTE Teat Dip. Submerge each teat full-length into the teat dip cup. Allow to air dry. Do not wipe. Always use fresh, full-strength FORTICEPT™ UDDER FORTE Teat Dip.

Storage and handling

Keep container closed when not in use. Do not store below 32F/0°C or above 95F/35°C for extended periods. In case of spill, flood area with water.

FIRST AID

Eye contact: In case of contact, immediately flush eyes with cool running water. Remove contact lenses and continue flushing with plenty of water for at least 15 minutes. Get medical attention if irritation persists.

Skin contact: Wash the skin with water and soap. Wash clothing before reuse.

Ingestion: DO NOT induce vomiting unless directed to do so by medical personnel. If large quantities of this material are swallowed, call a poison control center immediately.

For questions, please call:
+1(212)709-8133

MADE IN USA



FORTICEPT™ is registered Trade Mark of Lidan, Inc. all rights reserved 2016

Batch No.:
Exp. Date:

Forticept 3.785L



LIDAN, INC 30 WALL STR. 8th FLOOR NEW YORK, NY 10005 USA WWW.FORTICEPT.COM

Active Ingredients

Benzethonium Chloride0.1%
Thymol0.1%

Inactive Ingredients

Emollients, botanical extracts, and FD&C color.....99.80%

Uses

Teat dip is an aid in reducing the spread of microorganisms which may cause mastitis. Helps maintain udder hygiene and promote healthy skin

Caution

For external use only * Not for human use * Keep out of reach of children

Directions for use

IMPORTANT: Do not mix FORTICEPT™ UDDER FORTE Teat Dip with water, other teat dips, or any other products. When transferring from this container to any other, make sure that the other container is clean and is the proper container for dipping.

If product in dip cup becomes visibly dirty, discard contents and refill with undiluted product. Do not reuse or return any unused product to the original container.

PRIOR TO MILKING: Wash teats thoroughly right before milking with FORTICEPT™ UDDER WASH solution or pre-milking teat wash to avoid contamination of milk. Teats should be dried with single-service towels. Use proper procedures for udder washing.

POST-MILKING DIPPING: Immediately after each milking, use FORTICEPT™ UDDER FORTE Teat Dip. Submerge each teat full-length into the teat dip cup. Allow to air dry. Do not wipe. Always use fresh, full-strength FORTICEPT™ UDDER FORTE Teat Dip.

Storage and handling

Keep container closed when not in use. Do not store below 32F/0°C or above 95F/35°C for extended periods. In case of spill, flood area with water.

FIRST AID

Eye contact: In case of contact, immediately flush eyes with cool running water. Remove contact lenses and continue flushing with plenty of water for at least 15 minutes. Get medical attention if irritation persists.

Skin contact: Wash the skin with water and soap. Wash clothing before reuse.

Ingestion: DO NOT induce vomiting unless directed to do so by medical personnel.

If large quantities of this material are swallowed, call a poison control center immediately.

For questions, please call:

+1(212)709-8133

MADE IN USA



FORTICEPT™ is registered Trade Mark of Lidan, Inc. all rights reserved 2016

Batch No:
Exp. Date:

Forticept 5L



LIDAN, INC 30 WALL STR. 8th FLOOR NEW YORK, NY 10005 USA WWW.FORTICEPT.COM

Active Ingredients

Benzethonium Chloride0.1%
Thymol0.1%

Inactive Ingredients

Emollients, botanical extracts, and FD&C color.....99.80%

Uses

Teat dip is an aid in reducing the spread of microorganisms which may cause mastitis. Helps maintain udder hygiene and promote healthy skin

Caution

For external use only * Not for human use * Keep out of reach of children

Directions for use

IMPORTANT: Do not mix FORTICEPT™ UDDER FORTE Teat Dip with water, other teat dips, or any other products. When transferring from this container to any other, make sure that the other container is clean and is the proper container for dipping.

If product in dip cup becomes visibly dirty, discard contents and refill with undiluted product. Do not reuse or return any unused product to the original container.

PRIOR TO MILKING: Wash teats thoroughly right before milking with FORTICEPT™ UDDER WASH solution or pre-milking teat wash to avoid contamination of milk. Teats should be dried with single-service towels. Use proper procedures for udder washing.

POST-MILKING DIPPING: Immediately after each milking, use FORTICEPT™ UDDER FORTE Teat Dip. Submerge each teat full-length into the teat dip cup. Allow to air dry. Do not wipe. Always use fresh, full-strength FORTICEPT™ UDDER FORTE Teat Dip.

Storage and handling

Keep container closed when not in use. Do not store below 32F/0°C or above 95F/35°C for extended periods. In case of spill, flood area with water.

FIRST AID

Eye contact: In case of contact, immediately flush eyes with cool running water. Remove contact lenses and continue flushing with plenty of water for at least 15 minutes. Get medical attention if irritation persists.

Skin contact: Wash the skin with water and soap. Wash clothing before reuse.

Ingestion: DO NOT induce vomiting unless directed to do so by medical personnel.

If large quantities of this material are swallowed, call a poison control center immediately.

For questions, please call:

+1(212)709-8133

MADE IN USA



FORTICEPT™ is registered Trade Mark of Lidan, Inc. all rights reserved 2016

Batch No:
Exp. Date:

Forticept 10L



forticept™

FORTICEPT™

**UDDER FORTE
SANITIZING TEAT DIP**

IODINE FREE, PHOSPHORUS FREE
EMOLLIENT SYSTEM

With natural extracts of
Chamomile & Yarrow

*Teat dip is an aid in
reducing the spread of microorganisms
which may cause mastitis*

For Veterinary Use Only

2.64 Gal. / 10L

LIDAN, INC 30 WALL STR. 8th FLOOR NEW YORK, NY 10005 USA WWW.FORTICEPT.COM

Active Ingredients

Benzethonium Chloride0.1%
Thymol0.1%

Inactive Ingredients

Emollients, botanical extracts, and FD&C color.....99.80%

Uses

Teat dip is an aid in reducing the spread of microorganisms which may cause mastitis.
Helps maintain udder hygiene and promote healthy skin

Caution

For external use only * Not for human use * Keep out of reach of children

Directions for use

IMPORTANT: Do not mix FORTICEPT™ UDDER FORTE Teat Dip with water, other teat dips, or any other products. When transferring from this container to any other, make sure that the other container is clean and is the proper container for dipping.

If product in dip cup becomes visibly dirty, discard contents and refill with undiluted product. Do not reuse or return any unused product to the original container.

PRIOR TO MILKING: Wash teats thoroughly right before milking with FORTICEPT™ UDDER WASH solution or pre-milking teat wash to avoid contamination of milk. Teats should be dried with single-service towels. Use proper procedures for udder washing.

POST-MILKING DIPPING: Immediately after each milking, use FORTICEPT™ UDDER FORTE Teat Dip. Submerge each teat full-length into the teat dip cup. Allow to air dry. Do not wipe. Always use fresh, full-strength FORTICEPT™ UDDER FORTE Teat Dip.

Storage and handling

Keep container closed when not in use. Do not store below 32F/0°C or above 95F/35°C for extended periods. In case of spill, flood area with water.

FIRST AID

Eye contact: In case of contact, immediately flush eyes with cool running water. Remove contact lenses and continue flushing with plenty of water for at least 15 minutes. Get medical attention if irritation persists.

Skin contact: Wash the skin with water and soap. Wash clothing before reuse.

Ingestion: DO NOT induce vomiting unless directed to do so by medical personnel.

If large quantities of this material are swallowed, call a poison control center immediately.

For questions, please call:
+1(212)709-8133

MADE IN USA



FORTICEPT™ is registered Trade Mark of Lidan, Inc. all rights reserved 2016

Batch No.:
Exp. Date:

Forticept 20L



forticept™

FORTICEPT™

**UDDER FORTE
SANITIZING TEAT DIP**

IODINE FREE, PHOSPHORUS FREE
EMOLLIENT SYSTEM

With natural extracts of
Chamomile & Yarrow

*Teat dip is an aid in
reducing the spread of microorganisms
which may cause mastitis*

For Veterinary Use Only

5 Gal. / 20L

LIDAN, INC 30 WALL STR. 8th FLOOR NEW YORK, NY 10005 USA WWW.FORTICEPT.COM

Active Ingredients

Benzethonium Chloride0.1%
Thymol0.1%

Inactive Ingredients

Emollients, botanical extracts, and FD&C color.....99.80%

Uses

Teat dip is an aid in reducing the spread of microorganisms which may cause mastitis.
Helps maintain udder hygiene and promote healthy skin

Caution

For external use only * Not for human use * Keep out of reach of children

Directions for use

IMPORTANT: Do not mix FORTICEPT™ UDDER FORTE Teat Dip with water, other teat dips, or any other products. When transferring from this container to any other, make sure that the other container is clean and is the proper container for dipping.

If product in dip cup becomes visibly dirty, discard contents and refill with undiluted product. Do not reuse or return any unused product to the original container.

PRIOR TO MILKING: Wash teats thoroughly right before milking with FORTICEPT™ UDDER WASH solution or pre-milking teat wash to avoid contamination of milk. Teats should be dried with single-service towels. Use proper procedures for udder washing.

POST-MILKING DIPPING: Immediately after each milking, use FORTICEPT™ UDDER FORTE Teat Dip. Submerge each teat full-length into the teat dip cup. Allow to air dry. Do not wipe. Always use fresh, full-strength FORTICEPT™ UDDER FORTE Teat Dip.

Storage and handling

Keep container closed when not in use. Do not store below 32F/0°C or above 95F/35°C for extended periods. In case of spill, flood area with water.

FIRST AID

Eye contact: In case of contact, immediately flush eyes with cool running water. Remove contact lenses and continue flushing with plenty of water for at least 15 minutes. Get medical attention if irritation persists.

Skin contact: Wash the skin with water and soap. Wash clothing before reuse.

Ingestion: DO NOT induce vomiting unless directed to do so by medical personnel.

If large quantities of this material are swallowed, call a poison control center immediately.

For questions, please call:
+1(212)709-8133

MADE IN USA



FORTICEPT™ is registered Trade Mark of Lidan, Inc. all rights reserved 2016

Batch No.:
Exp. Date:

Forticept 1000L



LIDAN, INC 30 WALL STR. 8th FLOOR NEW YORK, NY 10005 USA WWW.FORTICEPT.COM

Active Ingredients

Benzethonium Chloride0.1%
Thymol0.1%

Inactive Ingredients

Emollients, botanical extracts, and FD&C color.....99.80%

Uses

Teat dip is an aid in reducing the spread of microorganisms which may cause mastitis.
Helps maintain udder hygiene and promote healthy skin

Caution

For external use only * Not for human use * Keep out of reach of children

Directions for use

IMPORTANT: Do not mix FORTICEPT™ UDDER FORTE Teat Dip with water, other teat dips, or any other products. When transferring from this container to any other, make sure that the other container is clean and is the proper container for dipping.

If product in dip cup becomes visibly dirty, discard contents and refill with undiluted product. Do not reuse or return any unused product to the original container.

PRIOR TO MILKING: Wash teats thoroughly right before milking with FORTICEPT™ UDDER WASH solution or pre-milking teat wash to avoid contamination of milk. Teats should be dried with single-service towels. Use proper procedures for udder washing.

POST-MILKING DIPPING: Immediately after each milking, use FORTICEPT™ UDDER FORTE Teat Dip. Submerge each teat full-length into the teat dip cup. Allow to air dry. Do not wipe. Always use fresh, full-strength FORTICEPT™ UDDER FORTE Teat Dip.

Storage and handling

Keep container closed when not in use. Do not store below 32F/0°C or above 95F/35°C for extended periods. In case of spill, flood area with water.

FIRST AID

Eye contact: In case of contact, immediately flush eyes with cool running water. Remove contact lenses and continue flushing with plenty of water for at least 15 minutes. Get medical attention if irritation persists.

Skin contact: Wash the skin with water and soap. Wash clothing before reuse.

Ingestion: DO NOT induce vomiting unless directed to do so by medical personnel. If large quantities of this material are swallowed, call a poison control center immediately.

For questions, please call:
+1(212)709-8133

MADE IN USA



FORTICEPT™ is registered Trade Mark of Lidan, Inc. all rights reserved 2016

Batch No:
Exp. Date:

FORTICEPT UDDER FORTE SANITIZING TEAT DIP

benzethonium chloride, thymol emulsion

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:72127-0209
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZETHONIUM CHLORIDE (UNII: PH41D05744) (BENZETHONIUM - UNII:1VU15B70BP)	BENZETHONIUM CHLORIDE	1 g in 1 L
THYMOL (UNII: 3J50XA376E) (THYMOL - UNII:3J50XA376E)	THYMOL	1 g in 1 L

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72127-0209-0	1 L in 1 BOTTLE		
2	NDC:72127-0209-1	3.785 L in 1 BOTTLE		
3	NDC:72127-0209-2	5 L in 1 BOTTLE		
4	NDC:72127-0209-3	10 L in 1 BOTTLE		
5	NDC:72127-0209-4	20 L in 1 BOTTLE		
6	NDC:72127-0209-5	1000 L in 1 BOTTLE		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		03/15/2018	

Labeler - LIDAN, INC. (014456716)

Registrant - LIDAN, INC. (014456716)

Establishment			
Name	Address	ID/FEI	Business Operations
Cosco International, Inc.		016433141	analysis, manufacture, pack

Establishment			
Name	Address	ID/FEI	Business Operations
DISHMAN CARBOGEN AMCIS LIMITED		915628142	api manufacture

Establishment			
Name	Address	ID/FEI	Business Operations
Univar USA Inc.		042953927	api manufacture

Establishment			
Name	Address	ID/FEI	Business Operations
Pharmex Group LLC		365309015	pack, repack

Revised: 4/2018

LIDAN, INC.