

SODIUM SULFACETAMIDE- sodium sulfacetamide liquid
Acella Pharmaceuticals, LLC


Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Sodium Sulfacetamide 10% Wash

Rx Only

DESCRIPTION:

Sodium sulfacetamide is $C_8H_9N_2NaO_3 \cdot H_2O$ with molecular weight of 254.24. Chemically, it is Acetamide N-[(4-aminophenyl) sulfonyl]-acetamide, monosodium salt, monohydrate. The structural formula is:



Each gram of Sodium Sulfacetamide 10% Wash contains 100 mg of Sodium Sulfacetamide, USP in a formulation containing cocamidopropyl betaine, disodium EDTA, PEG-150 pentaerythrityl tetrastearate and PEG 6 caprylic/capric glycerides, glyceryl monostearate, methyl paraben, PEG-60 almond triglyceride, polysorbate 60, purified water, sodium lauryl sulfate and sodium thiosulfate.

Sodium Sulfacetamide is an odorless, white, crystalline powder with a bitter taste. It is freely soluble in water, sparingly soluble in alcohol, while practically insoluble in benzene, in chloroform and in ether.

CLINICAL PHARMACOLOGY:

Sodium sulfacetamide exerts a bacteriostatic effect against sulfonamide sensitive Gram-positive and Gram-negative microorganisms commonly isolated from secondary cutaneous pyogenic infections. It acts by restricting the synthesis of folic acid required by bacteria for growth, by its competition with para-aminobenzoic acid. There is no clinical data available on the degree and rate of systemic absorption of Sodium Sulfacetamide 10% Wash when applied to the skin or scalp. However, significant absorption of sodium sulfacetamide through the skin has been reported.

The following in vitro data is available but the clinical significance is not known. Organisms which show susceptibility to sodium sulfacetamide are: *Streptococci*, *Staphylococci*, *E. coli*, *Klebsiella pneumonia*, *Pseudomonas pyocyanea*, *Salmonella species*, *Proteus vulgaris*, *Nocardia* and *Actinomyces* .

INDICATIONS:

Sodium Sulfacetamide 10% Wash is intended for topical application in the following scaling dermatoses: seborrheic dermatitis and seborrhea sicca (dandruff). It is also indicated for the treatment of secondary bacterial infections of the skin due to organisms susceptible to sulfonamides.

CONTRAINDICATIONS:

Sodium Sulfacetamide 10% Wash is contraindicated for use by patients having known hypersensitivity to sulfonamides or any other component of this product.

WARNINGS:

Sulfonamides are known to cause Stevens-Johnson syndrome in hypersensitive individuals. Stevens-Johnson syndrome also has been reported following the use of sodium sulfacetamide topically. Cases of drug induced systemic lupus erythematosus from topical sulfacetamide also have been reported. In one of these cases, there was a fatal outcome. **KEEP OUT OF THE REACH OF CHILDREN.**

PRECAUTIONS:

For external use only.

General - Nonsusceptible organisms, including fungi, may proliferate with the use of this preparation. Hypersensitivity reactions may recur when a sulfonamide is readministered, irrespective of the route of administration, and cross hypersensitivity between different sulfonamides may occur. If Sodium Sulfacetamide 10% Wash produces signs of hypersensitivity or other untoward reactions, discontinue use of the preparation. Systemic absorption of topical sulfonamides is greater following application to large, infected, abraded, denuded, or severely burned areas. Under these circumstances, potentially any of the adverse effects produced by the systemic administration of these agents could occur and appropriate observations and laboratory determinations should be performed.

Information for patients - Patients should discontinue use of Sodium Sulfacetamide 10% Wash if the condition worsens, or if a rash develops in the area being treated or elsewhere. Sodium Sulfacetamide 10% Wash also should be discontinued promptly and the physician notified if any arthritis, fever or sores in the mouth develop.

Drug Interactions:

Sodium Sulfacetamide 10% Wash is incompatible with silver preparations.

Carcinogenesis, Mutagenesis and Impairment of Fertility - Long-term studies in animals have not been performed to evaluate carcinogenic potential. Studies on reproduction and fertility also have not been performed. Chromosomal nondisjunction in the yeast, *Saccharomyces cerevisiae*, following application of sodium sulfacetamide has been reported. The significance of this finding to the topical use of sodium sulfacetamide in the human is not known.

PREGNANCY:

Category C. Animal reproduction studies have not been conducted with Sodium Sulfacetamide 10% Wash. It is not known whether Sodium Sulfacetamide 10% Wash can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Sodium Sulfacetamide 10% Wash should be given to a pregnant woman only if clearly needed or when potential benefits outweigh potential hazards to the fetus.

NURSING MOTHERS:

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be when Sodium Sulfacetamide 10% Wash is administered to a nursing woman.

PEDIATRIC USE:

Safety and effectiveness in children under the age of 12 have not been established.

ADVERSE REACTIONS:

Reports of irritation and hypersensitivity to sodium sulfacetamide are uncommon. The following adverse reactions, reported after administration of sterile ophthalmic sodium sulfacetamide, are noteworthy: instances of Stevens-Johnson syndrome and instances of local hypersensitivity which progressed to a syndrome resembling systemic lupus erythematosus; in one case a fatal outcome was reported. (See **warnings**.)

OVERDOSAGE:

The oral LD₅₀ of sulfacetamide in mice is 16.5 g/kg. In the event of overdosage, emergency treatment should be started immediately. Manifestations: Overdosage may cause nausea and vomiting. Large oral overdosage may cause hematuria, crystalluria and renal shutdown due to the precipitation of sulfa crystals in the renal tubules and the urinary tract. For treatment, contact your local Poison Control Center.

DOSAGE AND ADMINISTRATION:

Seborrheic dermatitis including seborrhea sicca - Wash affected areas twice daily (morning and evening), or as directed by your physician. Avoid contact with eyes or mucous membranes. Wet skin and liberally apply to areas to be cleansed, massage gently into skin working into a full lather, rinse thoroughly and pat dry. Rinsing with plain water will remove any excess medication. Repeat application as described for eight to ten days. If skin dryness occurs it may be controlled by rinsing cleanser off sooner or using less frequently. Regular shampooing following Sodium Sulfacetamide 10% Wash is not necessary, but the hair should be shampooed at least once a week. As the condition subsides, the interval between applications may be lengthened. Applications once or twice weekly or every other week may prevent recurrence. Should the condition recur

after stopping therapy, the application of Sodium Sulfacetamide 10% Wash should be reinstated as at the beginning of treatment.

Secondary Cutaneous Bacterial Infections - Wet skin and liberally apply to areas to be cleansed, massage gently into skin working into a full lather, rinse thoroughly and pat dry. Rinsing with plain water will remove any excess medication. Repeat application as described for eight to ten days. If skin dryness occurs it may be controlled by rinsing cleanser off sooner or using less often.

HOW SUPPLIED:

Sodium Sulfacetamide 10% Wash is available in 16 fluid ounce bottles, NDC 42192-129-16.

Store at controlled room temperature, 15° - 30°C (59° - 86°F). Do not freeze.

Note: Store upright. Protect from freezing and excessive heat. The wash may tend to darken slightly on storage. Slight discoloration does not impair the efficacy or safety of the product. Occasionally, a slight yellowish discoloration may occur when an excessive amount of the wash is used and comes in contact with white fabrics. This discoloration, however, presents no problem, as it is readily removed by ordinary laundering without bleaches.

All prescription substitutions using this product shall be made subject to state and federal statutes as applicable. **NOTE: this is not an Orange Book product and has not been subjected to FDA therapeutic equivalency or other equivalency testing. No representation is made as to generic status or bioequivalency.** Each person recommending a prescription substitution using this product shall make such recommendations based on each such person's professional opinion and knowledge, upon evaluating the active ingredients, excipients, inactive ingredients and chemical formulation information provided herein.

MANUFACTURED FOR: Acella Pharmaceuticals, LLC
Alpharetta, GA 30009
1-800-541-4802
Rev. 1010v2

PACKAGE LABEL - 10% Wash

NDC 42192-129-16

Sodium Sulfacetamide
10% Wash

Rx Only

For External Use Only

16 fl. oz. (480 mL)

Acella
PHARMACEUTICALS, LLC

NDC 42192-129-16

Sodium Sulfacetamide 10% Wash

Rx Only

For External Use Only

16 fl. oz. (480 mL)



Directions: Wash affected areas twice daily (morning and evening) or as directed by your physician. Rinse thoroughly and pat dry. See package insert for complete information.

Warnings: For external use only. Avoid contact with eyes.

KEEP OUT OF REACH OF CHILDREN. Keep bottle tightly closed.

Ingredients: Each mL of Sodium Sulfacetamide 10% Wash contains 100 mg of sodium sulfacetamide, USP in a formulation containing cocamidopropyl betaine, disodium EDTA, PEG-150 pentaerythrityl tetrastearate and PEG 6 caprylic/capric glycerides, glyceryl monostearate, methyl paraben, PEG-60 almond triglyceride, polysorbate 60, purified water, sodium lauryl sulfate and sodium thiosulfate.

Store at controlled room temperature 15° - 30°C (59° - 86°F). Protect from freezing.

Sodium Sulfacetamide 10% Wash is supplied in a 16 fluid ounce (480 mL) bottle (NDC 42192-129-16). For lot number and expiration date, see bottom of bottle.

All prescription substitutions and/or recommendations using this product shall be made subject to state and federal statutes as applicable. **NOTE: This is not an Orange Book product. No representation is made as to generic status or bioequivalency.** Please see package insert for more information.

Manufactured for:
Acella Pharmaceuticals, LLC
Alpharetta, GA 30009
1-800-541-4802
Rev. 1010d3



SODIUM SULFACETAMIDE

sodium sulfacetamide liquid

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:42192-129
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SULFACETAMIDE SODIUM (UNII: 4NRT660KJQ) (SULFACETAMIDE - UNII:4965G3J0F5)	SULFACETAMIDE SODIUM	100 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
COCAMIDOPROPYL BETAINE (UNII: 5OCF3011KX)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
PENTAERYTHRITYL TETRASTEARATE (UNII: W9Q3DZS0EG)	
CAPRYLIC/CAPRIC MONO/DIGLYCERIDES (UNII: U72Q2I8C85)	

GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PEG-60 ALMOND GLYCERIDES (UNII: 4Y0E651N0F)	
POLYSORBATE 60 (UNII: CAL22UVI4M)	
WATER (UNII: 059QF0KO0R)	
SODIUM LAURETH SULFATE (UNII: BPV390UAP0)	
SODIUM THIOSULFATE (UNII: HX1032V43M)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42192-129-16	480 mL in 1 BOTTLE; Type 0: Not a Combination Product	01/04/2011	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		01/04/2011	

Labeler - Acella Pharmaceuticals, LLC (825380939)

Registrant - Acella Pharmaceuticals, LLC (825380939)

Establishment

Name	Address	ID/FEI	Business Operations
Acella Pharmaceuticals, LLC		825380939	manufacture(42192-129)

Revised: 1/2024

Acella Pharmaceuticals, LLC