

BISACODYL ENTERIC COATED- bisacodyl tablet, coated
CVS Pharmacy

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient (in each tablet)

Bisacodyl USP 5 mg

Purpose

Stimulant laxative

Use

- for temporary relief of occasional constipation and irregularity
- this product generally produces a bowel movement in 6 to 12 hours

Warnings

Do not use

if you cannot swallow without chewing.

Ask a doctor before use if you have

- stomach pain, nausea or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

When using this product

- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- it may cause stomach discomfort, faintness and cramps

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take with a glass of water

adults and children 12 years of age and over | take 1 to 3 tablets in a single daily dose

children 6 to under 12 years of age	take 1 tablet in a single daily dose
children under 6 years of age	ask a doctor

Other information

- store between 20°-25°C (68°-77°F)

Inactive ingredients

acacia, anhydrous calcium sulfate, anhydrous lactose, carnauba wax, colloidal silicon dioxide, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, gelatin, iron oxide, iron oxide black, iron oxide yellow, magnesium stearate, microcrystalline cellulose, polyethylene glycol (PEG) 400, polyvinyl acetate phthalate, povidone, shellac, sodium starch glycolate, stearic acid, sugar, talc, titanium dioxide

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to the active ingredient in DULCOLAX®*

Gentle

LAXATIVE

BISACODYL, 5 mg

- Guaranteed relief
- Gentle yet effective

Tablets

Enteric coated

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.


*This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals Inc., distributor of Dulcolax® Laxative.

Distributed by: CVS Pharmacy, Inc.

One CVS Drive, Woonsocket, RI 02895

CVS.COM® 1-800-SHOP CVS

Product Label



pharmacy™

Gentle LAXATIVE

BISACODYL, 5 mg

CVS
pharmacy™

NDC 59779-990-25

Compare to the active ingredient in Dulcolax®*

Gentle LAXATIVE

BISACODYL, 5 mg

- **Guaranteed relief**
- **Gentle yet effective**

25 Tablets
Enteric coated



TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Lot No.:
Exp. Date:

*This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals, Inc., distributor of Dulcolax® Laxative.

Distributed by: **CVS Pharmacy, Inc.**
One CVS Drive, Woonsocket, RI 02895
© 2014 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-14921



PLEASE RECYCLE

#348639



PLD-C
FC002044

CVS Quality
Money Back Guarantee

<p>Warnings</p> <p>Do not use if you cannot swallow without chewing.</p> <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ■ stomach pain, nausea or vomiting ■ a sudden change in bowel habits that lasts more than 2 weeks <p>When using this product</p> <ul style="list-style-type: none"> ■ do not chew or crush tablet(s) ■ do not use within 1 hour after taking an antacid or milk ■ it may cause stomach discomfort, faintness and cramps <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition. ■ you need to use a laxative for more than 1 week <p>If pregnant or breast-feeding, ask a health professional before use.</p>	
<p>Other information</p> <ul style="list-style-type: none"> ■ store between 20°-25°C (68°-77°F) 	
<p>Inactive ingredients acacia, anhydrous calcium sulfate, anhydrous lactose, carnauba wax, colloidal silicon dioxide, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, gelatin, iron oxide, iron oxide black, iron oxide yellow, magnesium stearate, microcrystalline cellulose, polyethylene glycol (PEG) 400, polyvinyl acetate phthalate, povidone, shellac, sodium starch glycolate, stearic acid, sugar, talc, titanium dioxide</p>	<p>Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST</p>
<p>children 6 to under 12 take 1 tablet in a single daily dose years of age</p>	<p>children under 6 years ask a doctor of age</p>

Drug Facts		Use <ul style="list-style-type: none"> for temporary relief of occasional constipation and irregularity this product generally produces a bowel movement in 6 to 12 hours
Active ingredient (in each tablet) Bisacodyl USP 5 mg.....Stimulant laxative		
Purpose Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.		Directions <ul style="list-style-type: none"> take with a glass of water adults and children 12 years of age and over take 1 to 3 tablets in a single daily dose
Drug Facts (continued)		

CVS Enteric Coated Laxative Tablets

BISACODYL ENTERIC COATED
bisacodyl tablet, coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:59779-990
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
BISACODYL (UNII: 10 X0 709 Y6 I) (BISACODYL - UNII:10 X0 709 Y6 I)	BISACODYL	5 mg

Inactive Ingredients	
Ingredient Name	Strength
ACACIA (UNII: 5C5403N26O)	
CALCIUM SULFATE ANHYDROUS (UNII: E934B3V59H)	
ANHYDROUS LACTOSE (UNII: 3S Y5LH9 PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SHELLAC (UNII: 46N107B710)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
Polyvinyl Acetate Phthalate (UNII: 58QVG85GW3)	
SUCROSE (UNII: C151H8M554)	
POVIDONES (UNII: FZ989GH94E)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	

POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)

ALUMINUM OXIDE (UNII: LM26O6933)

Product Characteristics

Color	ORANGE	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	TCL003
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59779-990-50	2 in 1 CARTON		
1		25 in 1 BLISTER PACK		
2	NDC:59779-990-10	10 in 1 CARTON		
2		1 in 1 BLISTER PACK		
3	NDC:59779-990-25	25 in 1 CARTON		
3		1 in 1 BLISTER PACK		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part334	01/10/2013	

Labeler - CVS Pharmacy (062312574)

Revised: 5/2014

CVS Pharmacy