

HYDROMORPHONE HYDROCHLORIDE- hydromorphone hydrochloride tablet, extended release

Paddock Laboratories, LLC

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use HYDROMORPHONE HYDROCHLORIDE EXTENDED-RELEASE TABLETS safely and effectively. See full prescribing information for HYDROMORPHONE HYDROCHLORIDE EXTENDED-RELEASE TABLETS.

HYDROMORPHONE HYDROCHLORIDE extended-release tablets, for oral use, CII

Initial U.S. Approval: 1984

WARNING: ADDICTION, ABUSE, AND MISUSE; RISK EVALUATION AND MITIGATION STRATEGY (REMS); LIFE-THREATENING RESPIRATORY DEPRESSION; ACCIDENTAL INGESTION; NEONATAL OPIOID WITHDRAWAL SYNDROME; and RISKS FROM CONCOMITANT USE WITH BENZODIAZEPINES OR OTHER CNS DEPRESSANTS

See full prescribing information for complete boxed warning.

- Hydromorphone hydrochloride extended-release tablets expose users to risks of addiction, abuse, and misuse, which can lead to overdose and death. Assess patient's risk before prescribing, and monitor regularly for these behaviors and conditions (5.1).
- To ensure that the benefits of opioid analgesics outweigh the risks of addiction, abuse, and misuse, the Food and Drug Administration (FDA) has required a Risk Evaluation and Mitigation Strategy (REMS) for these products. (5.2)
- Serious, life-threatening, or fatal respiratory depression may occur. Monitor closely, especially upon initiation or following a dose increase. Instruct patients to swallow hydromorphone hydrochloride extended-release tablets whole to avoid exposure to a potentially fatal dose of hydromorphone (5.2).
- Accidental ingestion of hydromorphone hydrochloride extended-release tablets, especially by children, can result in fatal overdose of hydromorphone (5.2).
- Prolonged use of hydromorphone hydrochloride extended-release tablets during pregnancy can result in neonatal opioid withdrawal syndrome, which may be life-threatening if not recognized and treated. If opioid use is required for a prolonged period in a pregnant woman, advise the patient of the risk of neonatal opioid withdrawal syndrome and ensure that appropriate treatment will be available (5.3).
- Concomitant use of opioids with benzodiazepines or other central nervous system (CNS) depressants, including alcohol, may result in profound sedation, respiratory depression, coma, and death. Reserve concomitant prescribing for use in patients for whom alternative treatment options are inadequate; limit dosages and durations to the minimum required; and follow patients for signs and symptoms of respiratory depression and sedation (5.4, 7).

----- **RECENT MAJOR CHANGES** -----

Boxed Warning 09/2018

Warnings and Precautions (5.2) 09/2018

----- **INDICATIONS AND USAGE** -----

Hydromorphone hydrochloride extended-release tablets are an opioid agonist indicated in opioid-tolerant patients for the management of pain severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate. (1)

Patients considered opioid tolerant are those who are taking, for one week or longer, at least 60 mg oral morphine per day, 25 mcg transdermal fentanyl per hour, 30 mg oral oxycodone per day, 8 mg oral hydromorphone per day, 25 mg oral oxymorphone per day, 60 mg oral hydrocodone per day, or an equianalgesic dose of another opioid. (1)

Limitations of Use (1)

- Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses, and because of the greater risks of overdose and death with extended-release opioid formulations, reserve hydromorphone hydrochloride extended-release tablets for use in patients for whom alternative treatment options (e.g., non-opioid analgesics or immediate-release opioids) are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain. (1)

- Hydromorphone hydrochloride extended-release tablets are not indicated as an as-needed (prn) analgesic. (1)

----- DOSAGE AND ADMINISTRATION -----

- To be prescribed only by healthcare providers knowledgeable in use of potent opioids for management of chronic pain.
- For once daily administration IN OPIOID-TOLERANT PATIENTS (2.1).
- Use the lowest effective dosage for the shortest duration consistent with individual patient treatment goals (2.1).
- Individualize dosing based on the severity of pain, patients response, prior analgesic experience, and risk factors for addiction, abuse and misuse (2.3).
- Instruct patients to swallow hydromorphone hydrochloride extended-release tablets intact, and not to cut, break, chew, crush, or dissolve the tablets (risk of potentially fatal overdose) (2.1, 5.1).
- Dose may be increased using increments of 4 to 8 mg every 3 to 4 days as needed to achieve adequate analgesia (2.3).
- Do not abruptly discontinue hydromorphone hydrochloride extended-release tablets in a physically-dependent patient (2.4, 5.12).
- Moderate Hepatic Impairment: Initiate treatment with 25% of the dose that would be prescribed for patients with normal hepatic function. Monitor closely for respiratory and central nervous system depression (2.5).
- Moderate and Severe Renal Impairment: Initiate treatment in patients with moderate renal impairment with 50% and patients with severe renal impairment with 25% of the hydromorphone hydrochloride extended-release tablets dose that would be prescribed for patients with normal renal function. Monitor closely for respiratory and central nervous system depression (2.6).

----- DOSAGE FORMS AND STRENGTHS -----

Extended-release tablets: 8 mg, 12 mg, 16 mg, 32 mg (3)

----- CONTRAINDICATIONS -----

- Opioid non-tolerant patients (4)
- Significant respiratory depression (4)
- Acute or severe bronchial asthma in an unmonitored setting or in absence of resuscitative equipment (4)
- Known or suspected gastrointestinal obstruction, including paralytic ileus (4)
- Narrowed or obstructed gastrointestinal tract (4)
- Known hypersensitivity to any components including hydromorphone hydrochloride and sulfites (4, 5.13)

----- WARNINGS AND PRECAUTIONS -----

- Life-Threatening Respiratory Depression in Patients with Chronic Pulmonary Disease or Elderly Cachectic Debilitated Patients: Monitor closely, particularly during initiation and titration (5.6).
- Adrenal Insufficiency: If diagnosed, treat with physiologic replacement of corticosteroids, and wean patient off of the opioid (5.7).
- Severe Hypotension: Monitor during dose initiation and titration. Avoid use of hydromorphone hydrochloride extended-release tablets in patients with circulatory shock (5.8).
- Risks of Use in Patients with Increased Intracranial Pressure, Brain Tumors, Head Injury, or Impaired Consciousness: Monitor for sedation and respiratory depression. Avoid use of hydromorphone hydrochloride extended-release tablets in patients with impaired consciousness or coma (5.9).

----- ADVERSE REACTIONS -----

Most common adverse reactions (incidence >10%) are: constipation, nausea, vomiting, somnolence, headache, and dizziness (6.1).

To report SUSPECTED ADVERSE REACTIONS, contact Perrigo at 1-866-634-9120 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

----- DRUG INTERACTIONS -----

- Serotonergic Drugs: Concomitant use may result in serotonin syndrome. Discontinue hydromorphone hydrochloride extended-release tablets if serotonin syndrome is suspected (7).
- Monoamine Oxidase Inhibitors (MAOIs): Can potentiate the effects of hydromorphone. Avoid concomitant use in patients receiving MAOIs or within 14 days of stopping treatment with an MAOI (7).
- Mixed agonist/antagonist and partial agonist opioid analgesics: Avoid use with hydromorphone hydrochloride extended-release tablets because they may reduce analgesic effect of hydromorphone hydrochloride extended-release tablets or precipitate withdrawal symptoms (5.12, 7).

----- USE IN SPECIFIC POPULATIONS -----

- Pregnancy: May cause fetal harm (8.1).
- Lactation: Not recommended (8.2).
- Severe Hepatic Impairment: Use not recommended (8.6).
- Severe Renal Impairment: Consider an alternate analgesic (8.7).

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.

Revised: 1/2019

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: ADDICTION, ABUSE, AND MISUSE; RISK EVALUATION AND MITIGATION STRATEGY (REMS); LIFE-THREATENING RESPIRATORY DEPRESSION; ACCIDENTAL INGESTION; NEONATAL OPIOID WITHDRAWAL SYNDROME; and RISKS FROM CONCOMITANT USE WITH BENZODIAZEPINES OR OTHER CNS DEPRESSANTS

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Important Dosage and Administration Information
- 2.2 Initial Dosage
- 2.3 Titration and Maintenance of Therapy
- 2.4 Discontinuation of Hydromorphone Hydrochloride Extended-Release Tablets
- 2.5 Dosage Modifications in Patients with Moderate Hepatic Impairment
- 2.6 Dosage Modifications in Patients with Renal Impairment

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Addiction, Abuse, and Misuse
- 5.2 Opioid Analgesic Risk Evaluation and Mitigation Strategy (REMS)
- 5.3 Life-threatening Respiratory Depression
- 5.4 Neonatal Opioid Withdrawal Syndrome
- 5.5 Risks from Concomitant use with Benzodiazepines or Other CNS Depressants
- 5.6 Life-Threatening Respiratory Depression in Patients with Chronic Pulmonary Disease or in Elderly, Cachectic, or Debilitated Patients
- 5.7 Adrenal Insufficiency
- 5.8 Severe Hypotension
- 5.9 Risks of Use in Patients with Increased Intracranial Pressure, Brain Tumors, Head Injury, or Impaired Consciousness
- 5.10 Risks of Use in Patients with Gastrointestinal Conditions
- 5.11 Increased Risk of Seizures in Patients with Seizure Disorders
- 5.12 Withdrawal
- 5.13 Sulfites
- 5.14 Risks of Driving and Operating Machinery

6 ADVERSE REACTIONS

- 6.1 Clinical Trial Experience
- 6.2 Postmarketing Experience

7 DRUG INTERACTIONS

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.3 Females and Males of Reproductive Potential

8.4 Pediatric Use

8.5 Geriatric Use

8.6 Hepatic Impairment

8.7 Renal Impairment

9 DRUG ABUSE AND DEPENDENCE

9.1 Controlled Substance

9.2 Abuse

9.3 Dependence

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: ADDICTION, ABUSE, AND MISUSE; RISK EVALUATION AND MITIGATION STRATEGY (REMS); LIFE-THREATENING RESPIRATORY DEPRESSION; ACCIDENTAL INGESTION; NEONATAL OPIOID WITHDRAWAL SYNDROME; and RISKS FROM CONCOMITANT USE WITH BENZODIAZEPINES OR OTHER CNS DEPRESSANTS

Addiction, Abuse, and Misuse

Hydromorphone hydrochloride extended-release tablets expose patients and other users to the risks of opioid addiction, abuse, and misuse, which can lead to overdose and death. Assess each patient's risk prior to prescribing hydromorphone hydrochloride extended-release tablets, and monitor all patients regularly for the development of these behaviors and conditions [see Warnings and Precautions (5.1)].

Opioid Analgesic Risk Evaluation and Mitigation Strategy (REMS):

To ensure that the benefits of opioid analgesics outweigh the risks of addiction, abuse, and misuse, the Food and Drug Administration (FDA) has required a REMS for these products [see Warnings and Precautions (5.2)]. Under the requirements of the REMS, drug companies with approved opioid analgesic products must make REMS-compliant education programs available to healthcare providers. Healthcare providers are strongly encouraged to

- complete a REMS-compliant education program,
- counsel patients and/or their caregivers, with every prescription, on safe use, serious risks, storage, and disposal of these products,
- emphasize to patients and their caregivers the importance of reading the Medication Guide every time it is provided by their pharmacist, and
- consider other tools to improve patient, household, and community safety.

Life-threatening Respiratory Depression

Serious, life-threatening, or fatal respiratory depression may occur with use of hydromorphone hydrochloride extended-release tablets. Monitor for respiratory depression, especially during initiation of hydromorphone hydrochloride extended-release tablets or following a dose increase. Instruct patients to swallow hydromorphone hydrochloride extended-release tablets whole; crushing, chewing, or dissolving hydromorphone hydrochloride extended-release tablets can cause rapid release and absorption of a potentially fatal dose of hydromorphone [see Warnings and Precautions (5.3)].

Accidental Ingestion

Accidental ingestion of even one dose of hydromorphone hydrochloride extended release-tablets, especially by children, can result in a fatal overdose of hydromorphone [see Warnings and Precautions (5.3)].

Neonatal Opioid Withdrawal Syndrome

Prolonged use of hydromorphone hydrochloride extended-release tablets during pregnancy can result in neonatal opioid withdrawal syndrome, which may be life-threatening if not recognized and treated, and requires management according to protocols developed by neonatology experts. If opioid use is required for a prolonged period in a pregnant woman, advise the patient of the risk of neonatal opioid withdrawal syndrome and ensure that appropriate treatment will be available [see Warnings and Precautions (5.4)].

Risks From Concomitant Use With Benzodiazepines Or Other CNS Depressants

Concomitant use of opioids with benzodiazepines or other central nervous system (CNS) depressants, including alcohol, may result in profound sedation, respiratory depression,

coma, and death [see *Warnings and Precautions (5.5)*, *Drug Interactions (7)*].

- **Reserve concomitant prescribing of hydromorphone hydrochloride extended-release tablets and benzodiazepines or other CNS depressants for use in patients for whom alternative treatment options are inadequate.**
- **Limit dosages and durations to the minimum required.**
- **Follow patients for signs and symptoms of respiratory depression and sedation.**

1 INDICATIONS AND USAGE

Hydromorphone hydrochloride extended-release tablets are indicated for the management of pain in opioid-tolerant patients severe enough to require daily, around-the-clock, long-term opioid treatment and for which alternative treatment options are inadequate.

Patients considered opioid tolerant are those who are receiving, for one week or longer, at least 60 mg oral morphine per day, 25 mcg transdermal fentanyl per hour, 30 mg oral oxycodone per day, 8 mg oral hydromorphone per day, 25 mg oral oxymorphone per day, 60 mg oral hydrocodone per day, or an equianalgesic dose of another opioid.

Limitations of Use

- Because of the risks of addiction, abuse, and misuse with opioids, even at recommended doses, and because of the greater risks of overdose and death with extended-release opioid formulations, reserve hydromorphone hydrochloride extended-release tablets for use in patients for whom alternative treatment options (e.g., non-opioid analgesics or immediate-release opioids) are ineffective, not tolerated, or would be otherwise inadequate to provide sufficient management of pain.
- Hydromorphone hydrochloride extended-release tablets are not indicated as an as-needed (prn) analgesic.

2 DOSAGE AND ADMINISTRATION

2.1 Important Dosage and Administration Information

To avoid medication errors, prescribers and pharmacists must be aware that hydromorphone is available as both immediate-release 8 mg tablets and extended-release 8 mg tablets.

Hydromorphone hydrochloride extended-release tablets should be prescribed only by healthcare professionals who are knowledgeable in the use of potent opioids for the management of chronic pain.

Due to the risk of respiratory depression, hydromorphone hydrochloride extended-release tablets are only indicated for use in patients who are already opioid-tolerant. Discontinue or taper all other extended-release opioids when beginning hydromorphone hydrochloride extended-release tablets therapy. As hydromorphone hydrochloride extended-release tablets are only for use in opioid-tolerant patients, do not begin any patient on hydromorphone hydrochloride extended-release tablets as the first opioid.

Patients who are opioid-tolerant are those receiving, for one week or longer, at least 60 mg of oral morphine per day, at least 25 mcg transdermal fentanyl per hour, at least 30 mg of oral oxycodone per day, at least 8 mg of oral hydromorphone per day, at least 25 mg oral oxymorphone per day, at least 60 mg oral hydrocodone per day, or an equianalgesic dose of another opioid.

- Use the lowest effective dosage for the shortest duration consistent with individual patient treatment goals [see *Warnings and Precautions (5)*].

- Initiate the dosing regimen for each patient individually, taking into account the patient's prior analgesic treatment experience and risk factors for addiction, abuse, and misuse [see *Warnings and Precautions (5.1)*].
- Monitor patients closely for respiratory depression, especially within the first 24 to 72 hours of initiating therapy and following dosage increases with hydromorphone hydrochloride extended-release tablets and adjust the dosage accordingly [see *Warnings and Precautions (5.3)*].

Instruct patients to swallow hydromorphone hydrochloride extended-release tablets whole [see *Patient Counseling Information (17)*]. Crushing, chewing, or dissolving hydromorphone hydrochloride extended-release tablets will result in uncontrolled delivery of hydromorphone and can lead to overdose or death [see *Warnings and Precautions (5.1)*].

2.2 Initial Dosage

Conversion from Other Oral Hydromorphone Formulations to Hydromorphone Hydrochloride Extended-Release Tablets

Patients receiving oral immediate-release hydromorphone may be converted to hydromorphone hydrochloride extended-release tablets by administering a starting dose equivalent to the patient's total daily oral hydromorphone dose, taken once daily.

Conversion from Other Oral Opioids to Hydromorphone Hydrochloride Extended-Release Tablets

Discontinue all other around-the-clock opioid drugs when hydromorphone hydrochloride extended-release tablets therapy is initiated.

There is substantial inter-patient variability in the relative potency of different opioid drugs and opioid formulations. Therefore, a conservative approach is advised when determining the total daily dosage of hydromorphone hydrochloride extended-release tablets. It is safer to underestimate a patient's 24-hour oral hydromorphone dosage and provide rescue medication (e.g., immediate-release opioid) than to overestimate the 24-hour oral hydromorphone dosage and manage an adverse reaction due to overdose.

In a hydromorphone hydrochloride extended-release tablets clinical trial with an open-label titration period, patients were converted from their prior opioid to hydromorphone hydrochloride extended-release tablets using the **Table 1** as a guide for the initial hydromorphone hydrochloride extended-release tablets dose. The recommended starting dose of hydromorphone hydrochloride extended-release tablets is 50% of the calculated estimate of daily hydromorphone requirement. Calculate the estimated daily hydromorphone requirement using **Table 1**.

Consider the following when using the information in **Table 1**:

- This is **not** a table of equianalgesic doses.
- The conversion factors in this table are only for the conversion **from** one of the listed oral opioid analgesics **to** hydromorphone hydrochloride extended release-tablets.
- The table **cannot** be used to convert **from** hydromorphone hydrochloride extended-release tablets to another opioid. Doing so will result in an overestimation of the dose of the new opioid and may result in fatal overdose.

Table 1.	
Conversion Factors to Hydromorphone Hydrochloride Extended-Release Tablets	
Prior Oral Opioid	Approximate Oral Conversion Factor
Hydromorphone	1
Codeine	0.06
Hydrocodone	0.4
Methadone	0.6

Morphine	0.2
Oxycodone	0.4
Oxymorphone	0.6

To calculate the estimated hydromorphone hydrochloride extended-release tablets dose using **Table 1**:

- For patients on a single opioid, sum the current total daily dose of the opioid and then multiply the total daily dose by the conversion factor to calculate the approximate oral hydromorphone daily dose.
- For patients on a regimen of more than one opioid, calculate the approximate oral hydromorphone dose for each opioid and sum the totals to obtain the approximate total hydromorphone daily dose.
- For patients on a regimen of fixed-ratio opioid/non-opioid analgesic products, use only the opioid component of these products in the conversion.

Always round the dose down, if necessary, to the appropriate hydromorphone hydrochloride extended-release tablet strength(s) available.

Example conversion from a single opioid to hydromorphone hydrochloride extended-release tablets:

Step 1: Sum the total daily dose of the opioid

- 30 mg of oxycodone 2 times daily = 60 mg total daily dose of oxycodone

Step 2: Calculate the approximate equivalent dose of oral hydromorphone based on the total daily dose of the current opioid using **Table 1**

- 60 mg total daily dose of oxycodone x Conversion Factor of 0.4 = 24 mg of oral hydromorphone daily

Step 3: Calculate the approximate starting dose of hydromorphone hydrochloride extended-release tablets to be given every 24 hours, which is 50% of the calculated oral hydromorphone dose. Round down, if necessary, to the appropriate hydromorphone hydrochloride extended-release tablet strengths available.

- 50% of 24 mg results in an initial dose of 12 mg of hydromorphone hydrochloride extended-release tablets once daily
- Adjust individually for each patient

Close observation and frequent titration are warranted until pain management is stable on the new opioid. Monitor patients for signs and symptoms of opioid withdrawal or for signs of over-sedation/toxicity after converting patients to hydromorphone hydrochloride extended-release tablets.

Conversion from Transdermal Fentanyl to Hydromorphone Hydrochloride Extended-Release Tablets

Eighteen hours following the removal of the transdermal fentanyl patch, hydromorphone hydrochloride extended-release tablets treatment can be initiated. To calculate the 24-hour hydromorphone hydrochloride extended-release tablets dose, use a conversion factor of 25 mcg/hr fentanyl transdermal patch to 12 mg of hydromorphone hydrochloride extended-release tablets. Then reduce the hydromorphone hydrochloride extended-release tablets dose by 50%.

For example:

Step 1: Identify the dose of transdermal fentanyl.

- 75 mg of transdermal fentanyl

Step 2: Use the conversion factor of 25 mcg/hr fentanyl transdermal patch to 12 mg of hydromorphone hydrochloride extended-release tablets.

- 75 mg of transdermal fentanyl : 36 mg total daily dose of hydromorphone hydrochloride extended-release tablets

Step 3: Calculate the approximate starting dose of hydromorphone hydrochloride extended-release tablets to be given every 24 hours, which is 50% of the converted dose. Round down, if necessary, to the appropriate hydromorphone hydrochloride extended-release tablet strengths available.

- 50% of 36 mg results in an initial dose of 18 mg, which would be rounded down to 16 mg of hydromorphone hydrochloride extended-release tablets once daily
- Adjust individually for each patient

Conversion from Methadone to Hydromorphone Hydrochloride Extended-Release Tablets

Close monitoring is of particular importance when converting from methadone to other opioid agonists. The ratio between methadone and other opioid agonists may vary widely as a function of previous dose exposure. Methadone has a long half-life and can accumulate in the plasma.

2.3 Titration and Maintenance of Therapy

Individually titrate hydromorphone hydrochloride extended-release tablets to a dose that provides adequate analgesia and minimizes adverse reactions. Continually reevaluate patients receiving hydromorphone hydrochloride extended-release tablets to assess the maintenance of pain control and the relative incidence of adverse reactions, as well as monitoring for the development of addiction, abuse, or misuse. Frequent communication is important among the prescriber, other members of the healthcare team, the patient, and the caregiver/family during periods of changing analgesic requirements, including initial titration. During chronic therapy, periodically reassess the continued need for opioid analgesics.

Plasma levels of hydromorphone hydrochloride extended-release tablets are sustained for 18 to 24 hours. Dosage adjustments of hydromorphone hydrochloride extended-release tablets may be made in increments of 4 to 8 mg every 3 to 4 days as needed to achieve adequate analgesia.

Patients who experience breakthrough pain may require a dose increase of hydromorphone hydrochloride extended-release tablets, or may need rescue medication with an appropriate dose of an immediate-release analgesic. If the level of pain increases after dose stabilization, attempt to identify the source of increased pain before increasing the hydromorphone hydrochloride extended-release tablets dose.

If unacceptable opioid-related adverse reactions are observed, the subsequent doses may be reduced. Adjust the dose to obtain an appropriate balance between management of pain and opioid-related adverse reactions.

2.4 Discontinuation of Hydromorphone Hydrochloride Extended-Release Tablets

When a patient no longer requires therapy with hydromorphone hydrochloride extended-release tablets, taper doses gradually, by 25% to 50% every 2 to 3 days while monitoring carefully for signs and symptoms of withdrawal. If the patient develops these signs or symptoms, raise the dose to the previous level and taper more slowly, either by increasing the interval between decreases, decreasing the amount of change in dose, or both. Do not abruptly discontinue hydromorphone hydrochloride extended-release tablets.

To dispose of unused hydromorphone hydrochloride extended-release tablets flush all remaining tablets down the toilet or remit to authorities at a certified drug take-back program.

2.5 Dosage Modifications in Patients with Moderate Hepatic Impairment

Start patients with moderate hepatic impairment on 25% of the hydromorphone hydrochloride extended-release tablets dose that would be prescribed for patients with normal hepatic function. Closely monitor patients with moderate hepatic impairment for respiratory and central nervous system depression during initiation of therapy with hydromorphone hydrochloride extended-release tablets and during dose titration. Use of alternate analgesics is recommended for patients with severe hepatic impairment [*see Use in Specific Populations (8.6)*].

2.6 Dosage Modifications in Patients with Renal Impairment

Start patients with moderate renal impairment on 50% of the hydromorphone hydrochloride extended-release tablets dose that would be prescribed for patients with normal renal function. Closely monitor patients with renal impairment for respiratory and central nervous system depression during initiation of therapy with hydromorphone hydrochloride extended-release tablets and during dose titration. As hydromorphone hydrochloride extended-release tablets are only intended for once daily administration, consider use of an alternate analgesic that may permit more flexibility with the dosing interval in patients with severe renal impairment [*see Use in Specific Populations (8.7)*].

3 DOSAGE FORMS AND STRENGTHS

Extended-release tablets: available in 8 mg, 12 mg, 16 mg or 32 mg dosage strengths.

8 mg tablets: round, biconvex, dark beige tablets imprinted with “P293” over “8” on one side.

12 mg tablets: round, biconvex, white tablets imprinted with “P294” over “12” on one side.

16 mg tablets: round, biconvex, light beige tablets imprinted with “P295” over “16” on one side.

32 mg tablets: round, biconvex, pink tablets imprinted with "P297" over "32" on one side.

4 CONTRAINDICATIONS

Hydromorphone hydrochloride extended-release tablets are contraindicated in:

- Opioid non-tolerant patients. Fatal respiratory depression could occur in patients who are not opioid tolerant.
- Patients with significant respiratory depression [*see Warnings and Precautions (5.3)*].
- Acute or severe bronchial asthma in an unmonitored setting or in the absence of resuscitative equipment [*see Warnings and Precautions (5.6)*].
- Known or suspected gastrointestinal obstruction, including paralytic ileus [*see Warnings and Precautions (5.10)*].
- Patients who have had surgical procedures and/or underlying disease resulting in narrowing of the gastrointestinal tract, or have “blind loops” of the gastrointestinal tract or gastrointestinal obstruction [*see Warnings and Precautions (5.10)*].
- Patients with hypersensitivity (e.g., anaphylaxis) to hydromorphone [*see Warnings and Precautions (5.13)*].

5 WARNINGS AND PRECAUTIONS

5.1 Addiction, Abuse, and Misuse

Hydromorphone hydrochloride extended-release tablets contain hydromorphone, a Schedule II controlled substance. As an opioid, hydromorphone hydrochloride extended-release tablets expose users to the risks of addiction, abuse, and misuse [*see Drug Abuse and Dependence (9)*]. As modified-release products such as hydromorphone hydrochloride extended-release tablets deliver the opioid

over an extended period of time, there is a greater risk for overdose and death due to the larger amount of hydromorphone present.

Although the risk of addiction in any individual is unknown, it can occur in patients appropriately prescribed hydromorphone hydrochloride extended-release tablets and in those who obtain the drug illicitly. Addiction can occur at recommended doses and if the drug is misused or abused.

Assess each patient's risk for opioid addiction, abuse, or misuse prior to prescribing hydromorphone hydrochloride extended-release tablets, and monitor all patients receiving hydromorphone hydrochloride extended-release tablets for the development of these behaviors and conditions. Risks are increased in patients with a personal or family history of substance abuse (including drug or alcohol addiction or abuse) or mental illness (e.g., major depression). The potential for these risks should not, however, prevent the prescribing of hydromorphone hydrochloride extended-release tablets for the proper management of pain in any given patient. Patients at increased risk may be prescribed modified-release opioid formulations such as hydromorphone hydrochloride extended-release tablets, but use in such patients necessitates intensive counseling about the risks and proper use of hydromorphone hydrochloride extended-release tablets along with intensive monitoring for signs of addiction, abuse, and misuse.

Abuse or misuse of hydromorphone hydrochloride extended-release tablets by crushing, chewing, snorting, or injecting the dissolved product will result in the uncontrolled delivery of hydromorphone and can result in overdose and death [*see Overdosage (10)*].

Opioids are sought by drug abusers and people with addiction disorders and are subject to criminal diversion. Consider these risks when prescribing or dispensing hydromorphone hydrochloride extended-release tablets. Strategies to reduce these risks include prescribing the drug in the smallest appropriate quantity and advising the patient on the proper disposal of unused drug [*see Patient Counseling Information (17)*]. Contact local state professional licensing board or state controlled substances authority for information on how to prevent and detect abuse or diversion of this product.

5.2 Opioid Analgesic Risk Evaluation and Mitigation Strategy (REMS)

To ensure that the benefits of opioid analgesics outweigh the risks of addiction, abuse, and misuse, the Food and Drug Administration (FDA) has required a Risk Evaluation and Mitigation Strategy (REMS) for these products. Under the requirements of the REMS, drug companies with approved opioid analgesic products must make REMS-compliant education programs available to healthcare providers. Healthcare providers are strongly encouraged to do all of the following:

- Complete a REMS-compliant education program offered by an accredited provider of continuing education (CE) or another education program that includes all the elements of the FDA Education Blueprint for Health Care Providers Involved in the Management or Support of Patients with Pain.
- Discuss the safe use, serious risks, and proper storage and disposal of opioid analgesics with patients and/or their caregivers every time these medicines are prescribed. The Patient Counseling Guide (PCG) can be obtained at this link: www.fda.gov/OpioidAnalgesicREMSPCG.
- Emphasize to patients and their caregivers the importance of reading the Medication Guide that they will receive from their pharmacist every time an opioid analgesic is dispensed to them.
- Consider using other tools to improve patient, household, and community safety, such as patient prescriber agreements that reinforce patient-prescriber responsibilities.

To obtain further information on the opioid analgesic REMS and for a list of accredited REMS CME/CE, call 1-800-503-0784, or log on to www.opioidanalgesicrems.com. The FDA Blueprint can be found at www.fda.gov/OpioidAnalgesicREMSBlueprint.

5.3 Life-threatening Respiratory Depression

Serious, life-threatening, or fatal respiratory depression has been reported with the use of modified-

release opioids, even when used as recommended. Respiratory depression from opioid use, if not immediately recognized and treated, may lead to respiratory arrest and death. Management of respiratory depression may include close observation, supportive measures, and use of opioid antagonists, depending on the patient's clinical status [see *Overdosage (10)*]. Carbon dioxide (CO₂) retention from opioid-induced respiratory depression can exacerbate the sedating effects of opioids.

While serious, life-threatening, or fatal respiratory depression can occur at any time during the use of hydromorphone hydrochloride extended-release tablets, the risk is greatest during the initiation of therapy or following a dosage increase. Closely monitor patients for respiratory depression, especially within the first 24 to 72 hours of initiating therapy with hydromorphone hydrochloride extended-release tablets and following dosage increases.

To reduce the risk of respiratory depression, proper dosing and titration of hydromorphone hydrochloride extended-release tablets are essential [see *Dosage and Administration (2)*]. Overestimating the hydromorphone hydrochloride extended-release tablets dose when converting patients from another opioid product can result in fatal overdose with the first dose.

Accidental ingestion of even one dose of hydromorphone hydrochloride extended-release tablets, especially by children, can result in respiratory depression and death due to an overdose of hydromorphone.

5.4 Neonatal Opioid Withdrawal Syndrome

Prolonged use of hydromorphone hydrochloride extended-release tablets during pregnancy can result in withdrawal in the neonate. Neonatal opioid withdrawal syndrome, unlike opioid withdrawal syndrome in adults, may be life-threatening if not recognized and treated, and requires management according to protocols developed by neonatology experts. Observe newborns for signs of neonatal opioid withdrawal syndrome and manage accordingly. Advise pregnant women using opioids for a prolonged period of the risk of neonatal opioid withdrawal syndrome and ensure that appropriate treatment will be available [see *Use in Specific Populations (8.1)*, *Patient Counseling Information (17)*].

5.5 Risks from Concomitant use with Benzodiazepines or Other CNS Depressants

Profound sedation, respiratory depression, coma, and death may result from the concomitant use of hydromorphone hydrochloride extended-release tablets with benzodiazepines or other CNS depressants (e.g., non-benzodiazepine sedatives/hypnotics, anxiolytics, tranquilizers, muscle relaxants, general anesthetics, antipsychotics, other opioids, alcohol). Because of these risks, reserve concomitant prescribing of these drugs for use in patients for whom alternative treatment options are inadequate.

Observational studies have demonstrated that concomitant use of opioid analgesics and benzodiazepines increases the risk of drug-related mortality compared to use of opioid analgesics alone. Because of similar pharmacological properties, it is reasonable to expect similar risk with the concomitant use of other CNS depressant drugs with opioid analgesics [see *Drug Interactions (7)*].

If the decision is made to prescribe a benzodiazepine or other CNS depressant concomitantly with an opioid analgesic, prescribe the lowest effective dosages and minimum durations of concomitant use. In patients already receiving an opioid analgesic, prescribe a lower initial dose of the benzodiazepine or other CNS depressant than indicated in the absence of an opioid, and titrate based on clinical response. If an opioid analgesic is initiated in a patient already taking a benzodiazepine or other CNS depressant, prescribe a lower initial dose of the opioid analgesic, and titrate based on clinical response. Follow patients closely for signs and symptoms of respiratory depression and sedation.

Advise both patients and caregivers about the risks of respiratory depression and sedation when hydromorphone hydrochloride extended-release tablets are used with benzodiazepines or other CNS depressants (including alcohol and illicit drugs). Advise patients not to drive or operate heavy machinery until the effects of concomitant use of the benzodiazepine or other CNS depressant have been determined. Screen patients for risk of substance use disorders, including opioid abuse and misuse, and warn them of the risk for overdose and death associated with the use of additional CNS

depressants including alcohol and illicit drugs [see *Drug Interactions (7), Patient Counseling Information (17)*].

5.6 Life-Threatening Respiratory Depression in Patients with Chronic Pulmonary Disease or in Elderly, Cachectic, or Debilitated Patients

The use of hydromorphone hydrochloride extended-release tablets in patients with acute or severe bronchial asthma in an unmonitored setting or in the absence of resuscitative equipment is contraindicated.

Patients with Chronic Pulmonary Disease: Hydromorphone hydrochloride extended-release tablets treated patients with significant chronic obstructive pulmonary disease or cor pulmonale, and those with a substantially decreased respiratory reserve, hypoxia, hypercapnia, or pre-existing respiratory depression are at increased risk of decreased respiratory drive including apnea, even at recommended dosages of hydromorphone hydrochloride extended-release tablets [see *Warnings and Precautions (5.3)*].

Elderly, Cachectic, or Debilitated Patients: Life-threatening respiratory depression is more likely to occur in elderly, cachectic, or debilitated patients because they may have altered pharmacokinetics or altered clearance compared to younger, healthier patients [see *Warnings and Precautions (5.3)*].

Monitor such patients closely, particularly when initiating and titrating hydromorphone hydrochloride extended-release tablets and when hydromorphone hydrochloride extended-release tablets are given concomitantly with other drugs that depress respiration [see *Warnings and Precautions (5.3, 5.5)*]. Alternatively, consider the use of non-opioid analgesics in these patients.

5.7 Adrenal Insufficiency

Cases of adrenal insufficiency have been reported with opioid use, more often following greater than one month of use. Presentation of adrenal insufficiency may include non-specific symptoms and signs including nausea, vomiting, anorexia, fatigue, weakness, dizziness, and low blood pressure. If adrenal insufficiency is suspected, confirm the diagnosis with diagnostic testing as soon as possible. If adrenal insufficiency is diagnosed, treat with physiologic replacement doses of corticosteroids. Wean the patient off of the opioid to allow adrenal function to recover and continue corticosteroid treatment until adrenal function recovers. Other opioids may be tried as some cases reported use of a different opioid without recurrence of adrenal insufficiency. The information available does not identify any particular opioids as being more likely to be associated with adrenal insufficiency.

5.8 Severe Hypotension

Hydromorphone hydrochloride extended-release tablets may cause severe hypotension including orthostatic hypotension and syncope in ambulatory patients. There is an increased risk in patients whose ability to maintain blood pressure has already been compromised by a reduced blood volume, or concurrent administration of certain CNS depressant drugs (e.g. phenothiazines or general anesthetics) [see *Drug Interactions (7)*]. Monitor these patients for signs of hypotension after initiating or titrating the dosage of hydromorphone hydrochloride extended-release tablets. In patients with circulatory shock, hydromorphone hydrochloride extended-release tablets may cause vasodilation that can further reduce cardiac output and blood pressure. Avoid the use of hydromorphone hydrochloride extended-release tablets in patients with circulatory shock.

5.9 Risks of Use in Patients with Increased Intracranial Pressure, Brain Tumors, Head Injury, or Impaired Consciousness

In patients who may be susceptible to the intracranial effects of CO₂ retention (e.g., those with evidence of increased intracranial pressure or brain tumors), hydromorphone hydrochloride extended-release tablets may reduce respiratory drive, and the resultant CO₂ retention can further increase intracranial pressure. Monitor such patients for signs of sedation and respiratory depression, particularly when

initiating therapy with hydromorphone hydrochloride extended-release tablets.

Opioids may also obscure the clinical course in a patient with a head injury. Avoid the use of hydromorphone hydrochloride extended-release tablets in patients with impaired consciousness or coma.

5.10 Risks of Use in Patients with Gastrointestinal Conditions

Hydromorphone hydrochloride extended-release tablets are contraindicated in patients with known or suspected gastrointestinal obstruction, including paralytic ileus. Avoid the use of hydromorphone hydrochloride extended-release tablets in patients with other GI obstruction.

Because the shell of the hydromorphone hydrochloride extended-release tablet is insoluble, it may remain intact during gastrointestinal transit and be eliminated in the feces. Hydromorphone hydrochloride extended-release tablets are contraindicated in patients with preexisting severe gastrointestinal narrowing (pathologic or iatrogenic, for example: esophageal motility disorders, small bowel inflammatory disease, “short gut” syndrome due to adhesions or decreased transit time, past history of peritonitis, cystic fibrosis, chronic intestinal pseudoobstruction, or Meckel’s diverticulum). There have been reports of obstructive symptoms in patients with known strictures or risk of strictures, such as previous GI surgery, in association with the ingestion of drugs in nondeformable extended-release formulations.

It is possible that hydromorphone hydrochloride extended-release tablets may be visible on abdominal x-rays under certain circumstances, especially when digital enhancing techniques are utilized.

The hydromorphone in hydromorphone hydrochloride extended-release tablets may cause spasm of the sphincter of Oddi. Opioids may cause increases in serum amylase. Monitor patients with biliary tract disease, including acute pancreatitis, for worsening symptoms.

5.11 Increased Risk of Seizures in Patients with Seizure Disorders

The hydromorphone in hydromorphone hydrochloride extended-release tablets may increase the frequency of seizures in patients with seizure disorders, and may increase the risk of seizures occurring in other clinical settings associated with seizures. Monitor patients with a history of seizure disorders for worsened seizure control during hydromorphone hydrochloride extended-release tablets therapy.

5.12 Withdrawal

Avoid the use of mixed agonist/antagonist (e.g., pentazocine, nalbuphine, and butorphanol) or partial agonist (e.g., buprenorphine) analgesics in patients who are receiving a full opioid agonist analgesic, including hydromorphone hydrochloride extended-release tablets. In these patients, mixed agonist/antagonist and partial agonist analgesics may reduce the analgesic effect and/or may precipitate withdrawal symptoms [*see Drug Interactions (7)*].

When discontinuing hydromorphone hydrochloride extended-release tablets, gradually taper the dose [*see Dosage and Administration (2.4)*]. Do not abruptly discontinue hydromorphone hydrochloride extended-release tablets [*see Drug Abuse and Dependence (9.3)*].

5.13 Sulfites

Hydromorphone hydrochloride extended-release tablets contain sodium metabisulfite, a sulfite that may cause allergic-type reactions including anaphylactic symptoms and life-threatening or less severe asthmatic episodes in certain susceptible people. The overall prevalence of sulfite sensitivity in the general population is unknown and probably low. Sulfite sensitivity is seen more frequently in asthmatic than in nonasthmatic people [*see Adverse Reactions (6.2)*].

5.14 Risks of Driving and Operating Machinery

Hydromorphone hydrochloride extended-release tablets may impair the mental and/or physical abilities needed to perform potentially hazardous activities such as driving a car or operating machinery. Warn patients not to drive or operate dangerous machinery unless they are tolerant to the effects of hydromorphone hydrochloride extended-release tablets and know how they will react to the medication [see Patient Counseling Information (17)].

6 ADVERSE REACTIONS

The following serious adverse reactions are discussed elsewhere in the labeling:

- Addiction, Abuse, and Misuse [see Warnings and Precautions (5.1)]
- Life Threatening Respiratory Depression [see Warnings and Precautions (5.3)]
- Neonatal Opioid Withdrawal Syndrome [see Warnings and Precautions (5.4)]
- Interactions with Benzodiazepine or Other CNS Depressants [see Warnings and Precautions (5.5)]
- Adrenal Insufficiency [see Warnings and Precautions (5.7)]
- Severe Hypotension [see Warnings and Precautions (5.8)]
- Gastrointestinal Adverse Reactions [see Warnings and Precautions (5.10)]
- Seizures [see Warnings and Precautions (5.11)]
- Withdrawal [see Warnings and Precautions (5.12)]

6.1 Clinical Trial Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

Hydromorphone hydrochloride extended-release tablets were administered to a total of 2,524 patients in 15 controlled and uncontrolled clinical studies. Of these, 423 patients were exposed to hydromorphone hydrochloride extended-release tablets for greater than 6 months and 141 exposed for greater than one year.

The most common adverse reactions leading to study discontinuation were nausea, vomiting, constipation, somnolence, and dizziness. The most common treatment-related serious adverse reactions from controlled and uncontrolled chronic pain studies were drug withdrawal syndrome, overdose, confusional state, and constipation.

The overall incidence of adverse reactions in patients greater than 65 years of age was higher, with a greater than 5% difference in rates for constipation and nausea when compared with younger patients. The overall incidence of adverse reactions in female patients was higher, with a greater than 5% difference in rates for nausea, vomiting, constipation and somnolence when compared with male patients.

A 12-week double-blind, placebo-controlled, randomized withdrawal study was conducted in opioid tolerant patients with moderate to severe low back pain [see Clinical Studies (14)]. A total of 447 patients were enrolled into the open-label titration phase with 268 patients randomized into the double-blind treatment phase. The adverse reactions that were reported in at least 2% of the patients are contained in **Table 2**.

Table 2.

Number (%) of Patients with Adverse Reactions Reported in \geq 2% of Patients with Moderate to Severe Low Back Pain During the Open-Label Titration Phase or Double-Blind Treatment Phase by Preferred Term

Preferred Term	Open-Label Titration	Double-Blind
----------------	----------------------	--------------

	Phase	Treatment Phase	
	Hydromorphone Hydrochloride Extended-Release Tablets (N=447)	Hydromorphone Hydrochloride Extended-Release Tablets (N=134)	Placebo (N=134)
Constipation	69 (15)	10 (7)	5 (4)
Nausea	53 (12)	12 (9)	10 (7)
Somnolence	39 (9)	1 (1)	0 (0)
Headache	35 (8)	7 (5)	10 (7)
Vomiting	29 (6)	8 (6)	6 (4)
Pruritus	21 (5)	1 (1)	0 (0)
Dizziness	17 (4)	3 (2)	2 (1)
Insomnia	13 (3)	7 (5)	5 (4)
Dry Mouth	13 (3)	2 (1)	0 (0)
Edema Peripheral	13 (3)	3 (2)	1 (1)
Hyperhidrosis	13 (3)	2 (1)	2 (1)
Anorexia/Decreased Appetite	10 (2)	2 (1)	0 (0)
Arthralgia	9 (2)	8 (6)	3 (2)
Abdominal pain	9 (2)	4 (3)	3 (2)
Muscle Spasms	5 (1)	3 (2)	1 (1)
Weight Decreased	3 (1)	4 (3)	3 (2)

The adverse reactions that were reported in at least 2% of the total treated patients (N=2,474) in the 14 chronic clinical trials are contained in **Table 3**.

Table 3.

Number (%) of Patients with Adverse Reactions Reported in \geq 2% of Patients with Chronic Pain Receiving Hydromorphone Hydrochloride Extended-Release Tablets in 14 Clinical Studies by Preferred Term

Preferred Term	All Patients (N=2,474)
Constipation	765 (31)
Nausea	684 (28)
Vomiting	337 (14)
Somnolence	367 (15)
Headache	308 (12)
As thenia/Fatigue	272 (11)
Dizziness	262 (11)
Diarrhea	201 (8)
Pruritus	193 (8)
Insomnia	161 (7)
Hyperhidrosis	143 (6)
Edema Peripheral	135 (5)
Anorexia/Decreased Appetite	139 (6)
Dry Mouth	121 (5)
Abdominal Pain	115 (5)
Anxiety	95 (4)

Back Pain	95 (4)
Dyspepsia*	88 (4)
Depression	81 (3)
Dyspnea	76 (3)
Muscle Spasms	74 (3)
Arthralgia	72 (3)
Rash	64 (3)
Pain in Extremity	63 (3)
Pain	58 (2)
Drug Withdrawal Syndrome	55 (2)
Pyrexia	52 (2)
Fall	51 (2)
Chest Pain	51 (2)

* Reflux esophagitis, gastroesophageal reflux disease and Barrett's esophagus were grouped and reported with dyspepsia

The following Adverse Reactions occurred in patients with an overall frequency of <2% and are listed in descending order within each System Organ Class:

Cardiac disorders: palpitations, tachycardia, bradycardia, extrasystoles

Ear and labyrinth disorders: vertigo, tinnitus

Endocrine disorders: hypogonadism

Eye disorders: vision blurred, diplopia, dry eye, miosis

Gastrointestinal disorders: flatulence, dysphagia, hematochezia, abdominal distension, hemorrhoids, abnormal feces, intestinal obstruction, eructation, diverticulum, gastrointestinal motility disorder, large intestine perforation, anal fissure, bezoar, duodenitis, ileus, impaired gastric emptying, painful defecation

General disorders and administration site conditions: chills, malaise, feeling abnormal, feeling of body temperature change, feeling jittery, hangover, gait disturbance, feeling drunk, body temperature decreased

Infections and infestations: gastroenteritis, diverticulitis

Injury, poisoning and procedural complications: contusion, overdose

Investigations: weight decreased, hepatic enzyme increased, blood potassium decreased, blood amylase increased, blood testosterone decreased

Metabolism and nutrition disorders: dehydration, fluid retention, increased appetite, hyperuricemia

Musculoskeletal and connective tissue disorders: myalgia

Nervous system disorders: tremor, sedation, hypoesthesia, paresthesia, disturbance in attention, memory impairment, dysarthria, syncope, balance disorder, dysgeusia, depressed level of consciousness, coordination abnormal, hyperesthesia, myoclonus, dyskinesia, crying, hyperreflexia, encephalopathy, cognitive disorder, convulsion, psychomotor hyperactivity

Psychiatric disorders: confusional state, nervousness, restlessness, abnormal dreams, mood altered, hallucination, panic attack, euphoric mood, paranoia, dysphoria, listless, suicide ideation, libido decreased, aggression

Renal and urinary disorders: dysuria, urinary retention, urinary frequency, urinary hesitation, micturition disorder

Reproductive system and breast disorders: erectile dysfunction, sexual dysfunction

Respiratory, thoracic and mediastinal disorders: rhinorrhea, respiratory distress, hypoxia, bronchospasm, sneezing, hyperventilation, respiratory depression

Skin and subcutaneous tissue disorders: erythema

Vascular disorders: flushing, hypertension, hypotension

6.2 Postmarketing Experience

The following adverse reactions have been identified during post approval use of hydromorphone. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Serotonin syndrome: Cases of serotonin syndrome, a potentially life-threatening condition, have been reported during concomitant use of opioids with serotonergic drugs [see *Drug Interactions (7)*].

Adrenal insufficiency: Cases of adrenal insufficiency have been reported with opioid use, more often following greater than one month of use [see *Warnings and Precautions (5.7)*].

Anaphylaxis: Anaphylactic reaction has been reported with ingredients contained in hydromorphone hydrochloride extended-release tablets [see *Contraindications (4) and Warnings and Precautions (5.13)*].

Androgen deficiency: Cases of androgen deficiency have occurred with chronic use of opioids [see *Clinical Pharmacology (12.2)*].

7 DRUG INTERACTIONS

Table 4 includes clinically significant drug interactions with hydromorphone hydrochloride extended-release tablets.

Table 4: Clinically Significant Drug Interactions with Hydromorphone Hydrochloride Extended-Release Tablets

Benzodiazepines and Other Central Nervous System (CNS) Depressants	
<i>Clinical Impact:</i>	Due to additive pharmacologic effect, the concomitant use of benzodiazepines or other CNS depressants, including alcohol, can increase the risk of hypotension, respiratory depression, profound sedation, coma, and death.
<i>Intervention:</i>	Reserve concomitant prescribing of these drugs for use in patients for whom alternative treatment options are inadequate. Limit dosages and durations to the minimum required. Follow patients closely for signs of respiratory depression and sedation [see <i>Warnings and Precautions (5.5)</i>].
<i>Examples:</i>	Benzodiazepines and other sedatives/hypnotics, anxiolytics, tranquilizers, muscle relaxants, general anesthetics, antipsychotics, other opioids, alcohol.
Serotonergic Drugs	
<i>Clinical Impact:</i>	The concomitant use of opioids with other drugs that affect the serotonergic neurotransmitter system has resulted in serotonin syndrome.
<i>Intervention:</i>	If concomitant use is warranted, carefully observe the patient, particularly during treatment initiation and dose adjustment. Discontinue hydromorphone hydrochloride extended-release tablet if serotonin syndrome is suspected.
<i>Examples:</i>	Selective serotonin reuptake inhibitors (SSRIs), serotonin and norepinephrine reuptake inhibitors (SNRIs), tricyclic antidepressants (TCAs), triptans, 5-HT ₃ receptor antagonists, drugs that affect the serotonin neurotransmitter system (e.g., mirtazapine, trazodone, tramadol), monoamine oxidase (MAO) inhibitors (those intended to treat psychiatric disorders and also others, such as linezolid and intravenous methylene blue).

Monoamine Oxidase Inhibitors (MAOIs)	
<i>Clinical Impact:</i>	MAOI interactions with opioids may manifest as serotonin syndrome or opioid toxicity (e.g., respiratory depression, coma) [see <i>Warnings and Precautions (5.5)</i>].
<i>Intervention:</i>	The use of hydromorphone hydrochloride extended-release tablets is not recommended for patients taking MAOIs or within 14 days of stopping such treatment.
<i>Examples:</i>	phenelzine, tranylcypromine, linezolid
Mixed Agonist/Antagonist and Partial Agonist Opioid Analgesics	
<i>Clinical Impact:</i>	May reduce the analgesic effect of hydromorphone hydrochloride extended-release tablets and/or precipitate withdrawal symptoms [see <i>Warnings and Precautions (5.12)</i>].
<i>Intervention:</i>	Avoid concomitant use.
<i>Examples:</i>	butorphanol, nalbuphine, pentazocine, buprenorphine
Muscle Relaxants	
<i>Clinical Impact:</i>	Hydromorphone may enhance the neuromuscular blocking action of skeletal muscle relaxants and produce an increase degree of respiratory depression [see <i>Warnings and Precautions (5.5)</i>].
<i>Intervention:</i>	Monitor patients for signs of respiratory depression that may be greater than otherwise expected and decrease the dosage of hydromorphone hydrochloride extended-release tablets and/or the muscle relaxant as necessary.
Diuretics	
<i>Clinical Impact:</i>	Opioids can reduce the efficacy of diuretics by inducing the release of antidiuretic hormone.
<i>Intervention:</i>	Monitor patients for signs of diminished diuresis and/or effects on blood pressure and increase the dosage of the diuretic as needed.
Anticholinergic Drugs	
<i>Clinical Impact:</i>	The concomitant use of anticholinergic drugs may increase risk of urinary retention and/or severe constipation, which may lead to paralytic ileus.
<i>Intervention:</i>	Monitor patients for signs of urinary retention or reduced gastric motility when hydromorphone hydrochloride extended-release tablets are used concomitantly with anticholinergic drugs.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Prolonged use of opioid analgesics during pregnancy may cause neonatal opioid withdrawal syndrome [see *Warnings and Precautions (5.4)*]. There are no adequate and well-controlled studies in pregnant women. Based on animal data, advise pregnant women of the potential risk to a fetus.

In animal reproduction studies, reduced postnatal survival of pups, developmental delays, and altered behavioral responses were noted following oral treatment of pregnant rats with hydromorphone during gestation and through lactation at doses 2.1 times the human daily dose of 32 mg/day (HDD), respectively. In published studies, neural tube defects were noted following subcutaneous injection of hydromorphone to pregnant hamsters at doses 4.8 times the HDD and soft tissue and skeletal abnormalities were noted following subcutaneous continuous infusion of 2.3 times the HDD to pregnant mice. No malformations were noted at 2.1 or 17 times the HDD in pregnant rats or rabbits, respectively [see *Data*]. Based on animal data, advise pregnant women of the potential risk to a fetus.

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. In

the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

Clinical Considerations

Fetal/Neonatal Adverse Reactions

Prolonged use of opioid analgesics during pregnancy for medical or nonmedical purposes can result in physical dependence in the neonate and neonatal opioid withdrawal syndrome shortly after birth. Neonatal opioid withdrawal syndrome presents as irritability, hyperactivity and abnormal sleep pattern, high pitched cry, tremor, vomiting, diarrhea, and failure to gain weight. The onset, duration, and severity of neonatal opioid withdrawal syndrome vary based on the specific opioid used, duration of use, timing and amount of last maternal use, and rate of elimination of the drug by the newborn. Observe newborns for symptoms of neonatal opioid withdrawal syndrome, and manage accordingly [see *Warnings and Precautions* (5.4)].

Labor or Delivery

Opioids cross the placenta and may produce respiratory depression and psycho-physiologic effects in neonates. An opioid antagonist, such as naloxone, must be available for reversal of opioid-induced respiratory depression in the neonate. Hydromorphone hydrochloride extended-release tablets are not recommended for use in pregnant women during or immediately prior to labor, when use of shorter-acting analgesics or other analgesic techniques are more appropriate. Opioid analgesics, including hydromorphone hydrochloride extended-release tablets can prolong labor through actions which temporarily reduce the strength, duration, and frequency of uterine contractions. However, this effect is not consistent and may be offset by an increased rate of cervical dilation, which tends to shorten labor. Monitor neonates exposed to opioid analgesics during labor for signs of excess sedation and respiratory depression.

Data

Animal Data

Pregnant rats were treated with hydromorphone hydrochloride from Gestation Day 6 to 17 via oral gavage doses of 1.75, 3.5, or 7 mg/kg/day (0.5, 1.1, or 2.1 times the HDD of 32 mg/day based on body surface area, respectively). Maternal toxicity was noted in all treatment groups (reduced food consumption and body weights in the two highest dose groups). There was no evidence of malformations or embryotoxicity reported.

Pregnant rabbits were treated with hydromorphone hydrochloride from Gestation Day 6 to 20 via oral gavage doses of 10, 25, or 50 mg/kg/day (4.3, 8.5, or 17 times the HDD of 32 mg/day based on body surface area, respectively). Maternal toxicity was noted in the highest dose group (reduced food consumption and body weights). There was no evidence of malformations or embryotoxicity reported.

In a published study, neural tube defects (exencephaly and cranioschisis) were noted following subcutaneous administration of hydromorphone hydrochloride (19 to 258 mg/kg) on Gestation Day 8 to pregnant hamsters (4.8 to 65.4 times the HDD of 32 mg/day based on body surface area). The findings cannot be clearly attributed to maternal toxicity. No neural tube defects were noted at 14 mg/kg (3.5 times the human daily dose of 32 mg/day). In a published study, CF-1 mice were treated subcutaneously with continuous infusion of 7.5, 15, or 30 mg/kg/day hydromorphone hydrochloride (1.1, 2.3, or 4.6 times the human daily dose of 32 mg based on body surface area) via implanted osmotic pumps during organogenesis (Gestation Days 7 to 10). Soft tissue malformations (cryptorchidism, cleft palate, malformed ventricles and retina), and skeletal variations (split supraoccipital, checkerboard and split sternbrae, delayed ossification of the paws and ectopic ossification sites) were observed at doses 2.3 times the human dose of 32 mg/day based on body surface area. The findings cannot be clearly attributed to maternal toxicity.

Pregnant rats were treated with hydromorphone hydrochloride from Gestation Day 6 to Lactation Day 21 via oral gavage doses of 1.75, 3.5, or 7 mg/kg/day (0.5, 1.1, or 2.1 times the HDD of 32 mg/day based

on body surface area, respectively). Reduced pup weights were noted at 1.1 and 2.1 times the human daily dose of 32 mg/day and increased pup deaths, delayed ear opening, reduced auditory startle reflex, and reduced open-field activity were also noted at 2.1 times the HDD. Maternal toxicity was noted in all treatment groups (reduced food consumption and body weights in all groups) and decreased maternal care in the high dose group.

8.2 Lactation

Risk Summary

Because of the potential for serious adverse reactions, including excess sedation and respiratory depression in a breastfed infant, advise patients that breastfeeding is not recommended during treatment with hydromorphone hydrochloride extended-release tablets. Low concentrations of hydromorphone have been detected in human milk in clinical trials. Withdrawal symptoms can occur in breastfeeding infants when maternal administration of an opioid analgesic is stopped. Nursing should not be undertaken while a patient is receiving hydromorphone hydrochloride extended-release tablets since hydromorphone is excreted in the milk.

Clinical Considerations

Monitor infants exposed to hydromorphone hydrochloride extended-release tablets through breast milk for excess sedation and respiratory depression. Withdrawal symptoms can occur in breastfed infants when maternal administration of an opioid analgesic is stopped, or when breast-feeding is stopped.

8.3 Females and Males of Reproductive Potential

Infertility

Chronic use of opioids may cause reduced fertility in females and males of reproductive potential. It is not known whether these effects on fertility are reversible [*see Adverse Reactions (6.2), Nonclinical Toxicology (13.1)*].

8.4 Pediatric Use

The safety and effectiveness of hydromorphone hydrochloride extended-release tablets in patients 17 years of age and younger have not been established.

8.5 Geriatric Use

Elderly patients (aged 65 years or older) may have increased sensitivity to hydromorphone. In general, use caution when selecting a dosage for an elderly patient, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function and of concomitant disease or other drug therapy.

Respiratory depression is the chief risk for elderly patients treated with opioids and has occurred after large initial doses were administered to patients who were not opioid-tolerant or when opioids were coadministered with other agents that depress respiration. Titrate the dosage of hydromorphone hydrochloride extended-release tablets slowly in geriatric patients and monitor closely for signs of central nervous system and respiratory depression [*see Warnings and Precautions (5.3)*].

Hydromorphone is known to be substantially excreted by the kidney, and the risk of adverse reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

8.6 Hepatic Impairment

In a study that used a single 4 mg oral dose of immediate-release hydromorphone tablets, four-fold increases in plasma levels of hydromorphone (C_{max} and $AUC_{0-\infty}$) were observed in patients with moderate hepatic impairment (Child-Pugh Group B). Start patients with moderate hepatic impairment on

25% of the hydromorphone hydrochloride extended-release tablets dose that would be used in patients with normal hepatic function. Closely monitor patients with moderate hepatic impairment for respiratory and central nervous system depression during initiation of therapy with hydromorphone hydrochloride extended-release tablets and during dose titration. The pharmacokinetics of hydromorphone in severe hepatic impairment patients have not been studied. As further increases in C_{\max} and $AUC_{0-\infty}$ of hydromorphone in this group are expected, use of alternate analgesics is recommended [see *Dosage and Administration (2.5)*].

8.7 Renal Impairment

Administration of a single 4 mg dose of immediate-release hydromorphone tablets resulted in two-fold and four-fold increases in plasma levels of hydromorphone (C_{\max} and AUC_{0-48h}) in moderate ($CL_{cr} = 40$ to 60 mL/min) and severe ($CL_{cr} < 30$ mL/min) impairment, respectively. In addition, in patients with severe renal impairment hydromorphone appeared to be more slowly eliminated with longer terminal elimination half-life. Start patients with moderate renal impairment on 50% and patients with severe renal impairment on 25% of the hydromorphone hydrochloride extended-release tablets dose that would be prescribed for patients with normal renal function. Closely monitor patients with renal impairment for respiratory and central nervous system depression during initiation of therapy with hydromorphone hydrochloride extended-release tablets and during dose titration. As hydromorphone hydrochloride extended-release tablets are only intended for once daily administration, consider use of an alternate analgesic that may permit more flexibility with the dosing interval in patients with severe renal impairment [see *Dosage and Administration (2.6)*].

9 DRUG ABUSE AND DEPENDENCE

9.1 Controlled Substance

Hydromorphone hydrochloride extended-release tablets contain hydromorphone, a Schedule II controlled substance.

9.2 Abuse

Hydromorphone hydrochloride extended-release tablets contain hydromorphone, a substance with a high potential for abuse similar to other opioids including fentanyl, hydrocodone, oxycodone, methadone, morphine, oxymorphone and tapentadol. Hydromorphone hydrochloride extended-release tablets can be abused and is subject to misuse, abuse, addiction, and criminal diversion [see *Warnings and Precautions (5.1)*].

The high drug content in extended-release formulations adds to the risk of adverse outcomes from abuse and misuse.

All patients treated with opioids require careful monitoring for signs of abuse and addiction, because use of opioid analgesic products carries the risk of addiction even under appropriate medical use.

Prescription drug abuse is the intentional non-therapeutic use of a prescription drug, even once, for its rewarding psychological or physiological effects.

Drug addiction is a cluster of behavioral, cognitive, and physiological phenomena that develop after repeated substance use and includes: a strong desire to take the drug, difficulties in controlling its use, persisting in its use despite harmful consequences, a higher priority given to drug use than to other activities and obligations, increased tolerance, and sometimes a physical withdrawal.

"Drug-seeking" behavior is very common in persons with substance use disorders. Drug-seeking tactics include emergency calls or visits near the end of office hours, refusal to undergo appropriate examination, testing or referral, repeated "loss" of prescriptions, tampering with prescriptions, and reluctance to provide prior medical records or contact information for other treating healthcare provider. "Doctor shopping" (visiting multiple prescribers to obtain additional prescriptions) is

common among drug abusers and people suffering from untreated addiction. Preoccupation with achieving pain relief can be appropriate behavior in a patient with poor pain control.

Abuse and addiction are separate and distinct from physical dependence and tolerance. Healthcare providers should be aware that addiction may not be accompanied by concurrent tolerance and symptoms of physical dependence in all addicts. In addition, abuse of opioids can occur in the absence of true addiction.

Hydromorphone hydrochloride extended-release tablets, like other opioids, can be diverted for non-medical use into illicit channels of distribution. Careful record-keeping of prescribing information, including quantity, frequency, and renewal requests, as required by state and federal law, is strongly advised.

Proper assessment of the patient, proper prescribing practices, periodic re-evaluation of therapy, and proper dispensing and storage are appropriate measures that help to limit abuse of opioid drugs.

Risks Specific to Abuse of Hydromorphone Hydrochloride Extended-Release Tablets

Hydromorphone hydrochloride extended-release tablets are for oral use only.

Abuse of hydromorphone hydrochloride extended-release tablets poses a risk of overdose and death. This risk is increased with concurrent abuse of hydromorphone hydrochloride extended-release tablets with alcohol and other central nervous system depressants. Taking cut, broken, chewed, crushed, or dissolved hydromorphone hydrochloride extended-release tablets enhances drug release and increases the risk of overdose and death.

With intravenous abuse, the inactive ingredients in hydromorphone hydrochloride extended-release tablets, may result in local tissue necrosis, infection, pulmonary granulomas, embolism and death, and increased risk of endocarditis and valvular heart injury. Parenteral drug abuse is commonly associated with transmission of infectious disease such as hepatitis and HIV.

9.3 Dependence

Both tolerance and physical dependence can develop during chronic opioid therapy. Tolerance is the need for increasing doses of opioids to maintain a defined effect such as analgesia (in the absence of disease progression or other external factors). Tolerance may occur to both the desired and undesired effects of drugs, and may develop at different rates for different effects.

Physical dependence results in withdrawal symptoms after abrupt discontinuation or a significant dose reduction of a drug. Withdrawal also may be precipitated through the administration of drugs with opioid antagonist activity, e.g., naloxone, nalmefene, mixed agonist/antagonist analgesics (e.g., pentazocine, butorphanol, nalbuphine), or partial agonists (e.g., buprenorphine). Physical dependence may not occur to a clinically significant degree until after several days to weeks of continued opioid usage.

Hydromorphone hydrochloride extended-release tablets should not be abruptly discontinued [*see Dosage and Administration (2.3)*]. If hydromorphone hydrochloride extended-release tablets are abruptly discontinued in a physically-dependent patient, an abstinence syndrome may occur. Some or all of the following can characterize this syndrome: restlessness, lacrimation, rhinorrhea, yawning, perspiration, chills, piloerection, myalgia, mydriasis, irritability, anxiety, backache, joint pain, weakness, abdominal cramps, insomnia, nausea, anorexia, vomiting, diarrhea, increased blood pressure, respiratory rate, or heart rate.

Infants born to mothers physically dependent on opioids will also be physically dependent and may exhibit respiratory difficulties and withdrawal signs [*see Warnings and Precautions (5.4)*].

10 OVERDOSAGE

Clinical Presentation

Acute overdosage with hydromorphone hydrochloride extended-release tablets can be manifested by respiratory depression, somnolence progressing to stupor or coma, skeletal muscle flaccidity, cold and clammy skin, constricted pupils, and, in some cases pulmonary edema, bradycardia, hypotension, partial or complete airway obstruction, atypical snoring, and death. Marked mydriasis rather than miosis may be seen with hypoxia in overdose situations.

Treatment of Overdose

In case of overdose, priorities are the re-establishment of a patent and protected airway and institution of assisted or controlled ventilation, if needed. Employ other supportive measures (including oxygen, vasopressors) in the management of circulatory shock and pulmonary edema as indicated. Cardiac arrest or arrhythmias will require advanced life support techniques.

The opioid antagonists, naloxone and nalmefene, are specific antidotes to respiratory depression resulting from opioid overdose. For clinically significant respiratory or circulatory depression secondary to hydromorphone overdose, administer an opioid antagonist. Opioid antagonists should not be administered in the absence of clinically significant respiratory or circulatory depression secondary to hydromorphone overdose.


Because the duration of reversal is expected to be less than the duration of action of hydromorphone in hydromorphone hydrochloride extended-release tablets, carefully monitor the patient until spontaneous respiration is reliably reestablished. Hydromorphone hydrochloride extended-release tablets will continue to release hydromorphone and add to the hydromorphone load for up to 24 to 48 hours or longer following ingestion, necessitating prolonged monitoring. If the response to opioid antagonist is suboptimal or only brief in nature, administer additional antagonist as directed by the product's prescribing information.

In an individual physically dependent on opioids, administration of the recommended usual dosage of the antagonist will precipitate an acute withdrawal syndrome. The severity of the withdrawal symptoms experienced will depend on the degree of physical dependence and the dose of the antagonist administered. If a decision is made to treat serious respiratory depression in the physically dependent patient, administration of the antagonist should be initiated with care and by titration with smaller than usual doses of the antagonist.

11 DESCRIPTION

Hydromorphone hydrochloride extended-release tablets are for oral use and contain hydromorphone hydrochloride, an opioid agonist.

Hydromorphone hydrochloride USP is 4,5 α -epoxy-3-hydroxy-17-methylmorphinan-6-one hydrochloride. Hydromorphone hydrochloride is a white or almost white crystalline powder that is freely soluble in water, very slightly soluble in ethanol (96%), and practically insoluble in methylene chloride. Its molecular formula is C₁₇H₁₉NO₃•HCl. The compound has the following structural formula:



Hydromorphone hydrochloride extended-release tablets also contain the following inactive ingredients: ammonio methacrylate copolymer, colloidal silicon dioxide, ethylcellulose, lactose monohydrate (8, 12, and 16 mg only), magnesium stearate, mannitol (32 mg only), microcrystalline cellulose, povidone, talc and triethyl citrate.

The color coating of the tablets contains FD&C Blue No.1, ferric oxide red (8, 16, and 32 mg only), ferric oxide yellow (8 and 16 mg only), hypromellose, polyethylene glycol, polysorbate 80 (12 mg only), propylene glycol, and titanium dioxide.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Hydromorphone, a semi-synthetic morphine derivative, is a hydrogenated ketone of morphine. Hydromorphone is a full opioid agonist and is relatively selective for the mu-opioid receptor, although it can bind to other opioid receptors at higher doses. The principal therapeutic action of hydromorphone is analgesia. Like all full opioid agonists, there is no ceiling effect for analgesia with morphine. Clinically, dosage is titrated to provide adequate analgesia and may be limited by adverse reactions, including respiratory and CNS depression.

The precise mechanism of the analgesic action is unknown. However, specific CNS opioid receptors for endogenous compounds with opioid-like activity have been identified throughout the brain and spinal cord and are thought to play a role in the analgesic effects of this drug.

12.2 Pharmacodynamics

CNS Depressant/Alcohol Interaction

Additive pharmacodynamic effects may be expected when hydromorphone hydrochloride extended-release tablets are used in conjunction with alcohol, other opioids, legal or illicit drugs that cause central nervous system depression.

Effects on the Central Nervous System

Hydromorphone produces dose-related respiratory depression by direct action on brain stem respiratory centers. The respiratory depression involves a reduction in the responsiveness of the brain stem respiratory centers to both increases in carbon dioxide tension and to electrical stimulation.

Hydromorphone causes miosis, even in total darkness. Pinpoint pupils are a sign of opioid overdose but are not pathognomic (e.g., pontine lesions of hemorrhagic or ischemic origins may produce similar

findings). Marked mydriasis, rather than miosis, may be seen due to severe hypoxia in overdose situations.

Effects on the Gastrointestinal Tract and Other Smooth Muscle

Hydromorphone causes a reduction in motility associated with an increase in tone in the antrum of the stomach and duodenum. Digestion of food in the small intestine is delayed and propulsive contractions are decreased. Propulsive peristaltic waves in the colon are decreased, while tone is increased to the point of spasm, resulting in constipation. Other opioid-induced effects may include a reduction in biliary and pancreatic secretions, spasm of sphincter of Oddi, and transient elevations in serum amylase.

Effects on the Cardiovascular System

Hydromorphone produces peripheral vasodilation which may result in orthostatic hypotension or syncope. Release of histamine may be induced by hydromorphone and can contribute to opioid-induced hypotension. Manifestations of histamine release or peripheral vasodilation may include pruritus, flushing, red eyes, sweating, and/or orthostatic hypotension.

Effects on the Endocrine System

Opioids inhibit the secretion of ACTH, cortisol, and luteinizing hormone (LH) in humans. They also stimulate prolactin, growth hormone (GH) secretion, and pancreatic secretion of insulin and glucagon.

Chronic use of opioids may influence the hypothalamic-pituitary-gonadal axis, leading to androgen deficiency that may manifest as low libido, impotence, erectile dysfunction, amenorrhea, or infertility. The causal role of opioids in the clinical syndrome of hypogonadism is unknown because the various medical, physical, lifestyle, and psychological stressors that may influence gonadal hormone levels have not been adequately controlled for in studies conducted to date [see *Adverse Reactions (6.2)*].

Effects on the Immune System

Opioids have been shown to have a variety of effects on components of the immune system in *in vitro* and animal models. The clinical significance of these findings is unknown. Overall, the effects of opioids appear to be modestly immunosuppressive.

Concentration -Efficacy Relationships

The minimum effective analgesic concentration will vary widely among patients, especially among patients who have been previously treated with potent agonist opioids. The minimum effective analgesic concentration of hydromorphone for any individual patient may increase over time due to an increase in pain, the development of a new pain syndrome, and/or the development of analgesic tolerance [see *Dosage and Administration (2.1, 2.3)*].

Concentration -Adverse Reaction Relationships

There is a relationship between increasing hydromorphone plasma concentration and increasing frequency of dose-related opioid adverse reactions such as nausea, vomiting, CNS effects, and respiratory depression. In opioid-tolerant patients, the situation may be altered by the development of tolerance to opioid-related adverse reactions [see *Dosage and Administration (2.1, 2.2, 2.3)*].

12.3 Pharmacokinetics

Absorption

Hydromorphone hydrochloride extended-release tablets are an extended-release formulation of hydromorphone that produces a gradual increase in hydromorphone concentrations. Following a single-dose administration of hydromorphone hydrochloride extended-release tablets, plasma concentrations gradually increase over 6 to 8 hours, and thereafter concentrations are sustained for approximately 18 to 24 hours post-dose. The median T_{max} values ranged from 12 to 16 hours. The mean half-life was approximately 11 hours, ranging from 8 to 15 hours in most individual subjects. Linear pharmacokinetics has been demonstrated for hydromorphone hydrochloride extended-release tablets over the dose range

8 to 64 mg, with a dose-proportional increase in C_{max} and overall exposure ($AUC_{0-\infty}$) (see **Table 5**). Steady-state plasma concentrations are approximately twice those observed following the first dose, and steady state is reached after 3 to 4 days of once-daily dosing of hydromorphone hydrochloride extended-release tablets. At steady state, hydromorphone hydrochloride extended-release tablets given once daily maintained hydromorphone plasma concentrations within the same concentration range as the immediate-release tablet given 4 times daily at the same total daily dose and diminished the fluctuations between peak and trough concentrations seen with the immediate-release tablet (see **Figure 1**). The bioavailability of hydromorphone hydrochloride extended-release tablets once daily and immediate-release hydromorphone four times daily in adults is comparable, as presented in **Table 5**.

Figure 1.

Mean Steady-State Plasma Concentration Profile




Table 5.

Mean (\pm SD) Hydromorphone Hydrochloride Extended-Release Tablets Pharmacokinetic Parameters

Regimen	Dosage	T_{max}^* (hrs)	C_{max} (ng/mL)	AUC (ng•hr/mL)	$T_{1/2}$ (hr)
Single Dose (N=31)	8 mg	12 (4-30)	0.93 (1.01)	18.1 (5.8)	10.6 (4.3)
	16 mg	16 (6-30)	1.69 (0.78)	36.5 (11.3)	10.3 (2.4)
	32 mg	16 (4-24)	3.25 (1.37)	72.2 (24.3)	11.0 (3.2)
	64 mg	16 (6-30)	6.61 (1.75)	156.0 (30.6)	10.9 (3.8)
Multiple Dose† (N=29)	16 mg q24h	12 (6-24)	3.54 (0.96)‡	57.6 (16.3)	NA
	IR 4 mg q6h	0.75 (0.5-2)	5.28 (1.37)§	54.8 (14.8)	NA

NA = not applicable

* Median (range) reported for T_{max}

† Steady-state results on Day 5 (0-24 hours)

‡ C_{min} 2.15 (0.87) ng/mL

§ C_{min} 1.47 (0.42) ng/mL

Food Effect

The pharmacokinetics of hydromorphone hydrochloride extended-release tablets are not affected by food as indicated by bioequivalence when administered under fed and fasting conditions. Therefore, hydromorphone hydrochloride extended-release tablets may be administered without regard to meals. When a 16 mg dose of hydromorphone hydrochloride extended-release tablets was administered to healthy volunteers immediately following a high-fat meal, the median time to C_{max} (T_{max}) was minimally affected by the high-fat meal occurring at 16 hours compared to 18 hours while fasting.

Distribution

Following intravenous administration of hydromorphone to healthy volunteers, the mean volume of distribution was 2.9 (\pm 1.3) L/kg, suggesting extensive tissue distribution. The mean extent of binding of hydromorphone to human plasma proteins was determined to be 27% in an *in vitro* study.

Elimination

Metabolism

After oral administration of an immediate-release formulation, hydromorphone undergoes extensive first-pass metabolism and is metabolized primarily in the liver by glucuronidation to hydromorphone-3-glucuronide, which follows a similar time course to hydromorphone in plasma. Exposure to the glucuronide metabolite is 35 to 40 times higher than exposure to the parent drug. *In vitro* data suggest that hydromorphone in clinically relevant concentrations has minimal potential to inhibit the activity of human hepatic CYP450 enzymes including CYP1A2, 2C9, 2C19, 2D6, 3A4, and 4A11.

Excretion

Approximately 75% of the administered dose is excreted in urine. Most of the administered hydromorphone dose is excreted as metabolites. Approximately 7% and 1% of the dose are excreted as unchanged hydromorphone in urine and feces, respectively.

Specific Populations

Age: Geriatric Patients

Population PK analysis performed on plasma concentration data from 407 osteoarthritis (OA) patients using hydromorphone hydrochloride extended-release tablets showed an average 11% increase in hydromorphone AUC in the elderly group (65 to 75 years of age) when compared to the younger age group (less than or equal to 65 years of age).

Sex

Females appeared to have approximately 10% higher mean systemic exposure in terms of C_{max} and AUC values.

Hepatic Impairment

In a study that used a single 4 mg oral dose of immediate-release hydromorphone tablets, four-fold increases in plasma levels of hydromorphone (C_{max} and $AUC_{0-\infty}$) were observed in patients with moderate hepatic impairment (Child-Pugh Group B). Pharmacokinetics of hydromorphone in severe hepatic impairment patients has not been studied. Further increase in C_{max} and $AUC_{0-\infty}$ of hydromorphone in this group is expected. Start patients with moderate hepatic impairment on 25% of the

usual dose of hydromorphone hydrochloride extended-release tablets and closely monitor for respiratory and central nervous system depression during dose titration. Consider alternate analgesic therapy for patients with severe hepatic impairment [see *Dosage and Administration (2.5) and Specific Populations (8.6)*].

Renal Impairment

Renal impairment affected the pharmacokinetics of hydromorphone and its metabolites following administration of a single 4 mg dose of immediate-release tablets. The effects of renal impairment on hydromorphone pharmacokinetics were two-fold and four-fold increases in plasma levels of hydromorphone (C_{max} and AUC_{0-48h}) in moderate ($CL_{cr} = 40$ to 60 mL/min) and severe ($CL_{cr} < 30$ mL/min) impairment, respectively. In addition, in patients with severe renal impairment hydromorphone appeared to be more slowly eliminated with longer terminal elimination half-life (40 hr) compared to subjects with normal renal function (15 hr). Start patients with moderate renal impairment on 50% of the usual hydromorphone hydrochloride extended-release tablets dose for patients with normal renal function and closely monitor for respiratory and central nervous system depression during dose titration. As hydromorphone hydrochloride extended-release tablets are only intended for once-daily administration, consider use of an alternate analgesic that may permit more flexibility with the dosing interval in patients with severe renal impairment [see *Dosage and Administration (2.6) and Use in Specific Populations (8.7)*].

Drug Interaction Studies

Alcohol Interaction

An *in vivo* study examined the effect of alcohol (40%, 20%, 4% and 0%) on the bioavailability of a single dose of 16 mg of hydromorphone hydrochloride extended-release tablets in healthy, fasted or fed volunteers. The results showed that the hydromorphone mean $AUC_{0-\infty}$ was 5% higher and 4% lower (not statistically significant) in the fasted and fed groups respectively after co-administration of 240 mL of 40% alcohol. The $AUC_{0-\infty}$ was similarly unaffected in subjects following the co-administration of hydromorphone hydrochloride extended-release tablets and alcohol (240 mL of 20% or 4% alcohol).

The change in geometric mean C_{max} with concomitant administration of alcohol and hydromorphone hydrochloride extended-release tablets ranged from an increase of 10% to 31% across all conditions studied. The change in mean C_{max} was greater in the fasted group of subjects. Following concomitant administration of 240 mL of 40% alcohol while fasting, the mean C_{max} increased by 37% and up to 151% in an individual subject. Following the concomitant administration of 240 mL of 20% alcohol while fasting, the mean C_{max} increased by 35% and up to 139% in an individual subject. Following the concomitant administration of 240 mL of 4% alcohol while fasting, the mean C_{max} increased by 19% on average and as much as 73% for an individual subject. The range of median T_{max} for the fed and fasted treatments with 4%, 20% and 40% alcohol was 12 to 16 hours compared to 16 hours for the 0% alcohol treatments.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Long-term studies to evaluate the carcinogenic potential of hydromorphone hydrochloride were completed in both Han-Wistar rats and Crl:CD1® (ICR) mice. Hydromorphone HCl was administered to Han-Wistar rats (2, 5, and 15 mg/kg/day for males, and 8, 25 and 75 mg/kg/day for females) for 2 years by oral gavage. In female rats, incidences of hibernoma (tumor of brown fat) were increased at 10.5 times the maximum recommended daily exposure based on AUC at the mid dose (2 tumor, 25 mg/kg/day) and 53.7 times the maximum recommended human daily exposure based on AUC at the maximum dose (4 tumors, 75 mg/kg/day). The clinical relevance of this finding to humans has not been established. There

was no evidence of carcinogenicity in male rats. The systemic drug exposure (AUC, ng·h/mL) at the 15 mg/kg/day in male rats was 7.6 times greater than the human exposure at a single dose of 32 mg/day of hydromorphone hydrochloride extended-release tablets. There was no evidence of carcinogenic potential in CrI:CD1® (ICR) mice administered hydromorphone HCl at doses up to 15 mg/kg/day for 2 years by oral gavage. The systemic drug exposure (AUC, ng·h/mL) at the 15 mg/kg/day in mice was 1.1 (in males) and 1.2 (in females) times greater than the human exposure at a single dose of 32 mg/day of hydromorphone hydrochloride extended-release tablets.

Mutagenesis

Hydromorphone was not mutagenic in the *in vitro* bacterial reverse mutation assay (Ames assay). Hydromorphone was not clastogenic in either the *in vitro* human lymphocyte chromosome aberration assay or the *in vivo* mouse micronucleus assay.

Impairment of Fertility

Reduced implantation sites and viable fetuses were noted at 2.1 times the human daily dose of 32 mg/day in a study in which female rats were treated orally with 1.75, 3.5, or 7 mg/kg/day hydromorphone hydrochloride (0.7, 1.4, or 2.8 times a human daily dose of 24 mg/day (HDD) based on body surface area) beginning 14 days prior to mating through Gestation Day 7 and male rats were treated with the same hydromorphone hydrochloride doses beginning 28 days prior to and throughout mating.

14 CLINICAL STUDIES


Hydromorphone hydrochloride extended-release tablets were investigated in a double-blind, placebo-controlled, randomized withdrawal study in opioid tolerant patients with moderate-to-severe low back pain. Patients were considered opioid tolerant if they were currently on opioid therapy that was ≥ 60 mg/day of oral morphine equivalent for at least 2 months prior to screening. Patients entered an open-label conversion and titration phase with hydromorphone hydrochloride extended-release tablets, were converted to a starting dose that was approximately 75% of their total daily morphine equivalent dose, and were dosed once daily until adequate pain control was achieved while exhibiting tolerable side effects. Supplemental immediate-release hydromorphone tablets were allowed throughout the study. Patients who achieved a stable dose entered a 12-week, double-blind, placebo-controlled, randomized treatment phase. Mean daily dose at randomization was 37.8 mg/day (range of 12 mg/day to 64 mg/day). Fifty-eight (58) percent of patients were successfully titrated to a stable dose of hydromorphone hydrochloride extended-release tablets during the open-label conversion and titration phase.

During the double-blind treatment phase, patients randomized to hydromorphone hydrochloride extended-release tablets continued with the stable dose achieved in the conversion and titration phase of the study. Patients randomized to placebo received, in a blinded manner, hydromorphone hydrochloride extended-release tablets and matching placebo in doses tapering from the stable dose achieved in conversion and titration. During the taper down period, patients were allowed immediate-release hydromorphone tablets as supplemental analgesia to minimize opioid withdrawal symptoms in placebo patients. After the taper period, the number of immediate-release hydromorphone tablets was limited to two tablets per day. Forty-nine (49) percent of patients treated with hydromorphone hydrochloride extended-release tablets and 33% of patients treated with placebo completed the 12-week treatment period.

Hydromorphone hydrochloride extended-release tablets provided superior analgesia compared to placebo. There was a significant difference between the mean changes from Baseline to Week 12 or Final Visit in average weekly pain intensity Numeric Rating Scale (NRS) scores obtained from patient diaries between the two groups. The proportion of patients with various degrees of improvement from screening to Week 12 or Final Visit is shown in **Figure 2**. For this analysis, patients who discontinued treatment for any reason prior to Week 12 were assigned a value of zero improvement.

Figure 2.

Percent Reduction in Average Pain Intensity from Screening to Week 12 or Final Visit



16 HOW SUPPLIED/STORAGE AND HANDLING

How Supplied

Hydromorphone Hydrochloride Extended-Release Tablet Strengths				
Strength	Color	Tablet Description	Bottle Count	NDC
8 mg	Dark Beige	Round, biconvex, imprinted with "P293" over "8" on one side	100	0574-0293-01
12 mg	White	Round, biconvex, imprinted with "P294" over "12" on one side	100	0574-0294-01
16 mg	Light Beige	Round, biconvex, imprinted with "P295" over "16" on one side	100	0574-0295-01
32 mg	Pink	Round, biconvex, imprinted with "P297" over "32" on one side	100	0574-0297-01

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide)

Addiction, Abuse, and Misuse

Inform patients that the use of hydromorphone hydrochloride extended-release tablets, even when taken as recommended, can result in addiction, abuse, and misuse, which can lead to overdose or death [see *Warnings and Precautions (5.1)*]. Instruct patients not to share hydromorphone hydrochloride extended-release tablets with others and to take steps to protect hydromorphone hydrochloride extended-release tablets from theft or misuse.

Life-threatening Respiratory Depression

Inform patients of the risk of life-threatening respiratory depression, including information that the risk is greatest when starting hydromorphone hydrochloride extended-release tablets or when the dose is increased, and that it can occur even at recommended doses [see *Warnings and Precautions (5.3)*]. Advise

patients how to recognize respiratory depression and to seek medical attention if breathing difficulties develop.

Accidental Ingestion

Inform patients that accidental ingestion, especially by children, may result in respiratory depression or death [see *Warnings and Precautions (5.3)*]. Instruct patients to take steps to store hydromorphone hydrochloride extended-release tablets securely and to dispose of unused hydromorphone hydrochloride extended-release tablets by flushing the tablets down the toilet.

Interactions with Benzodiazepines and Other CNS Depressants

Inform patients and caregivers that potentially fatal additive effects may occur if hydromorphone hydrochloride extended-release tablets are used with benzodiazepines or other CNS depressants, including alcohol, and not to use these concomitantly unless supervised by a health care provider [see *Warnings and Precautions (5.5)*, *Drug Interactions (7)*].

Serotonin Syndrome

Inform patients that hydromorphone hydrochloride extended-release tablets could cause a rare but potentially life-threatening condition resulting from concomitant administration of serotonergic drugs. Warn patients of the symptoms of serotonin syndrome and to seek medical attention right away if symptoms develop. Instruct patients to inform their healthcare providers if they are taking, or plan to take serotonergic medications [see *Drug Interactions 7*].

MAOI Interaction

Inform patients to avoid taking hydromorphone hydrochloride extended-release tablets while using any drugs that inhibit monoamine oxidase. Patients should not start MAOIs while taking hydromorphone hydrochloride extended release tablets [see *Drug Interactions 7*].

Adrenal Insufficiency

Inform patients that hydromorphone hydrochloride extended-release tablets could cause adrenal insufficiency, a potentially life-threatening condition. Adrenal insufficiency may present with non-specific symptoms and signs such as nausea, vomiting, anorexia, fatigue, weakness, dizziness, and low blood pressure. Advise patients to seek medical attention if they experience a constellation of these symptoms [see *Warnings and Precautions (5.7)*].

Important Administration Instructions

Instruct patients how to properly take hydromorphone hydrochloride extended-release tablets, including the following:

- Hydromorphone hydrochloride extended-release tablets are designed to work properly only if swallowed intact. Taking cut, broken, chewed, crushed, or dissolved hydromorphone hydrochloride extended-release tablets can result in a fatal overdose [see *Dosage and Administration (2.1)*].
- Using hydromorphone hydrochloride extended-release tablets exactly as prescribed to reduce the risk of life-threatening adverse reactions (e.g., respiratory depression)
- Do not discontinue hydromorphone hydrochloride extended-release tablets without first discussing the need for a tapering regimen with the prescriber [see *Dosage and Administration (2.4)*].

Gastrointestinal Blockage

Advise patients that people with certain stomach or intestinal problems such as narrowing of the intestines or previous surgery may be at higher risk of developing a blockage. Symptoms include abdominal distension, abdominal pain, severe constipation, or vomiting. Instruct patients to contact their healthcare provider immediately if they develop these symptoms.

Hypotension

Inform patients that hydromorphone hydrochloride extended-release tablets may cause orthostatic hypotension and syncope. Instruct patients how to recognize symptoms of low blood pressure and how to reduce the risk of serious consequences should hypotension occur (e.g., sit or lie down, carefully rise from a sitting or lying position).

Anaphylaxis

Inform patients that anaphylaxis has been reported with ingredients contained in hydromorphone hydrochloride extended-release tablets. Advise patients how to recognize such a reaction and when to seek medical attention [see *Contraindications (4)*, *Warnings and Precautions (5.13)*, and *Adverse Reactions (6.2)*].

Pregnancy

Neonatal Opioid Withdrawal Syndrome

Inform female patients of reproductive potential that prolonged use of hydromorphone hydrochloride extended-release tablets during pregnancy can result in neonatal opioid withdrawal syndrome, which may be life-threatening if not recognized and treated [see *Warnings and Precautions (5.4)*, *Use in Specific Populations (8.1)*].

Embryo-Fetal Toxicity

Inform female patients of reproductive potential that hydromorphone hydrochloride extended-release tablets can cause fetal harm and to inform their healthcare provider of a known or suspected pregnancy [see *Use in Specific Populations (8.1)*].

Lactation

Advise patients that breastfeeding is not recommended during treatment with hydromorphone hydrochloride extended-release tablets [see *Use in Specific Populations (8.2)*].

Infertility

Inform patients that chronic use of opioids may cause reduced fertility. It is not known whether these effects on fertility are reversible [see *Use in Specific Populations (8.3)*].

Driving or Operating Heavy Machinery

Inform patients that hydromorphone hydrochloride extended-release tablets may impair the ability to perform potentially hazardous activities such as driving a car or operating heavy machinery. Advise patients not to perform such tasks until they know how they will react to the medication [see *Warnings and Precautions (5.14)*].

Constipation

Advise patients of the potential for severe constipation, including management instructions and when to seek medical attention [see *Adverse Reactions (6.1)*, *Clinical Pharmacology (12.2)*].

Disposal Of Unused Hydromorphone Hydrochloride Extended-Release Tablets

Advise patients to flush the unused tablets down the toilet when hydromorphone hydrochloride extended-release tablets are no longer needed.

Manufactured By

Perrigo®

Minneapolis, MN 55427

2203457

1B500 RC J12

Rev 10-18 D

PHARMACIST: Dispense a separate Medication Guide to each patient. Medication Guides available at: www.perrigo.com or call 1-866-634-9120

Medication Guide

Hydromorphone Hydrochloride (hy-druh-mor-fon hy-druh-klawr-ahyd) Extended-Release Tablets, CII

Hydromorphone hydrochloride extended-release tablets are:

- A strong prescription pain medicine that contains an opioid (narcotic) that is used to manage pain severe enough to require daily around-the-clock, long-term treatment with an opioid, when other pain treatments such as non-opioid pain medicines or immediate-release opioid medicines do not treat your pain well enough or you cannot tolerate them.
- A long-acting (extended-release) opioid pain medicine that can put you at risk for overdose and death. Even if you take your dose correctly as prescribed you are at risk for opioid addiction, abuse, and misuse that can lead to death.
- Not for use to treat pain that is not around-the-clock.

Important information about hydromorphone hydrochloride extended-release tablets:

- **Get emergency help right away if you take too many hydromorphone hydrochloride extended-release tablets (overdose).** When you first start taking hydromorphone hydrochloride extended-release tablets, when your dose is changed, or if you take too much (overdose), serious or life threatening breathing problems that can lead to death may occur.
- Taking hydromorphone hydrochloride extended-release tablets with other opioid medicines, benzodiazepines, alcohol, or other central nervous system depressants (including street drugs) can cause severe drowsiness, decreased awareness, breathing problems, coma, and death.
- Never give anyone else your hydromorphone hydrochloride extended-release tablets. They could die from taking it. Store hydromorphone hydrochloride extended-release tablets away from children and in a safe place to prevent stealing or abuse. Selling or giving away hydromorphone hydrochloride extended-release tablets is against the law.

Do not take hydromorphone hydrochloride extended-release tablets if you have:

- severe asthma, trouble breathing, or other lung problems.
- a bowel blockage or have narrowing of the stomach or intestines.

Before taking hydromorphone hydrochloride extended-release tablets, tell your healthcare provider if you have a history of:

- head injury, seizures

- liver, kidney, thyroid problems
- allergy to sulfites
- problems urinating
- pancreas or gallbladder problems
- abuse of street or prescription drugs, alcohol addiction, or mental health problems.

Tell your healthcare provider if you are:

- **pregnant or planning to become pregnant.** Prolonged use of hydromorphone hydrochloride extended-release tablets during pregnancy can cause withdrawal symptoms in your newborn baby that could be life-threatening if not recognized and treated.
- **breastfeeding.** Not recommended during treatment with hydromorphone hydrochloride extended-release tablets. It may harm your baby.
- taking prescription or over-the-counter medicines, vitamins, or herbal supplements. Taking hydromorphone hydrochloride extended-release tablets with certain other medicines can cause serious side effects.

When taking hydromorphone hydrochloride extended-release tablets:

- Do not change your dose. Take hydromorphone hydrochloride extended-release tablets exactly as prescribed by your healthcare provider. Use the lowest dose possible for the shortest time needed.
- Take your prescribed dose every 24 hours, at the same time every day. Do not take more than your prescribed dose in 24 hours. If you miss a dose, take your next dose at your usual time the next day.
- Swallow hydromorphone hydrochloride extended-release tablets whole. Do not cut, break, chew, crush, dissolve, snort, or inject hydromorphone hydrochloride extended-release tablets because this may cause you to overdose and die.
- **Call your healthcare provider if the dose you are taking does not control your pain.**
- **Do not stop taking hydromorphone hydrochloride extended-release tablets without talking to your healthcare provider.**
- Hydromorphone hydrochloride extended-release tablets are contained in a hard tablet shell that you may see in your bowel movement; this is normal.
- After you stop taking hydromorphone hydrochloride extended-release tablets, flush any unused tablets down the toilet.

While taking hydromorphone hydrochloride extended-release tablets, DO NOT:

- Drive or operate heavy machinery, until you know how hydromorphone hydrochloride extended release tablets affect you. Hydromorphone hydrochloride extended-release tablets can make you sleepy, dizzy, or lightheaded.
- Drink alcohol or use prescription or over-the-counter medicines containing alcohol. Using products containing alcohol during treatment with hydromorphone hydrochloride extended-release tablets may cause you to overdose and die.

The possible side effects of hydromorphone hydrochloride extended-release tablets are:

- constipation, nausea, sleepiness, vomiting, tiredness, headache, dizziness, abdominal pain.

Call your healthcare provider if you have any of these symptoms and they are severe.

Get emergency medical help if you have:

- trouble breathing, shortness of breath, fast heartbeat, chest pain, swelling of your face, tongue, or throat, extreme drowsiness, light-headedness when changing positions, feeling faint, agitation, high body temperature, trouble walking, stiff muscles, or mental changes such as confusion.

These are not all the possible side effects of hydromorphone hydrochloride extended-release tablets. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. **For more information go to dailymed.nlm.nih.gov**

Manufactured By

Perrigo®

Minneapolis, MN 55427

2203068 1B500 RC IC3 Rev 11-17 C

www.perrigo.com or call 1-866-634-9120

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Revised November 2017

PACKAGE/LABEL PRINCIPAL DISPLAY PANEL – 8 mg

Rx Only

NDC 0574-0293-01

Once Daily

HYDROmorphone HCl Extended-Release Tablets 8 mg

CII

For opioid tolerant patients only

Dispense the accompanying Medication Guide to each patient

100 Tablets

Rx Only

NDC 0574-0293-01

Once Daily

HYDROmorphone HCl

Extended-Release 

Tablets 8 mg

For opioid tolerant patients only

Dispense the accompanying Medication Guide to each patient

100 Tablets



Each tablet contains:
Hydromorphone Hydrochloride USP, 8 mg

Swallow tablets whole. Do not break, crush or chew tablet.

See package insert for dosing information.

Store at 20° to 25°C (68° to 77°F)
[see USP Controlled Room Temperature].

Manufactured By Perrigo
Minneapolis, MN 55427
2203633 1B578 RC F3 Rev 05-17 A

GTIN 00305740293010

Lot # / Exp date area
No Varnish

N 3 0574-0293-01 0

The following image is a placeholder representing the product identifier that is either affixed or imprinted on the drug package label during the packaging operation.

S/N [insert product's serial number]
Lot [insert product's lot number]
Exp [insert product's expiration date]

PACKAGE/LABEL PRINCIPAL DISPLAY PANEL – 12 mg

0574-0294-01

Once Daily

HYDROmorphone HCl Extended-Release Tablets 12 mg

CII

For opioid tolerant patients only

Dispense the accompanying Medication Guide to each patient

100 Tablets

Rx Only



The following image is a placeholder representing the product identifier that is either affixed or imprinted on the drug package label during the packaging operation.

S/N [insert product's serial number]
Lot [insert product's lot number]
Exp [insert product's expiration date]

PACKAGE/LABEL PRINCIPAL DISPLAY PANEL – 16 mg

NDC 0574-0295-01

Once Daily

HYDROmorphone HCl Extended-Release Tablets 16 mg

CII

For opioid tolerant patients only


Dispense the accompanying Medication Guide to each patient

100 Tablets

Rx Only



The following image is a placeholder representing the product identifier that is either affixed or imprinted on the drug package label during the packaging operation.



Package/Label Display Panel – 32 mg

NDC 0574-0297-01

Once Daily

HYDROmorphone HCl Extended-Release Tablets 32 mg

CII

For opioid tolerant patients only

Dispense the accompanying Medication Guide to each patient

100 Tablets

Rx Only

Rx Only

NDC 0574-0297-01
Once Daily
HYDROMORPHONE HCl
Extended-Release **Tablets 32 mg** **Ⓜ**

For opioid tolerant patients only

Dispense the accompanying Medication Guide to each patient

100 Tablets **Perrigo**

Each tablet contains:
Hydromorphone Hydrochloride USP, 32 mg
Swallow tablets whole. Do not break, crush or chew tablet.
See package insert for dosing information.
Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

Manufactured By Perrigo
Minneapolis, MN 55427
2203630 5P178 RC F2 Rev 05-17 A

GTIN 00305740297018

3 0574-0297-01 8

Lot # / Exp date area
No Varnish

The following image is a placeholder representing the product identifier that is either affixed or imprinted on the drug package label during the packaging operation.

S/N [insert product's serial number]
Lot [insert product's lot number]
Exp [insert product's expiration date]

HYDROMORPHONE HYDROCHLORIDE

hydromorphone hydrochloride tablet, extended release

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0574-0293
Route of Administration	ORAL	DEA Schedule	CII

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HYDROMORPHONE HYDROCHLORIDE (UNII: L960UP2KRW) (HYDROMORPHONE - UNII:Q812464R06)	HYDROMORPHONE HYDROCHLORIDE	8 mg

Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

MAGNESIUM STEARATE (UNII: 70097M6I30)
ETHYLCELLULOSE (100 MPA.S) (UNII: 47MLB0F1MV)
POVIDONE K30 (UNII: U725QWY32X)
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)
AMMONIO METHACRYLATE COPOLYMER TYPE A (UNII: 8GQS4E66YY)
TALC (UNII: 7SEV7J4R1U)
HYPROMELLOSE 2910 (3 MPA.S) (UNII: 0VUT3PMY82)
HYPROMELLOSE 2910 (6 MPA.S) (UNII: 0WZ8WG20P6)
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)
FERRIC OXIDE RED (UNII: 1K09F3G675)

Product Characteristics

Color	BROWN (Dark Beige)	Score	no score
Shape	ROUND (biconvex)	Size	7mm
Flavor		Imprint Code	P293;8
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0574-0293-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	05/22/2015	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA204278	05/22/2015	

HYDROMORPHONE HYDROCHLORIDE

hydromorphone hydrochloride tablet, extended release

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0574-0294
Route of Administration	ORAL	DEA Schedule	CII

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HYDROMORPHONE HYDROCHLORIDE (UNII: L960UP2KRW) (HYDROMORPHONE - UNII:Q812464R06)	HYDROMORPHONE HYDROCHLORIDE	12 mg

Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
ETHYLCELLULOSE (100 MPA.S) (UNII: 47MLB0F1MV)	
POVIDONE K30 (UNII: U725QWY32X)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
AMMONIO METHACRYLATE COPOLYMER TYPE A (UNII: 8GQS4E66YY)	
TALC (UNII: 7SEV7J4R1U)	
HYPROMELLOSE 2910 (3 MPA.S) (UNII: 0VUT3PMY82)	
HYPROMELLOSE 2910 (6 MPA.S) (UNII: 0WZ8WG20P6)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	

Product Characteristics

Color	WHITE	Score	no score
Shape	ROUND (biconvex)	Size	8mm
Flavor		Imprint Code	P294;12
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0574-0294-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	05/22/2015	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA204278	05/22/2015	

HYDROMORPHONE HYDROCHLORIDE

hydromorphone hydrochloride tablet, extended release

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0574-0295
Route of Administration	ORAL	DEA Schedule	CII

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

HYDROMORPHONE HYDROCHLORIDE (UNII: L960UP2KRW) (HYDROMORPHONE - UNII:Q812464R06)	HYDROMORPHONE HYDROCHLORIDE	16 mg
---	--------------------------------	-------

Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
ETHYLCELLULOSE (100 MPA.S) (UNII: 47MLB0F1MV)	
POVIDONE K30 (UNII: U725QWY32X)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
AMMONIO METHACRYLATE COPOLYMER TYPE A (UNII: 8GQS4E66YY)	
TALC (UNII: 7SEV7J4R1U)	
HYPROMELLOSE 2910 (3 MPA.S) (UNII: 0VUT3PMY82)	
HYPROMELLOSE 2910 (6 MPA.S) (UNII: 0WZ8WG20P6)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	

Product Characteristics

Color	BROWN (Light Beige)	Score	no score
Shape	ROUND (biconvex)	Size	9mm
Flavor		Imprint Code	P295;16
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0574-0295-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	05/22/2015	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA204278	05/22/2015	

HYDROMORPHONE HYDROCHLORIDE

hydromorphone hydrochloride tablet, extended release

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0574-0297
--------------	-------------------------	--------------------	---------------

Route of Administration

ORAL

DEA Schedule

CII

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HYDROMORPHONE HYDROCHLORIDE (UNII: L960UP2KRW) (HYDROMORPHONE - UNII:Q812464R06)	HYDROMORPHONE HYDROCHLORIDE	32 mg

Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
MANNITOL (UNII: 3OWL53L36A)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
ETHYLCELLULOSE (100 MPA.S) (UNII: 47MLB0F1MV)	
POVIDONE K30 (UNII: U725QWY32X)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
AMMONIO METHACRYLATE COPOLYMER TYPE A (UNII: 8GQS4E66YY)	
TALC (UNII: 7SEV7J4R1U)	
HYPROMELLOSE 2910 (3 MPA.S) (UNII: 0VUT3PMY82)	
HYPROMELLOSE 2910 (6 MPA.S) (UNII: 0WZ8WG20P6)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	

Product Characteristics

Color	PINK	Score	no score
Shape	ROUND (biconvex)	Size	9mm
Flavor		Imprint Code	P297;32
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0574-0297-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	09/29/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA204278	05/22/2015	

Labeler - Paddock Laboratories, LLC (967694121)

