

**COLD PLUS HEAD CONGESTION NIGHTTIME- acetaminophen,
chlorpheniramine maleate, dextromethorphan hydrobromide, and
phenylephrine hydrochloride tablet, coated
TopCo Associates LLC**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

TCR - 1161 - 2019-1016

Drug Facts

Active ingredients (in each caplet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Chlorpheniramine maleate 2 mg	Antihistamine
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- for the temporary relief of the following cold symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - runny nose and sneezing
 - cough
 - sinus congestion and pressure
- helps clear nasal passages
- relieves cough to help you sleep
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- high blood pressure
- heart disease
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days

- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)

adults and children 12 years and over	<ul style="list-style-type: none"> ▪ take 2 caplets every 4 hours ▪ swallow whole – do not crush, chew, or dissolve ▪ do not take more than 10 caplets in 24 hours
children under 12 years	<ul style="list-style-type: none"> ▪ ask a doctor

Other information

- store between 20-25C (68°-77°F) in a dry place
- retain carton for complete product information and warnings

Inactive ingredients

acesulfame potassium, colloidal silicon dioxide, croscarmellose sodium, FD&C blue #1 aluminum lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide

PRINCIPAL DISPLAY PANEL

NDC 36800-925-02

TopCare® Health

NIGHTTIME

Cold + Head Congestion Nighttime

ACETAMINOPHEN, DEXTROMETHORPHAN HBr, PHENYLEPHRINE HCl,
CHLORPHENIRAMINE MALEATE

PAIN RELIEVER/FEVER REDUCER, COUGH SUPPRESSANT, NASAL CONGESTION,
ANTIHISTAMINE

Relief of:

- Headache + Body Aches
- Fever + Sore Throat
- Cough
- Nasal Congestion
- Runny Nose

24 COOL TASTE CAPLETS

NC

NC

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are:

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

■ do not exceed recommended dosage

■ excitability may occur, especially in children

■ marked drowsiness may occur

■ avoid alcoholic drinks

■ avoid sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

■ be careful when driving a motor vehicle or operating machinery

■ pain, nasal congestion, or cough gets worse or lasts more than 7 days

■ fever gets worse or lasts more than 3 days

■ redness or swelling is present

■ new symptoms occur

■ cough comes back or occurs with rash or headache that lasts

■ These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warnings: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ do not take more than directed (see overdose warning)

■ take 2 caplets every 4 hours

■ swallow whole; do not crush, chew, or dissolve

■ do not take more than 10 caplets in 24 hours

■ ask a doctor

Adults and children 12 years and over

Children under 12 years

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are:

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis

Other information

■ store between 20°-25° (68°-77°) in a dry place

■ retain carton for complete product information and warnings

Inactive ingredients

aspartame, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, croscarmellose sodium, croscopolone, FD&C blue #1

Drug Facts (continued)

aluminum lake, fluor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, croscarmellose sodium, croscopolone, FD&C blue #1

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are:

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis

Other information

■ store between 20°-25° (68°-77°) in a dry place

■ retain carton for complete product information and warnings

Inactive ingredients

aspartame, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, croscarmellose sodium, croscopolone, FD&C blue #1

Drug Facts (continued)

aluminum lake, fluor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, croscarmellose sodium, croscopolone, FD&C blue #1

TopCare[®] health

NIGHTTIME

Cold + Head Congestion Nighttime

For Adults

TopCare[®] health

NIGHTTIME

Cold + Head Congestion Nighttime

ACETAMINOPHEN, DEXTROMETHORPHAN HBr, PHENYLEPHRINE HCl, CHLORPHENIRAMINE MALEATE
PAIN RELIEVER/FEVER REDUCER, COUGH SUPPRESSANT, NASAL DECONGESTANT, ANTIHISTAMINE

RELIEF OF:

- Headache + Body Aches
- Fever + Sore Throat
- Cough
- Nasal Congestion
- Runny Nose

For Adults

24 COOL TASTE CAPLETS

actual size

F116102TCR_R2



0 36800 11802 7

Drug Facts

Active ingredients (in each caplet)

Purpose

Acetaminophen 325 mg.....Pain reliever/fever reducer

Chlorpheniramine maleate 2 mg.....Antihistamine

Dextromethorphan HBr 10 mg.....Cough suppressant

Phenylephrine HCl 5 mg.....Nasal decongestant

Drug Facts (continued)

Uses

■ for the temporary relief of the following cold symptoms:

- headache
- sore throat
- minor aches and pains
- nasal congestion
- runny nose and sneezing
- sinus congestion and pressure
- helps clear nasal passages
- relieves cough to help you sleep
- temporarily reduces fever

COLD PLUS HEAD CONGESTION NIGHTTIME

acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, and phenylephrine hydrochloride tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-925
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1VS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	blue	Score	no score
Shape	OVAL	Size	17mm
Flavor	MINT	Imprint Code	AAA;1139
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-925-02	2 in 1 CARTON	08/30/2011	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	08/30/2011	

Labeler - TopCo Associates LLC (006935977)

Revised: 11/2021

TopCo Associates LLC