DAYTIME NIGHTTIME COLD AND FLU RELIEF MULTI-SYMPTOMacetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl

Family Dollar (FAMILY WELLNESS)

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredients in Daytime (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

Active ingredients in Nighttime (in each softgel)

Acetaminophen 325 mg

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

Purpose for Daytime

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Purpose for Nighttime

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

DAYTIME

- temporarily relieves common cold and flu symptoms:
 - cough due to minor throat and bronchial irritation
 - nasal congestion
 - headache

- minor aches and pain
- fever
- sore throat

NIGHTTIME

- temporarily relieves common cold and flu symptoms:
 - cough due to minor throat and bronchial irritation
 - sore throat
 - headache
 - minor aches and pain
 - fever
 - runny nose & sneezing

Warnings

Liver warning: These products contain acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

DAYTIME

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure

- cough that occurs with too much phlegm (mucus)
- trouble urinating due to an enlarged prostate gland
- · persistent or chronic cough such as occurs with smoking, asthma, or emphysema

NIGHTTIME

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

DAYTIME

taking the blood thinning drug warfarin.

NIGHTTIME

- taking the blood thinning drug warfarin
- taking sedative or tranquilizers

When using this product,

DAYTIME

do not exceed recommended dosage.

NIGHTTIME

- do not exceed recommended dosage
- excitability may occur, especially in children
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

DAYTIME

- nervousness, dizziness or sleeplessness occur
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

NIGHTTIME

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

cough comes back or occurs with rash or headache that lasts

these could be signs of a serious condition

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- swallow whole; do not crush, chew, or dissolve

Daytime: adults and children 12 years and over: take 2 softgels with water every 4 hours

Nighttime: adults and children 12 years and over: take 2 softgels with water every 6 hours

children under 12 years: do not use

Other information

- store between 15-30°C (59-86°F)
- avoid excessive heat

Inactive ingredients

Daytime butylated hydroxyanisole, butylated hydroxytoluene, FD&C yellow #6, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

Nighttime D&C yellow #10, FD&C blue #1, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

COMPARE TO THE ACTIVE INGREDIENTS IN VICKS® DAYQUIL® AND

NYQUIL® LIQUICAPS®*

Multi-Symptom Cold & Flu Relief

DAYTIME

Acetaminophen, Dextromethorphan HBr, Phenylephrine HCL

Relieves

- Aches, Fever, Sore Throat
- Cough
- Nasal Congestion

Alcohol-Free

Antihistamine-Free

Softgels**

(**Liquid-Filled Capsules)

When using Daytime and Nighttime products, carefully read the labeling to ensure correct dosing

NIGHTTIME

Acetaminophen, Dextromethorphan HBr, Doxylamine Succinate

Relieves

- Aches, Fever, Sore Throat
- Cough
- Runny Nose, Sneezing

Alcohol-Free

Softgels**

(**Liquid-Filled Capsules)

*This product is not manufactured or distributed by Procter & Gamble Company. Vicks® DayQuil®, NyQuil®, LiquiCaps® are registered trademarks of The Procter and Gamble Company.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

DISTRIBUTED BY: MIDWOOD BRANDS, LLC

500 VOLVO PARKWAY, CHESAPEAKE, VA 23320

Product Label

notice any signs or symptoms. medical attention is critical for adults as well as for children even if you do not recommended dose can cause liver damage, in case of overdose, get medical help or confact a Poison Confroi Center (1-600-222-1222) right away, Quick If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warming: Taking more than the These could be signs of a serious condition. congh comes back or occurs with rash or headache that lasts. Ledness or swelling is present in new symptoms occur. ■ beju or cough gets worse or lasts more than 7 days Stop use and ask a doctor if

 ве свъели мине пили в шоло, лешств от обесали в шасинету sequence sequence and tranquilizers may increase drowsiness

■ marked drowsiness may occur ■ avoid alcoholic drinks

When using this product at do not exceed recommended dosage at excitability may occur, especially in children

 taking the blood thinning drug warfarin
 taking sedatives or tranquilizers Ask a doctor or pharmacist before use if you are

■ trouble unnating due to an enlarged prostate gland sathma, chronic bronchitis, or emphysema

 a breathing problem or chronic cough that lasts such as occurs with smoking, condu just occruz with too much buiedm (mucus)

Ask a doctor before use if you have I liver disease I glaucoma

pharmacist before taking this product. If you are now taking a prescription monoamine oxidase inhibitor (MAOI)
(cartain drugs for depression, psychiathic, or emotional conditions, or
Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do
not know if your prescription drug contains an MOI), ask a doctor or
not know if your prescription drug contains an MOI), ask a doctor or
not know if your prescription drug contains an MOI), ask a doctor or
not have prepared.

sax a doctor or pharmacist.

nonprescription). If you are not sure whether a drug contains acetaminophen, with any other drug containing acetaminophen (prescription or

Drug Facts (continued)

Nighttime Cold & Flu Softgel

uogos suà sidus or symptoms. medical attention is critical for adults as well as for children even if you do not neilo of contact a Poison Control Center (1-804-222-1222) right away, clinick recommended dose can cause liver damage, in case of overdose, get medical It pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children, Overdose warming: Taking more than the

Luese conia de signs of a senous condidon. conditionmes back or occurs with rash or headache that lasts.

■ fever gets worse or lasts more than 3 days ■ Lequese or swelling is present ■ new symbioms occur ■ psin, cough, or nasal congestion gets worse or lasts more than 7 days ■ uervousness, dizziness, or sieepiessness occur Stop use and ask a doctor if

When using this product, do not exceed recommended dosage.

CITY WAITERING

Ask a doctor or pharmacist before use if you are taking the blood thinning

seu ton od

 bersistent or chronic cough such as occurs with smoking, asthma, or m trouble unnating due to enlarged prostate gland condy that occurs with too much phiegm (mucus) aunssaud poorg ufiru assesib bionyriti = Saladeid .

■ liver disease ■ heart disease Ask a doctor before use if you have

pharmacist before taking this product.

Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or If you are now baking a prescription monoamine codasse inhibitor (MADI) (certain drugs for depression, psychiatinc, or emotional conditions, or

sax a doctor or pharmacist. nonprescription). If you are not sure whether a drug contains acetaminophen, with any other drug containing acetaminophen (prescription or

accompanied or followed by fever, headache, rash, nausea, or vomiting, consult

Sore throat warning: If sore throat is severe, persists for more than 2 days, is

occur if you take: more than 4,000 mg of acetaminophen in 24 hours Liver warning: This product contains acetaminophen. Severe liver damage may

Dund Facts (continued) Daytime Cold & Flu Softgel

TROUGH BUOS III

accompanied or followed by fever, headache, rash, nausea, or vomiting, consult Sore throat warning: If sore throat is severe, persists for more than 2 days, is misy actions. Symptoms may cause severe sidn reactions. Symptoms symptoms in may bitcher at the man and severe and severe measurements and severe measurements and severe measurements. The property severe is severe measurements and severe is cause a process in the measurements and severe is severe measurements.

3 or more alcoholic drinks every day while using this product ■ with other drugs containing acetaminophen

cocur if you take: In more than 4,000 mg of acetaminophen in 24 hours Liver warming: This product contains acetaminophen. Severe liver damage may

minor achee and pains artever around mose & sneezing a sone throat cough due to minor throat and bronchial irritation nzes . Frembousuji) reijeves common cold and flu symptoms:

> Doxylamine succinate 6.25 mg. Dextromethorphan HBr 15 mg.

> > Drug Facts

Active ingredients (in each softgel)

Nighttime Cold & Flu Softgel

COMPARE TO THE ACTIVE INGREDIENTS IN VICKS" DAYQUIL" AND NYQUIL" COLD & FLU LIQUICAPS"*

cough due to minor throat and pronchial irritation in headache minor aches and pains in tever шазај соодварои nass # pemporarily relieves common cold and thu symptoms: -рт с кин эпппарилаги Cough suppressant Dextromethorphan HBr 10 mg condu anbbuesasur səsodund Active ingredients (in each softgel) sasodina Drug Facts Daytime Cold & Flu Softgel

■ with other drugs containing acetaminophen

FAMILY Ness™

Multi-Symptom Cold & Flu

Acetaminophen, Dextromethorphan HBr.

Cough

Runny Nose, Sneezing

Alcohol-Free

8 Softgels** (**Liquid-Filled Capsules)





DISTRIBUTED BY: MIDWOOD BRANDS, LLC 500 VOLVO PARKWAY, CHESAPEAKE, VA 23320

NIGHTTIME

Doxylamine Succinate

Aches, Fever, Sore Throat

Actual Size

DAYTIME

Acetaminophen, Dextromethorphan HBr,

"Liquid-Filled Capsules)

When using Daytime and Nighttime products, carefully read the labeling to ensure correct dosing.

TOTAL 24 Softgels



id unused product w for a refund (with n

r 100% SATISFIED? urn package and unus nily Dollar store for a r

DO NOT USE IF CARTON IS OPENED OR IF BLISTER BROKEN OR SHOWS ANY SIGNS OF TAMPERING. within 30 days to any receipt) or exchange.

AMPER EVIDENT: D UNIT IS TORN, BR

TAMPER

Vicks*, DayQuil*, MyQuil*, and LiquiCaps* are registered trademarks of The Procter and Gamble Company. "This product is not manufactured or distributed by The Procter & Gamble Company. Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST Call 1-877-753-3935 Monday-Friday 9AM-5PM EST Questions or comments? glycol, povidone, propylene glycol, purfied water, sorbitan, sorbitol, white ink manntol, polyethylene glycol, povidone, propylene glycol, purifled water, sorbitan, sorbitol, white ink hydroxytoluene, FD&C yellow #6, gelatin, glycerin, mannitol, polyethylene INSCINE INGREDIENTS Dutylated hydroxyanisole, butylated INACTIVE INGICEDIEN D&C yellow #10, FD&C blue #1, geletin, glycerin, ■ store between 15-30°C (59-86°F)
■ avoid excessive heat ■ store between 15-30°C (59-86°F)
■ avoid excessive heat Other information Other information ■ cyjjqueu nuqeu 15 years: do not use children under 12 years; do not use ■ swallow whole; do not crush, chew, or dissolve ■ adults and children 12 years and over: take 2 sofigels with water every 4 hours ■ adults and children 12 years and over: take 2 softgels with water every 6 hours swallow whole; do not crush, chew, or dissolve ■ do not take more than 4 doses in 24 hours ■ do not take more than 4 doses in 24 hours ■ do not take more than directed (see Overdose warning) ■ do not take more than directed (see Overdose warning) Directions Directions Drug Facts (continued) Drug Facts (continued) Daytime Cold & Flu Softgel Nighttime Cold & Flu Softgel

FAMILY WELLNESS Multi-Symptom Cold & Flu

DAYTIME NIGHTTIME COLD AND FLU RELIEF MULTI-SYMPTOM

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl kit

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:55319-850

Packagir	ıg

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	NDC:55319-850- 24	1 in 1 KIT; Type 0: Not a Combination Product	09/30/2019	09/30/2025

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	16 BLISTER PACK	16
Part 2	32 BLISTER PACK	32

Part 1 of 2

NIGHTTIME COLD FLU RELIEF

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate capsule

Product Information

Route of Administration ORAL

Active Ingredient/Active Moiety

l	Ingredient Name	Basis of Strength	Strength
l	ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII: 36209ITL9D)	ACETAMINOPHEN	325 mg

	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients			
Ingredient Name	Strength		
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)			
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)			
GELATIN (UNII: 2G86QN327L)			
GLYCERIN (UNII: PDC6A3C0OX)			
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)			
POVIDONE (UNII: FZ 989GH94E)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
WATER (UNII: 059QF0KO0R)			
SORBITAN (UNII: 6092ICV9RU)			
SORBITOL (UNII: 506T60A25R)			
MANNITOL (UNII: 3OWL53L36A)			

Product Characteristics				
Color	green	Score	no score	
Shape	CAPSULE	Size	20mm	
Flavor		Imprint Code	P30	
Contains				

П	Packaging				
-	tem Code	Package Description	Marketing Start Date	Marketing End Date	
	1	8 in 1 CARTON			
	1	1 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph final	part341	09/30/2019		

Part 2 of 2

DAYTIME COLD FLU RELIEF

acetaminophen, dextromethorphan hbr, phenylephrine hcl capsule

Product Information

Route of Administration

ORAL

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg		
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg		
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg		

Inactive Ingredients		
Ingredient Name	Strength	
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)		
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)		
MANNITOL (UNII: 30WL53L36A)		
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)		
GELATIN (UNII: 2G86QN327L)		
GLYCERIN (UNII: PDC6A3C0OX)		
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)		
POVIDONE (UNII: FZ 989GH94E)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
WATER (UNII: 059QF0KO0R)		
SORBITAN (UNII: 6092ICV9RU)		
SORBITOL (UNII: 506T60A25R)		

Product Characteristics				
Color	orange	Score	no score	
Shape	CAPSULE	Size	20mm	
Flavor		Imprint Code	P19	
Contains				

Packaging							
#	ltem Code	Package Description	Marketing Start Date	Marketing End Date			
1		16 in 1 CARTON					
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product					

Marketing Information							
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date				
OTC monograph final	part341	09/30/2019					

Marketing Information							
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date				
OTC monograph final	part341	09/30/2019	09/30/2025				

Labeler - Family Dollar (FAMILY WELLNESS) (024472631)

Revised: 1/2023 Family Dollar (FAMILY WELLNESS)