

**TRIAMTERENE AND HYDROCHLOROTHIAZIDE- triamterene and hydrochlorothiazide capsule**  
**AvKARE**

-----  
**Triamterene and Hydrochlorothiazide Capsules, USP**


**Rx Only**

**DESCRIPTION**

Each triamterene and hydrochlorothiazide capsule for oral use contains triamterene 37.5 mg and hydrochlorothiazide 25 mg. Hydrochlorothiazide is a diuretic/antihypertensive agent and triamterene is an antikaliuretic agent.

Hydrochlorothiazide is slightly soluble in water. It is soluble in dilute ammonia, dilute aqueous sodium hydroxide, and dimethylformamide. It is sparingly soluble in methanol.


Hydrochlorothiazide is 6-chloro-3,4-dihydro-2 H-1, 2, 4-benzothiadiazine-7- sulfonamide 1,1-dioxide, and its structural formula is:



Molecular Formula: C<sub>7</sub>H<sub>8</sub>ClN<sub>3</sub>O<sub>4</sub>S<sub>2</sub>      M.W. 297.74

At 50°C, triamterene is practically insoluble in water (less than 0.1%). It is soluble in formic acid, sparingly soluble in methoxyethanol, and very slightly soluble in alcohol.

Triamterene is 2, 4, 7-triamino-6-phenylpteridine and its structural formula is:



Molecular Formula: C<sub>12</sub>H<sub>11</sub>N<sub>7</sub> M.W. 253.26

Inactive ingredients consist of lactose monohydrate, pregelatinized starch, sodium starch glycolate, polysorbate 80, citric acid anhydrous, povidone, and magnesium stearate. The capsule shell consists of titanium dioxide and gelatin. The capsule imprinting ink consists of shellac glaze in ethanol, iron oxide black, n-butyl alcohol, propylene glycol, ethanol, methanol, FD&C Blue # 2 Aluminum Lake, FD&C Red # 40 Aluminum Lake, FD&C Blue # 1 Aluminum Lake, and D&C Yellow # 10 Aluminum Lake.

Triamterene and hydrochlorothiazide capsules, USP meet USP Dissolution Test 3 as published in the current USP monograph for triamterene and hydrochlorothiazide capsules.

## **CLINICAL PHARMACOLOGY**

The triamterene and hydrochlorothiazide capsule is a diuretic/antihypertensive drug product that combines natriuretic and antikaliuretic effects. Each component complements the action of the other. The hydrochlorothiazide component blocks the reabsorption of sodium and chloride ions, and thereby increases the quantity of sodium traversing the distal tubule and the volume of water excreted. A portion of the additional sodium presented to the distal tubule is exchanged there for potassium and hydrogen ions. With continued use of hydrochlorothiazide and depletion of sodium, compensatory mechanisms tend to increase this exchange and may produce excessive loss of potassium, hydrogen, and chloride ions. Hydrochlorothiazide also decreases the excretion of calcium and uric acid, may increase the excretion of iodide, and may reduce glomerular filtration rate. The exact mechanism of the antihypertensive effect of hydrochlorothiazide is not known.

The triamterene component of triamterene and hydrochlorothiazide capsules, USP exerts its diuretic effect on the distal renal tubule to inhibit the reabsorption of sodium in exchange for potassium and hydrogen ions. Its natriuretic activity is limited by the amount of sodium reaching its site of action. Although it blocks the increase in this exchange that is stimulated by mineralocorticoids (chiefly aldosterone), it is not a competitive antagonist of aldosterone and its activity can be demonstrated in adrenalectomized rats and patients with Addison's disease. As a result, the dose of triamterene required is not proportionally related to the level of mineralocorticoid activity, but is dictated by the response of the individual patients, and the kaliuretic effect of concomitantly administered drugs. By inhibiting the distal tubular exchange mechanism, triamterene maintains or increases the sodium excretion and reduces the excess loss of potassium, hydrogen and chloride ions induced by hydrochlorothiazide. As with hydrochlorothiazide, triamterene may reduce glomerular filtration and renal plasma flow. Via this mechanism it may reduce uric acid excretion although it has no tubular effect on uric acid reabsorption or secretion. Triamterene does not affect calcium excretion. No predictable antihypertensive effect has been demonstrated for triamterene.

Duration of diuretic activity and effective dosage range of the hydrochlorothiazide and triamterene components of triamterene and hydrochlorothiazide capsules, USP are similar. Onset of diuresis with triamterene and hydrochlorothiazide capsules, USP takes place within 1 hour, peaks at 2 to 3 hours and tapers off during the subsequent 7 to 9 hours.

Triamterene and hydrochlorothiazide capsules, USP are well absorbed.

Upon administration of a single oral dose to fasted normal male volunteers, the following mean pharmacokinetic parameters were determined:

	<b>AUC (0-48) ng*hrs/mL (± SD)</b>	<b>C<sub>max</sub> ng/mL (± SD)</b>	<b>Median T<sub>max</sub> Hrs</b>	<b>Ae Mg (± SD )</b>
triamterene	148.7 (87.9)	46.4 (29.4)	1.1	2.7 (1.4)
hydroxytriamterene sulfate	1,865 (471)	720 (364)	1.3	19.7 (6.1)
hydrochlorothiazide	834 (177)	135.1 (35.7)	2.0	14.3 (3.8)

where AUC (0-48), C<sub>max</sub>, T<sub>max</sub> and Ae represent area under the plasma concentration versus time plot, maximum plasma concentration, time to reach C<sub>max</sub>, and amount excreted in urine over 48 hours.

One triamterene and hydrochlorothiazide capsule is bioequivalent to a single-entity 25 mg hydrochlorothiazide tablet and 37.5 mg triamterene capsule used in the double-blind clinical trial below (see Clinical Trials).

In a limited study involving 12 subjects, coadministration of triamterene and hydrochlorothiazide capsules, USP with a high-fat meal resulted in: (1) an increase in the mean bioavailability of triamterene by about 67% (90% confidence interval = 0.99, 1.90), p-hydroxytriamterene sulfate by about 50% (90% confidence interval = 1.06, 1.77), hydrochlorothiazide by about 17% (90% confidence interval = 0.90, 1.34); (2) increases in the peak concentrations of triamterene and p-hydroxytriamterene; and (3) a delay of up to 2 hours in the absorption of the active constituents.

## Clinical Trials

A placebo-controlled, double-blind trial was conducted to evaluate the efficacy of triamterene and hydrochlorothiazide capsules, USP. This trial demonstrated that triamterene and hydrochlorothiazide capsules, USP (37.5 mg triamterene/25 mg hydrochlorothiazide) were effective in controlling blood pressure while reducing the incidence of hydrochlorothiazide-induced hypokalemia. This trial involved 636 patients with mild to moderate hypertension controlled by hydrochlorothiazide 25 mg daily and who had hypokalemia (serum potassium <3.5 mEq/L) secondary to the hydrochlorothiazide. Patients were randomly assigned to 4 weeks' treatment with once-daily regimens of 25 mg hydrochlorothiazide plus placebo, or 25 mg hydrochlorothiazide combined with one of the following doses of triamterene: 25 mg, 37.5 mg, 50 mg, or 75 mg.

Blood pressure and serum potassium were monitored at baseline and throughout the trial. All five treatment groups had similar mean blood pressure and serum potassium concentrations at baseline (mean systolic blood pressure range: 137±14 mmHg to 140±16 mmHg; mean diastolic blood pressure range: 86±9 mmHg to 88±8 mmHg; mean serum potassium range: 2.3 to 3.4 mEq/L with the majority of patients having

values between 3.1 and 3.4 mEq/L).

While all triamterene regimens reversed hypokalemia, at week 4 the 37.5 mg regimen proved optimal compared with the other tested regimens. On this regimen, 81% of the patients had a significant ( $p < 0.05$ ) reversal of hypokalemia vs. 59% of patients on the placebo/hydrochlorothiazide regimen. The mean serum potassium concentration on 37.5 mg triamterene went from  $3.2 \pm 0.2$  mEq/L at baseline to  $3.7 \pm 0.3$  mEq/L at week 4, a significantly greater ( $p < 0.05$ ) improvement than that achieved with placebo/hydrochlorothiazide (i.e.,  $3.2 \pm 0.2$  mEq/L at baseline and  $3.5 \pm 0.4$  mEq/L at week 4). Also, 51% of patients in the 37.5 mg triamterene group had an increase in serum potassium of  $\geq 0.5$  mEq/L at week 4 vs. 33% in the placebo group. The 37.5 mg triamterene/25 mg hydrochlorothiazide regimen also maintained control of blood pressure; mean supine systolic blood pressure at week 4 was  $138 \pm 21$  mmHg while mean supine diastolic blood pressure was  $87 \pm 13$  mmHg.

## INDICATIONS AND USAGE

**This fixed combination drug is not indicated for the initial therapy of edema or hypertension except in individuals in whom the development of hypokalemia cannot be risked.**

Triamterene and hydrochlorothiazide capsules, USP are indicated for the treatment of hypertension or edema in patients who develop hypokalemia on hydrochlorothiazide alone.

Triamterene and hydrochlorothiazide capsules, USP are also indicated for those patients who require a thiazide diuretic and in whom the development of hypokalemia cannot be risked.

Triamterene and hydrochlorothiazide capsules, USP may be used alone or as an adjunct to other antihypertensive drugs, such as beta-blockers. Since triamterene and hydrochlorothiazide capsules, USP may enhance the action of these agents, dosage adjustments may be necessary.

**Usage in Pregnancy:** The routine use of diuretics in an otherwise healthy woman is inappropriate and exposes mother and fetus to unnecessary hazard. Diuretics do not prevent development of toxemia of pregnancy, and there is no satisfactory evidence that they are useful in the treatment of developed toxemia.

Edema during pregnancy may arise from pathological causes or from the physiologic and mechanical consequences of pregnancy. Diuretics are indicated in pregnancy when edema is due to pathologic causes, just as they are in the absence of pregnancy. Dependent edema in pregnancy resulting from restriction of venous return by the expanded uterus is properly treated through elevation of the lower extremities and use of support hose; use of diuretics to lower intravascular volume in this case is illogical and unnecessary. There is hypervolemia during normal pregnancy which is harmful to neither the fetus nor the mother (in the absence of cardiovascular disease), but which is associated with edema, including generalized edema in the majority of pregnant women. If this edema produces discomfort, increased recumbency will often provide relief. In rare instances this edema may cause extreme discomfort which is not relieved by rest. In these cases a short course of diuretics may provide relief and may be appropriate.

## CONTRAINDICATIONS

**Antikaliuretic Therapy and Potassium Supplementation:** Triamterene and hydrochlorothiazide capsules, USP should not be given to patients receiving other potassium-sparing agents such as spironolactone, amiloride, or other formulations containing triamterene. Concomitant potassium-containing salt substitutes should also not be used.

Potassium supplementation should not be used with triamterene and hydrochlorothiazide capsules, USP except in severe cases of hypokalemia. Such concomitant therapy can be associated with rapid increases in serum potassium levels. If potassium supplementation is used, careful monitoring of the serum potassium level is necessary.

**Impaired Renal Function:** Triamterene and hydrochlorothiazide capsules, USP are contraindicated in patients with anuria, acute and chronic renal insufficiency or significant renal impairment.

**Hypersensitivity:** Hypersensitivity to either drug in the preparation or to other sulfonamide-derived drugs is a contraindication.

**Hyperkalemia:** Triamterene and hydrochlorothiazide capsules, USP should not be used in patients with preexisting elevated serum potassium.

## WARNINGS

### Hyperkalemia

Abnormal elevation of serum potassium levels (greater than or equal to 5.5 mEq/liter) can occur with all potassium-sparing diuretic combinations, including triamterene and hydrochlorothiazide capsules, USP. Hyperkalemia is more likely to occur in patients with renal impairment and diabetes (even without evidence of renal impairment), and in the elderly or severely ill. Since uncorrected hyperkalemia may be fatal, serum potassium levels must be monitored at frequent intervals especially in patients first receiving triamterene and hydrochlorothiazide capsules, USP, when dosages are changed or with any illness that may influence renal function.

If hyperkalemia is suspected (warning signs include paresthesias, muscular weakness, fatigue, flaccid paralysis of the extremities, bradycardia, and shock), an electrocardiogram (ECG) should be obtained. However, it is important to monitor serum potassium levels because hyperkalemia may not be associated with ECG changes.

If hyperkalemia is present, triamterene and hydrochlorothiazide capsules, USP should be discontinued immediately and a thiazide alone should be substituted. If the serum potassium exceeds 6.5 mEq/liter more vigorous therapy is required. The clinical situation dictates the procedures to be employed. These include the intravenous administration of calcium chloride solution, sodium bicarbonate solution, and/or the oral or parenteral administration of glucose with a rapid-acting insulin preparation. Cationic exchange resins such as sodium polystyrene sulfonate may be orally or rectally administered. Persistent hyperkalemia may require dialysis.

The development of hyperkalemia associated with potassium-sparing diuretics is accentuated in the presence of renal impairment (see CONTRAINDICATIONS section). Patients with mild renal functional impairment should not receive this drug without frequent and continuing monitoring of serum electrolytes. Cumulative drug effects may be observed in patients with impaired renal function. The renal clearances of hydrochlorothiazide and the pharmacologically active metabolite of triamterene, the sulfate ester of hydroxytriamterene, have been shown to be reduced and the plasma levels increased following administration of triamterene and hydrochlorothiazide capsules, USP to elderly patients and patients with impaired renal function.

Hyperkalemia has been reported in diabetic patients with the use of potassium-sparing agents even in the absence of apparent renal impairment. Accordingly, serum electrolytes must be frequently monitored if triamterene and hydrochlorothiazide capsules, USP are used in diabetic patients.

**Metabolic or Respiratory Acidosis:** Potassium-sparing therapy should also be avoided in severely ill patients in whom respiratory or metabolic acidosis may occur. Acidosis may be associated with rapid elevations in serum potassium levels. If triamterene and hydrochlorothiazide capsules, USP are employed, frequent evaluations of acid/base balance and serum electrolytes are necessary.

**Acute Myopia and Secondary Angle-Closure Glaucoma:** Hydrochlorothiazide, a sulfonamide, can cause an idiosyncratic reaction, resulting in acute transient myopia and acute angle-closure glaucoma. Symptoms include acute onset of decreased visual acuity or ocular pain and typically occur within hours to weeks of drug initiation. Untreated acute angle-closure glaucoma can lead to permanent vision loss. The primary treatment is to discontinue hydrochlorothiazide as rapidly as possible. Prompt medical or surgical treatments may need to be considered if the intraocular pressure remains uncontrolled. Risk factors for developing acute angle-closure glaucoma may include a history of sulfonamide or penicillin allergy.

## PRECAUTIONS

### Diabetes

Caution should be exercised when administering triamterene and hydrochlorothiazide capsules, USP to patients with diabetes, since thiazides may cause hyperglycemia, glycosuria, and alter insulin requirements in diabetes. Also, diabetes mellitus may become manifest during thiazide administration.

### Impaired Hepatic Function

Thiazides should be used with caution in patients with impaired hepatic function. They can precipitate hepatic coma in patients with severe liver disease. Potassium depletion induced by the thiazide may be important in this connection. Administer triamterene and hydrochlorothiazide capsules, USP cautiously and be alert for such early signs of impending coma as confusion, drowsiness, and tremor; if mental confusion increases discontinue triamterene and hydrochlorothiazide capsules, USP for a few days. Attention must be given to other factors that may precipitate hepatic coma, such as blood in the gastrointestinal tract or preexisting potassium depletion.

## Hypokalemia

Hypokalemia is uncommon with triamterene and hydrochlorothiazide capsules, USP; but, should it develop, corrective measures should be taken such as potassium supplementation or increased intake of potassium-rich foods. Institute such measures cautiously with frequent determinations of serum potassium levels, especially in patients receiving digitalis or with a history of cardiac arrhythmias. If serious hypokalemia (serum potassium less than 3.0 mEq/L) is demonstrated by repeat serum potassium determinations, triamterene and hydrochlorothiazide capsules, USP should be discontinued and potassium chloride supplementation initiated. Less serious hypokalemia should be evaluated with regard to other coexisting conditions and treated accordingly.

## Electrolyte Imbalance

Electrolyte imbalance, often encountered in such conditions as heart failure, renal disease or cirrhosis of the liver, may also be aggravated by diuretics and should be considered during therapy with triamterene and hydrochlorothiazide capsules, USP when using high doses for prolonged periods or in patients on a salt-restricted diet.

Serum determinations of electrolytes should be performed, and are particularly important if the patient is vomiting excessively or receiving fluids parenterally. Possible fluid and electrolyte imbalance may be indicated by such warning signs as: dry mouth, thirst, weakness, lethargy, drowsiness, restlessness, muscle pain or cramps, muscular fatigue, hypotension, oliguria, tachycardia, and gastrointestinal symptoms.

## Hypochloremia

Although any chloride deficit is generally mild and usually does not require specific treatment except under extraordinary circumstances (as in liver disease or renal disease), chloride replacement may be required in the treatment of metabolic alkalosis. Dilutional hyponatremia may occur in edematous patients in hot weather; appropriate therapy is water restriction, rather than administration of salt, except in rare instances when the hyponatremia is life threatening. In actual salt depletion, appropriate replacement is the therapy of choice.

## Renal Stones

Triamterene has been found in renal stones in association with the other usual calculus components. Triamterene and hydrochlorothiazide capsules, USP should be used with caution in patients with a history of renal stones.

## Laboratory Tests

**Serum Potassium:** The normal adult range of serum potassium is 3.5 to 5.0 mEq per liter with 4.5 mEq often being used for a reference point. If hypokalemia should develop, corrective measures should be taken such as potassium supplementation or increased dietary intake of potassium-rich foods.

Institute such measures cautiously with frequent determinations of serum potassium levels. Potassium levels persistently above 6 mEq per liter require careful observation and treatment.

Serum potassium levels do not necessarily indicate true body potassium concentration.

A rise in plasma pH may cause a decrease in plasma potassium concentration and an increase in the intracellular potassium concentration. Discontinue corrective measures for hypokalemia immediately if laboratory determinations reveal an abnormal elevation of serum potassium. Discontinue triamterene and hydrochlorothiazide capsules, USP and substitute a thiazide diuretic alone until potassium levels return to normal.

**Serum Creatinine and BUN:** Triamterene and hydrochlorothiazide capsules, USP may produce an elevated blood urea nitrogen level, creatinine level or both. This apparently is secondary to a reversible reduction of glomerular filtration rate or a depletion of intravascular fluid volume (prerenal azotemia) rather than renal toxicity; levels usually return to normal when triamterene and hydrochlorothiazide capsules, USP are discontinued. If azotemia increases, discontinue triamterene and hydrochlorothiazide capsules, USP. Periodic BUN or serum creatinine determinations should be made, especially in elderly patients and in patients with suspected or confirmed renal insufficiency.

**Serum PBI:** Thiazide may decrease serum PBI levels without sign of thyroid disturbance.

**Parathyroid Function:** Thiazides should be discontinued before carrying out tests for parathyroid function. Calcium excretion is decreased by thiazides. Pathologic changes in the parathyroid glands with hypercalcemia and hypophosphatemia have been observed in a few patients on prolonged thiazide therapy. The common complications of hyperparathyroidism such as bone resorption and peptic ulceration have not been seen.

## **Drug Interactions**

**Angiotensin-converting Enzyme Inhibitors:** Potassium-sparing agents should be used with caution in conjunction with angiotensin-converting enzyme (ACE) inhibitors due to an increased risk of hyperkalemia.

**Oral Hypoglycemic Drugs:** Concurrent use with chlorpropamide may increase the risk of severe hyponatremia.

**Nonsteroidal Anti-inflammatory Drugs:** A possible interaction resulting in acute renal failure has been reported in a few patients on triamterene and hydrochlorothiazide capsules, USP when treated with indomethacin, a nonsteroidal anti-inflammatory agent. Caution is advised in administering nonsteroidal anti-inflammatory agents with triamterene and hydrochlorothiazide capsules, USP.

**Lithium:** Lithium generally should not be given with diuretics because they reduce its renal clearance and increase the risk of lithium toxicity. Read circulars for lithium preparations before use of such concomitant therapy with triamterene and hydrochlorothiazide capsules, USP.

**Surgical Considerations:** Thiazides have been shown to decrease arterial responsiveness to norepinephrine (an effect attributed to loss of sodium). This diminution is not sufficient to preclude effectiveness of the pressor agent for therapeutic use. Thiazides have also been shown to increase the paralyzing effect of nondepolarizing muscle relaxants such as tubocurarine (an effect attributed to potassium loss); consequently caution should be observed in patients undergoing surgery.

**Other Considerations:** Concurrent use of hydrochlorothiazide with amphotericin B or

corticosteroids or corticotropin (ACTH) may intensify electrolyte imbalance, particularly hypokalemia, although the presence of triamterene minimizes the hypokalemic effect.

Thiazides may add to or potentiate the action of other antihypertensive drugs. See INDICATIONS AND USAGE for concomitant use with other antihypertensive drugs.

The effect of oral anticoagulants may be decreased when used concurrently with hydrochlorothiazide; dosage adjustments may be necessary.

Triamterene and hydrochlorothiazide capsules, USP may raise the level of blood uric acid; dosage adjustments of antigout medication may be necessary to control hyperuricemia and gout.

The following agents given together with triamterene may promote serum potassium accumulation and possibly result in hyperkalemia because of the potassium-sparing nature of triamterene, especially in patients with renal insufficiency: blood from blood bank (may contain up to 30 mEq of potassium per liter of plasma or up to 65 mEq per liter of whole blood when stored for more than 10 days); low-salt milk (may contain up to 60 mEq of potassium per liter); potassium-containing medications (such as parenteral penicillin G potassium); salt substitutes (most contain substantial amounts of potassium).

Exchange resins, such as sodium polystyrene sulfonate, whether administered orally or rectally, reduce serum potassium levels by sodium replacement of the potassium; fluid retention may occur in some patients because of the increased sodium intake.

Chronic or overuse of laxatives may reduce serum potassium levels by promoting excessive potassium loss from the intestinal tract; laxatives may interfere with the potassium-retaining effects of triamterene.

The effectiveness of methenamine may be decreased when used concurrently with hydrochlorothiazide because of alkalinization of the urine.

### **Drug/Laboratory Test Interactions**

Triamterene and quinidine have similar fluorescence spectra; thus, triamterene and hydrochlorothiazide capsules, USP will interfere with the fluorescent measurement of quinidine.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

**Carcinogenesis:** Long-term studies have not been conducted with triamterene and hydrochlorothiazide capsules, USP (the triamterene/hydrochlorothiazide combination), or with triamterene alone.

**Hydrochlorothiazide:** Two-year feeding studies in mice and rats, conducted under the auspices of the National Toxicology Program (NTP), treated mice and rats with doses of hydrochlorothiazide up to 600 and 100 mg/kg/day, respectively. On a body-weight basis, these doses are 600 times (in mice) and 100 times (in rats) the Maximum Recommended Human Dose (MRHD) for the hydrochlorothiazide component of triamterene and hydrochlorothiazide capsules, USP at 50 mg/day (or 1.0 mg/kg/day based on 50 kg individuals). On the basis of body-surface area, these doses are 56 times (in mice) and 21 times (in rats) the MRHD. These studies uncovered no evidence of carcinogenic potential of hydrochlorothiazide in rats or female mice, but there was equivocal evidence of hepatocarcinogenicity in male mice.

**Mutagenesis:** Studies of the mutagenic potential of triamterene and hydrochlorothiazide capsules, USP (the triamterene/hydrochlorothiazide combination), or of triamterene alone have not been performed.

**Hydrochlorothiazide:** Hydrochlorothiazide was not genotoxic in in vitro assays using strains TA 98, TA 100, TA 1535, TA 1537 and TA 1538 of *Salmonella typhimurium* (the Ames test); in the Chinese Hamster Ovary (CHO) test for chromosomal aberrations; or in in vivo assays using mouse germinal cell chromosomes, Chinese hamster bone marrow chromosomes, and the *Drosophila* sex-linked recessive lethal trait gene. Positive test results were obtained in the in vitro CHO Sister Chromatid Exchange (clastogenicity) test, and in the mouse Lymphoma Cell (mutagenicity) assays, using concentrations of hydrochlorothiazide of 43 to 1300 mcg/mL.

Positive test results were also obtained in the *Aspergillus nidulans* nondisjunction assay, using an unspecified concentration of hydrochlorothiazide.

**Impairment of Fertility:** Studies of the effects of triamterene and hydrochlorothiazide capsules, USP (the triamterene/hydrochlorothiazide combination), or of triamterene alone on animal reproductive function have not been conducted.

**Hydrochlorothiazide:** Hydrochlorothiazide had no adverse effects on the fertility of mice and rats of either sex in studies wherein these species were exposed, via their diet, to doses of up to 100 and 4 mg/kg/day, respectively, prior to mating and throughout gestation. Corresponding multiples of the MRHD are 100 (mice) and 4 (rats) on the basis of body-weight and 9.4 (mice) and 0.8 (rats) on the basis of body-surface area.

## Pregnancy

Teratogenic Effects: Category C

**Triamterene and hydrochlorothiazide capsules, USP:** Animal reproduction studies to determine the potential for fetal harm by triamterene and hydrochlorothiazide capsules, USP have not been conducted. However, a One Generation Study in the rat approximated composition of triamterene and hydrochlorothiazide capsules, USP by using a 1:1 ratio of triamterene to hydrochlorothiazide (30:30 mg/kg/day); there was no evidence of teratogenicity at those doses which were, on a body-weight basis, 15 and 30 times, respectively, the MRHD, and on the basis of body-surface area, 3.1 and 6.2 times, respectively, the MRHD.

The safe use of triamterene and hydrochlorothiazide capsules, USP in pregnancy has not been established since there are no adequate and well-controlled studies with triamterene and hydrochlorothiazide capsules, USP in pregnant women. Triamterene and hydrochlorothiazide capsules, USP should be used during pregnancy only if the potential benefit justifies the risk to the fetus.

**Triamterene:** Reproduction studies have been performed in rats at doses as high as 20 times the MRHD on the basis of body-weight, and 6 times the human dose on the basis of body-surface area without evidence of harm to the fetus due to triamterene.

Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

**Hydrochlorothiazide:** Hydrochlorothiazide was orally administered to pregnant mice and rats during respective periods of major organogenesis at doses up to 3,000 and

1,000 mg/kg/day, respectively. At these doses, which are multiples of the MRHD equal to 3,000 for mice and 1,000 for rats, based on body-weight, and equal to 282 for mice and 206 for rats, based on body-surface area, there was no evidence of harm to the fetus.

There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

### Nonteratogenic Effects

Thiazides and triamterene have been shown to cross the placental barrier and appear in cord blood. The use of thiazides and triamterene in pregnant women requires that the anticipated benefit be weighed against possible hazards to the fetus. These hazards include fetal or neonatal jaundice, pancreatitis, thrombocytopenia, and possible other adverse reactions which have occurred in the adult.

### Nursing Mothers

Thiazides and triamterene in combination have not been studied in nursing mothers. Triamterene appears in animal milk; this may occur in humans. Thiazides are excreted in human breast milk. If use of the combination drug product is deemed essential, the patient should stop nursing.

### Pediatric Use

Safety and effectiveness in pediatric patients have not been established.

## ADVERSE REACTIONS

Adverse effects are listed in decreasing order of severity.

**Hypersensitivity:** Anaphylaxis, rash, urticaria, subacute cutaneous lupus erythematosus-like reactions, photosensitivity.

**Cardiovascular:** Arrhythmia, postural hypotension.

**Metabolic:** Diabetes mellitus, hyperkalemia, hypokalemia, hyponatremia, acidosis, hypercalcemia, hyperglycemia, glycosuria, hyperuricemia, hypochloremia.

**Gastrointestinal:** Jaundice and/or liver enzyme abnormalities, pancreatitis, nausea and vomiting, diarrhea, constipation, abdominal pain.

**Renal:** Acute renal failure (one case of irreversible renal failure has been reported), interstitial nephritis, renal stones composed primarily of triamterene, elevated BUN, and serum creatinine, abnormal urinary sediment.

**Hematologic:** Leukopenia, thrombocytopenia and purpura, megaloblastic anemia.

**Musculoskeletal:** Muscle cramps.

**Central Nervous System:** Weakness, fatigue, dizziness, headache, dry mouth.

**Miscellaneous:** Impotence, sialadenitis.

Thiazides alone have been shown to cause the following additional adverse reactions:

**Central Nervous System:** Paresthesias, vertigo.

**Ophthalmic:** Xanthopsia, transient blurred vision.

**Respiratory:** Allergic pneumonitis, pulmonary edema, respiratory distress.

**Other:** Necrotizing vasculitis, exacerbation of lupus.

**Hematologic:** Aplastic anemia, agranulocytosis, hemolytic anemia.

**Neonate and infancy:** Thrombocytopenia and pancreatitis—rarely, in newborns whose mothers have received thiazides during pregnancy.

**Skin:** Erythema multiforme including Stevens-Johnson syndrome, exfoliative dermatitis including toxic epidermal necrolysis.

## **OVERDOSAGE**

Electrolyte imbalance is the major concern (see WARNINGS section). Symptoms reported include: polyuria, nausea, vomiting, weakness, lassitude, fever, flushed face, and hyperactive deep tendon reflexes. If hypotension occurs, it may be treated with pressor agents such as levarterenol to maintain blood pressure. Carefully evaluate the electrolyte pattern and fluid balance. Induce immediate evacuation of the stomach through emesis or gastric lavage. There is no specific antidote.

Reversible acute renal failure following ingestion of 50 tablets of a product containing a combination of 50 mg triamterene and 25 mg hydrochlorothiazide has been reported.

Although triamterene is largely protein-bound (approximately 67%), there may be some benefit to dialysis in cases of overdose.

## **DOSAGE AND ADMINISTRATION**

**The usual dose of triamterene and hydrochlorothiazide capsules, USP is one or two capsules given once daily, with appropriate monitoring of serum potassium and of the clinical effect (see WARNINGS, Hyperkalemia).**

## **HOW SUPPLIED**

Triamterene and Hydrochlorothiazide Capsules, USP contain 37.5 mg triamterene and 25 mg hydrochlorothiazide, and are available in size # 4 white opaque/white opaque capsules imprinted with Logo “LANNETT” on the cap and “1632” on the body. They are supplied as follows:

Bottles of 1000, NDC 42291-841-10

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Protect from light. Dispense in a tight, light-resistant container as defined in the USP, with a child-resistant closure.

Manufactured for:  
AvKARE, Inc.  
Pulaski, TN 38478

Mfg. Rev. 04/11

AV 06/12

**PRINCIPAL DISPLAY PANEL**

**AvKARE**

NDC 42291-841-10

**TRIAMTERENE AND HYDROCHLOROTHIAZIDE CAPSULES, USP**

**37.5 mg / 25 mg**

1000 Capsules      Rx Only

Each capsule contains:

Triamterene, USP.....37.5 mg

Hydrochlorothiazide, USP.....25 mg

Usual Dosage:

See package insert for full prescribing information.

Triamterene and hydrochlorothiazide capsules USP, 37.5 mg/25 mg meet USP Dissolution Test 3 as published in the current USP monograph for triamterene and hydrochlorothiazide capsules.

Inactive ingredients:

Lactose monohydrate, pregelatinized starch, sodium starch glycolate, polysorbate 80, citric acid anhydrous, povidone, magnesium stearate, titanium dioxide, gelatin, shellac glaze in ethanol, iron oxide black, n-butyl alcohol, propylene glycol, ethanol, methanol, FD&C Blue # 2 Aluminum Lake, FD&C Red # 40 Aluminum Lake, FD&C Blue # 1 Aluminum Lake, and D&C Yellow # 10 Aluminum Lake.

Protect from light. Dispense in a tight, light-resistant container as defined in the USP with a child-resistant closure.

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Manufactured for:

AvKARE, Inc.

Pulaski, TN 38478

Mfg. Rev. 08/10

AV 07/12

**N3 42291 84110 9**



NDC 42291-841-10

# TRIAMTERENE AND HYDROCHLOROTHIAZIDE CAPSULES, USP

37.5 mg/25 mg

1000 Capsules Rx Only

Each capsule contains:  
Triamterene, USP ..... 37.5 mg  
Hydrochlorothiazide, USP ..... 25 mg

Usual Dosage:  
See package insert for full prescribing information.  
Triamterene and hydrochlorothiazide capsules USP, 37.5 mg/25 mg meet USP Dissolution Test 3 as published in the current USP monograph for triamterene and hydrochlorothiazide capsules.

Inactive Ingredients:  
Lactose monohydrate, pregelatinized starch, sodium starch glycolate, poly sorbate 80, citric acid anhydrous, povidone, magnesium stearate, titanium dioxide, gelatin, shellac glaze in ethanol, iron oxide black, n-butyl alcohol, propylene glycol, Red # 40 Aluminum Lake, FD&C Blue # 1 Aluminum Lake, and D&C Yellow # 10 Aluminum Lake.

Protect from light. Dispense in a tight, light-resistant container as defined in the USP with a child-resistant closure. Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Keep this and all drugs out of the reach of children.

Manufactured for:  
AVKARE, Inc.  
Pulaski, TN 38478

Mfg. Rev. 08/10 AV 07/12



N 42291 84110 9

## TRIAMTERENE AND HYDROCHLOROTHIAZIDE

triamterene and hydrochlorothiazide capsule

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:42291-841(NDC:0527-1632)
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
TRIAMTERENE (UNII: WS821Z52LQ) (TRIAMTERENE - UNII:WS821Z52LQ)	TRIAMTERENE	37.5 mg
HYDROCHLOROTHIAZIDE (UNII: 0J48LPH2TH) (HYDROCHLOROTHIAZIDE - UNII:0J48LPH2TH)	HYDROCHLOROTHIAZIDE	25 mg

### Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
STARCH, CORN (UNII: O8232NY35J)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
GELATIN (UNII: 2G86QN327L)	
SHELLAC (UNII: 46N107B71O)	
ALCOHOL (UNII: 3K9958V90M)	
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	
BUTYL ALCOHOL (UNII: 8PJ61P6TS3)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
METHYL ALCOHOL (UNII: Y4S76JWM15)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	

<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	14mm
<b>Flavor</b>		<b>Imprint Code</b>	LANNETT;1632
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42291-841-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	07/03/2012	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA201407	07/03/2012	

**Labeler** - AvKARE (796560394)

Revised: 1/2022

AvKARE