

**MENSTRUAL COMPLETE- acetaminophen, caffeine, pyrilamine maleate tablet**  
**Strive Pharmaceuticals Inc.**

-----  
**RIGHT REMEDIES Menstrual Complete**

**Drug Facts**

**Active ingredients (in each gelcap)**

Acetaminophen 500 mg  
Caffeine 60 mg  
Pyrilamine maleate 15 mg

**Purposes**

Pain reliever  
Diuretic  
Antihistamine

**Uses**

for the temporary relief of these symptoms associated with menstrual periods:  
• cramps • bloating • water-weight gain • headache • backache • muscle aches •  
fatigue

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 gelcaps in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include: • skin reddening • blisters • rash • hives • facial swelling • asthma (wheezing) • shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

- liver disease
- glaucoma
- difficulty in urination due to enlargement of the prostate gland

- a breathing problem such as emphysema or chronic bronchitis

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

**When using this product**

- you may get drowsy
- avoid alcoholic drinks
- excitability may occur, especially in children
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- limit the use of caffeine-containing medications, foods, or beverages because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat. The recommended dose of this product contains about as much caffeine as a cup of coffee.

**Stop use and ask a doctor if**

- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

**If pregnant or breast-feeding,** ask a health professional before use

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

***Directions***

- do not take more than the recommended dose
- adults and children 12 years and older: take 2 gelcaps with water. Repeat every 6 hours as needed. Do not exceed 6 gelcaps per day
- children under 12 years: consult a doctor

***Other information***

- each gelcap contains **sodium 3 mg** • store at room temperature.

***Inactive ingredients***

croscarmellose sodium, D&C red#33 aluminium lake, edible black printing ink, ethyl cellulose, FD&C blue #1 aluminum lake, gelatin, glycerin, hydroxypropyl methylcellulose, Isopropyl alcohol, magnesium stearate, microcrystalline cellulose, pregelatinized potato starch, purified water, PVPK-30, sorbitol sorbitan solution, talc, titanium dioxide, triacetin

***Questions or comments?***

1-888-577-8033 Monday - Friday 8am - 4pm EST

## RIGHT REMEDIES

Compare to the active ingredients in **Midol® Complete\***

### READ AND KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGN OF TAMPERING. DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BRPKEN OR MISSING**

\*This product is not manufactured or distributed by Bayer Healthcare LLC, owner of the registered trademark Midol®.

Distributed by: Strive Pharmaceuticals Inc., East Brunswick, NJ 08816

Product of United Arab Emirates

## Packaging



### Drug Facts (continued)

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

#### Ask a doctor before use if you have

- liver disease
- glaucoma
- difficulty in urination due to enlargement of the prostate gland
- a breathing problem such as emphysema or chronic bronchitis

#### Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

Active ingredients  
72 Complete\*  
XC 70692-131-46

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**Drug Facts**

Active ingredients (in each gelcap)	Purpose
Acetaminophen 500 mg	Pain reliever
Caffeine 60 mg	Diuretic
Pyrilamine maleate 15 mg	Antihistamine

**Uses**  
for the temporary relief of these symptoms associated with menstrual periods: ■ cramps ■ bloating ■ water-weight gain ■ headache ■ backache ■ muscle aches ■ fatigue

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take  
■ more than 6 gelcaps in 24 hours, which is the maximum daily amount for this product  
■ with other drugs containing acetaminophen  
■ 3 or more alcoholic drinks every day while using this product  
**Allergy Alert:** acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:  
■ skin reddening ■ blisters ■ rash ■ hives  
■ facial swelling ■ asthma (wheezing) ■ shock  
If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Drug Facts (continued under label)**




3 70692 00017 7

PEEL HERE

LOT:  
EXP:

NO VARNISH

**When using this product**

- you may get drowsy ■ avoid alcoholic drinks
- excitability may occur, especially in children
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- limit the use of caffeine-containing medications, foods, or beverages because too much caffeine may cause nervousness, irritability, sleeplessness, and occasionally rapid heartbeat. The recommended dose of this product contains about as much caffeine as a cup of coffee.

**Stop use and ask a doctor if**

- new symptoms occur ■ redness or swelling is present
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

**If pregnant or breast-feeding,** ask a health professional before use.  
**Keep out of reach of children.**

**Overdose warning:** in case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than the recommended dose
- adults and children 12 years and older: take 2 gelcaps with water. Repeat every 6 hours, as needed. Do not exceed 6 gelcaps per day.
- children under 12 years: consult a doctor

**Other information**

- each gelcap contains: sodium 3 mg ■ store at room temperature.

**Inactive ingredients**  
croscarmellose sodium, D&C red #33 aluminum lake, edible black printing ink, ethyl cellulose, FD&C blue #1 aluminum lake, gelatin, glycerin, hydroxypropyl methylcellulose, isopropyl alcohol, magnesium stearate, microcrystalline cellulose, pregelatinized potato starch, purified water, PVP K-30, sorbitol sorbitan solution, talc, titanium dioxide, triacetin

**Questions or comments?**  
1-888-577-8033 Monday-Friday 8am-4pm EST

Distributed by: Strive Pharmaceuticals, Inc. East Brunswick, NJ 08816  
Product of United Arab Emirates  
\*This product is not manufactured or distributed by Bayer Healthcare LLC, owner of the registered trademark Midol®.  
REV.00-022024 LB7069213146

## MENSTRUAL COMPLETE

acetaminophen, caffeine, pyrilamine maleate tablet

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:70692-131
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

<b>CAFFEINE</b> (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	60 mg
<b>PYRILAMINE MALEATE</b> (UNII: R35D29L3ZA) (PYRILAMINE - UNII:HPE317O9TL)	PYRILAMINE MALEATE	15 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>D&amp;C RED NO. 33</b> (UNII: 9DBA0SBB0L)	
<b>ETHYLCELLULOSE, UNSPECIFIED</b> (UNII: 7Z8S9VYZ4B)	
<b>FD&amp;C BLUE NO. 1 ALUMINUM LAKE</b> (UNII: J9EQA3S2JM)	
<b>GELATIN, UNSPECIFIED</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>ISOPROPYL ALCOHOL</b> (UNII: ND2M416302)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>STARCH, POTATO</b> (UNII: 8I089SAH3T)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>TRIACETIN</b> (UNII: XHX3C3X673)	

## Product Characteristics

<b>Color</b>	blue (Dark and Light Blue)	<b>Score</b>	no score
<b>Shape</b>	OVAL (oblong)	<b>Size</b>	18mm
<b>Flavor</b>		<b>Imprint Code</b>	S79
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70692-131-24	3 in 1 CARTON	04/05/2023	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:70692-131-46	72 in 1 BOTTLE; Type 0: Not a Combination Product	03/07/2024	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	04/05/2023	

Revised: 3/2024

Strive Pharmaceuticals Inc.