

FIBERCON- calcium polycarbophil tablet
Foundation Consumer Brands

Fibercon ®

Drug Facts

Active ingredient (in each caplet)

Calcium polycarbophil 625 mg equivalent to 500 mg polycarbophil

Purpose

Bulk-forming laxative

Uses

- relieves occasional constipation (irregularity)
- this product generally produces bowel movement in 12 to 72 hours

Warnings

Choking

Taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty in swallowing. If you experience chest pain, vomiting, or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.

Ask a doctor before use if you have

- abdominal pain, nausea, or vomiting
- a sudden change in bowel habits that lasts over a period of 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug. Take this product 2 or more hours before or after other drugs. All laxatives may affect how other drugs work.

When using this product do not use for more than 7 days unless directed by a doctor

Stop use and ask a doctor if rectal bleeding occurs or if you fail to have a bowel movement after use of this or any other laxative. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take each dose of this product with at least 8 ounces (a full glass) of water or other fluid. Taking this product without enough liquid may cause choking. See choking warning.
- FiberCon works naturally so continued use for one to three days is normally required to provide full benefit. Dosage may vary according to diet, exercise, previous laxative use or severity of constipation.
- adults and children 12 years of age and over: 2 caplets per dose. Do not take more than 8 caplets in 24 hours unless directed by a doctor.
- children under 12 years of age: ask a doctor

Other information

- **each caplet contains:** 140 mg calcium and 10 mg magnesium
- protect contents from moisture
- store at 20-25°C (68-77°F)

Inactive ingredients

caramel, crospovidone, hypromellose, magnesium stearate, microcrystalline cellulose, mineral oil, polyethylene glycol, silicon dioxide, sodium lauryl sulfate, stearic acid, talc

Questions or comments?

Call **1-888-594-0677** weekdays 9 AM to 5 PM EST

Distributed by: Foundation Consumer Brands, LLC
Pittsburgh, PA 15212

WARNING: Reproductive Harm - www.P65warnings.ca.gov

PRINCIPAL DISPLAY PANEL - 140 Caplet Bottle

NEW FORMULA

FiberCon®

Calcium Polycarbophil
Bulk-Forming Laxative

Simple Relief for Constipation

- Won't cause gas or bloating
- As effective as fiber powders

140 Coated Caplets

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take each dose of this product with at least 8 ounces (a full glass) of water or other fluid. Taking this product without enough liquid may cause choking. See choking warning.
- FiberCon works with your body so continued use for one to three days is normally required to provide full benefit. Dosage may vary according to diet, exercise, previous laxative use or severity of constipation.
- adults and children 12 years of age and over: 2 caplets per dose. Do not take more than 8 caplets in 24 hours unless directed by a doctor.
- children under 12 years of age: ask a doctor

Other information

- each caplet contains: 125 mg calcium
- protect contents from moisture
- store at 20-25°C (68-77°F)

Inactive ingredients caramel, crospovidone, hypromellose, magnesium stearate, microcrystalline cellulose, mineral oil, polyethylene glycol, silicon dioxide, sodium lauryl sulfate, stearic acid, talc

Questions or comments?
Call 1-888-594-0677 weekdays 9 AM to 5 PM EST

Distributed by: Foundation Consumer Brands, LLC
Pittsburgh, PA 15212
©2024 fibercon.com

WARNING:
Reproductive Harm –
www.P65Warnings.ca.gov.

NEW FORMULA

FiberCon[®]

CALCIUM POLYCARBOPHIL BULK-FORMING LAXATIVE

Simple Relief for Constipation

- Won't cause gas or bloating
- As effective as fiber powders

ACTUAL SIZE



TM

Bottle sealed with printed foil under cap.
Do Not Use if foil is torn or missing.

140 COATED CAPLETS

Drug Facts

Active ingredient (in each caplet) **Purpose**
Calcium polycarbophil 625 mg
equivalent to 500 mg polycarbophil Bulk-forming laxative

Uses

- relieves occasional constipation (irregularity)
- this product generally produces bowel movement in 12 to 72 hours

Warnings

Choking: Taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty in swallowing. If you experience chest pain, vomiting, or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.

Ask a doctor before use if you have

- abdominal pain, nausea, or vomiting
- a sudden change in bowel habits that lasts over a period of 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug. Take this product 2 or more hours before or after other drugs. All laxatives may affect how other drugs work.

When using this product do not use for more than 7 days unless directed by a doctor

Stop use and ask a doctor if rectal bleeding occurs or if you fail to have a bowel movement after use of this or any other laxative. These could be signs of a serious condition. ▶



R62820
07LA30013
052924

FIBERCON

calcium polycarbophil tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:80070-500
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CALCIUM POLYCARBOPHIL (UNII: 8F049NKY49) (POLYCARBOPHIL - UNII:W25LM17A4W)	CALCIUM POLYCARBOPHIL	625 mg

Inactive Ingredients

Ingredient Name	Strength
TALC (UNII: 7SEV7J4R1U)	
CARAMEL (UNII: T9D99G2B1R)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CROSPVIDONE (120 .MU.M) (UNII: 68401960MK)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
MINERAL OIL (UNII: T5L8T28FGP)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	brown	Score	2 pieces
Shape	OVAL	Size	18mm

Flavor		Imprint Code	LL;F1
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:80070-500-14	140 in 1 BOTTLE; Type 0: Not a Combination Product	04/04/2021	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	04/04/2021	

Labeler - Foundation Consumer Brands (117603632)

Revised: 9/2024

Foundation Consumer Brands