

PROTAMINE SULFATE - protamine sulfate injection, solution
Fresenius Kabi USA, LLC

Protamine Sulfate Injection, USP

For Intravenous Use

WARNING

Protamine sulfate can cause severe hypotension, cardiovascular collapse, noncardiogenic pulmonary edema, catastrophic pulmonary vasoconstriction, and pulmonary hypertension. Risk factors include high dose or overdose, rapid administration (see **WARNINGS** and **DOSAGE AND ADMINISTRATION**), repeated doses, previous administration of protamine, and current or previous use of protamine-containing drugs (NPH insulin, protamine zinc insulin, and certain beta-blockers). Allergy to fish, previous vasectomy, and severe left ventricular dysfunction and abnormal preoperative pulmonary hemodynamics also may be risk factors. In patients with any of these risk factors, the risk to benefit of administration of protamine sulfate should be carefully considered. Vasopressors and resuscitation equipment should be immediately available in case of a severe reaction to protamine. Protamine sulfate should not be given when bleeding occurs without prior heparin use.

DESCRIPTION:

Protamines are simple proteins of low molecular weight that are rich in arginine and strongly basic. They occur in the sperm of salmon and certain other species of fish.

Protamine sulfate occurs as fine white or off-white amorphous or crystalline powder. It is sparingly soluble in water. The pH is between 6.0 and 7.0. The cationic hydrogenated protamine at a pH of 6.8 to 7.1 reacts with anionic heparin at a pH of 5.0 to 7.5 to form an inactive complex.

Protamine Sulfate Injection, USP is a sterile, isotonic solution of protamine sulfate. It acts as a heparin antagonist. It is also a weak anticoagulant.

Each mL contains: Protamine sulfate 10 mg; sodium chloride 9 mg; Water for Injection q.s. Sulfuric acid and/or dibasic sodium phosphate (heptahydrate) may have been added for pH adjustment.

The preparation is preservative free.

Protamine sulfate is administered intravenously.

CLINICAL PHARMACOLOGY:

When administered alone, protamine has an anticoagulant effect. However, when it is given in the presence of heparin (which is strongly acidic), a stable salt is formed and the anticoagulant activity of both drugs is lost.

Protamine sulfate has a rapid onset of action. Neutralization of heparin occurs within five minutes after intravenous administration of an appropriate dose of protamine sulfate. Although the metabolic fate of the heparin-protamine complex has not been elucidated, it has been postulated that protamine sulfate in the heparin-protamine complex may be partially metabolized or may be attacked by fibrinolysin, thus freeing heparin.

INDICATIONS AND USAGE:

Protamine Sulfate Injection, USP is indicated in the treatment of heparin overdosage.

CONTRAINDICATIONS:

Protamine sulfate is contraindicated in patients who have shown previous intolerance to the drug.

WARNINGS:

Hyperheparinemia or bleeding has been reported in experimental animals and in some patients 30 minutes to 18 hours after cardiac surgery (under cardiopulmonary bypass) in spite of complete neutralization of heparin by adequate doses of protamine sulfate at the end of the operation. It is important to keep the patient under close observation after cardiac surgery. Additional doses of protamine sulfate should be administered if indicated by coagulation studies, such as the heparin titration test with protamine and the determination of plasma thrombin time.

Too-rapid administration of protamine sulfate can cause severe hypotensive and anaphylactoid reactions (see **DOSAGE AND ADMINISTRATION** and **WARNINGS**). Facilities to treat shock should be available.

PRECAUTIONS:

General

Because of the anticoagulant effect of protamine, it is unwise to give more than 50 mg over a short period unless a larger dose is clearly needed.

Patients with a history of allergy to fish may develop hypersensitivity reactions to protamine, although to date no relationship has been established between allergic reactions to protamine and fish allergy.

Previous exposure to protamine can induce a humoral immune response and predispose susceptible individuals to the development of untoward reactions from the subsequent use of this drug. Patients exposed to protamine through the use of protamine-containing insulin or during heparin neutralization may experience life-threatening reactions and fatal anaphylaxis upon receiving large doses of protamine intravenously. Severe reactions to intravenous protamine can occur in the absence of local or systemic allergic reactions to subcutaneous injection of protamine-containing insulin. Reports of the presence of antiprotamine antibodies in the sera of infertile or

vasectomized men suggest that some of these individuals may react to the use of protamine sulfate.

Fatal anaphylaxis has been reported in one patient with no prior history of allergies.

Drug Interactions

Protamine sulfate has been shown to be incompatible with certain antibiotics, including several of the cephalosporins and penicillins (see **DOSAGE AND ADMINISTRATION**).

Carcinogenesis, Mutagenesis, Impairment of Fertility

Studies have not been performed to determine potential for carcinogenicity, mutagenicity or impairment of fertility.

Pregnancy

Pregnancy Category C

Animal reproduction studies have not been conducted with protamine sulfate. It is also not known whether protamine sulfate can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Protamine sulfate should be given to a pregnant woman only if clearly needed.

Nursing Mothers

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when protamine sulfate is administered to a nursing woman.

Pediatric Use

Safety and effectiveness in children have not been established.

ADVERSE REACTIONS:

To report SUSPECTED ADVERSE REACTIONS, contact Fresenius Kabi USA, LLC at 1-800-551-7176 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

The intravenous administration of protamine sulfate may cause a sudden fall in blood pressure and bradycardia. Other reactions include transitory flushing and feeling of warmth, dyspnea, nausea, vomiting and lassitude. Back pain has been reported in conscious patients undergoing such procedures as cardiac catheterization.

Severe adverse reactions have been reported including: (1) Anaphylaxis that resulted in severe respiratory distress, circulation collapse and capillary leak (see **PRECAUTIONS**). Fatal anaphylaxis has been reported in one patient with no prior history of allergies; (2) Anaphylactoid reactions with circulatory collapse, capillary leak, and noncardiogenic pulmonary edema; acute pulmonary hypertension.

Complement activation by the heparin-protamine complexes, release of lysosomal enzymes from neutrophils, and prostaglandin and thromboxane generation have been associated with the development of anaphylactoid reactions.

Severe and potentially irreversible circulatory collapse associated with myocardial failure

and reduced cardiac output can also occur. The mechanism(s) of this reaction and the role played by concurrent factors are unclear.

High-protein, noncardiogenic pulmonary edema associated with the use of protamine has been reported in patients on cardiopulmonary bypass who are undergoing cardiovascular surgery. The etiologic role of protamine in the pathogenesis of this condition is uncertain, and multiple factors have been present in most cases. The condition has been reported in association with administration of certain blood products, other drugs, cardiopulmonary bypass alone, and other etiologic factors. It is difficult to treat, and it can be life-threatening. Because fatal anaphylactic and anaphylactoid reactions have been reported after the administration of protamine sulfate, the drug should be given only when resuscitation techniques and treatment of anaphylactic and anaphylactoid shock are readily available.

OVERDOSAGE:

Signs and Symptoms

Overdose of protamine sulfate may cause bleeding. Protamine has a weak anticoagulant effect due to an interaction with platelets and with many proteins including fibrinogen. This effect should be distinguished from the rebound anticoagulation that may occur 30 minutes to 18 hours following the reversal of heparin with protamine.

Rapid administration of protamine is more likely to result in bradycardia, dyspnea, a sensation of warmth, flushing, and severe hypotension. Hypertension has also occurred.

The median lethal dose of protamine sulfate is 50 mg/kg in mice. Serum concentrations of protamine sulfate are not clinically useful. Information is not available on the amount of drug in a single dose that is associated with overdose or is likely to be life-threatening.

Treatment

To obtain up-to-date information about the treatment of overdose, a good resource is your certified Regional Poison Control Center. Telephone numbers of certified poison control centers are listed in the *Physicians' Desk Reference (PDR)*. In managing overdose, consider the possibility of multiple drug overdoses, interaction among drugs and unusual drug kinetics in your patient.

Replace blood loss with blood transfusions or fresh frozen plasma.

If the patient is hypotensive, consider fluids, epinephrine, dobutamine or dopamine.

DOSAGE AND ADMINISTRATION:

Each mg of protamine sulfate, calculated on the dried basis, neutralizes not less than 100 USP Heparin Units.

Protamine sulfate injection should be given by very slow intravenous injection over a 10-minute period in doses not to exceed 50 mg (see WARNINGS).

Protamine sulfate is intended for injection without further dilution; however, if further dilution is desired, D5-W or normal saline may be used. Diluted solutions should not be stored since they contain no preservative.

Protamine sulfate should not be mixed with other drugs without knowledge of their compatibility, because protamine sulfate has been shown to be incompatible with certain antibiotics, including several of the cephalosporins and penicillins.

Because heparin disappears rapidly from the circulation, the dose of protamine sulfate required also decreases rapidly with the time elapsed following intravenous injection of heparin. For example, if the protamine sulfate is administered 30 minutes after the heparin, one-half the usual dose may be sufficient.

The dosage of protamine sulfate should be guided by blood coagulation studies (see **WARNINGS**).

Parenteral drug products should be visually inspected for particulate matter and discoloration prior to administration, whenever solution and container permit.

HOW SUPPLIED:

Product Code	Unit of Sale	Strength	Each
NP22905	NDC 63323-229-15 Unit of 25	50 mg per 5 mL (10 mg per mL)	NDC 63323-229-21 5 mL Single Dose Flip-top Vial
NP22930	NDC 63323-229-35 Individually Packaged	250 mg per 25 mL (10 mg per mL)	NDC 63323-229-35 25 mL Single Dose Flip-top Vial

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Do not permit to freeze.

CAUTION: The total dose of protamine sulfate contained in product No. NP22930 (250 mg in 25 mL) is 5 times greater than in product No. NP22905 (50 mg in 5 mL).

The large size 25 mL Vial is designed for antiheparin treatment only when large doses of heparin have been given during surgery and are to be neutralized by large doses of protamine sulfate after surgical procedures.

Novaplus is a registered trademark of Vizient, Inc.

Manufactured by:

Fresenius Kabi
Lake Zurich, IL 60047

www.fresenius-kabi.com/us

451124D

Revised: January 2024



PACKAGE LABEL - PRINCIPAL DISPLAY - Protamine 25mL Single Dose Vial Label

NDC 63323-229-21 NP22905

Protamine Sulfate

Injection, USP

For Intravenous Use Only

50 mg per 5 mL (10mg per mL)

5mL

Single Dose Vial

Rx only


Novaplus is a registered

Trademark of Vizient, Inc.

Sterile, Nonpyrogenic
Preservative Free.
Discard unused portion.
Each mL contains: Protamine sulfate 10 mg;
sodium chloride 9 mg; Water for Injection q.s.
Sulfuric acid and/or dibasic sodium
phosphate (heptahydrate) may have been
added for pH adjustment.
Usual Dosage: See insert.
Each mg of protamine sulfate, calculated
on the dried basis, neutralizes not less than
100 USP Heparin Units.

NDC 63323-229-21 NP22905
Protamine Sulfate Injection, USP
For Intravenous Use Only 50 mg per 5 mL (10 mg per mL)
5 mL Single Dose Vial Rx only
Novaplus is a registered trademark of Vizient, Inc.
novaplus

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].
Do not permit to freeze.
Mfd. by: **Fresenius Kabi**
Lake Zurich, IL 60047



3 63323-229-21 4
402417D
LOT/EXP

PACKAGE LABEL - PRINCIPAL DISPLAY - Protamine 25mL Single Dose Tray Label

NDC 63323-229-15 NP22905

Protamine Sulfate

Injection, USP

For Intravenous Use Only

50 mg per 5 mL (10mg per mL)

25 x 5 mL

Single Dose Vial

Rx only

Novaplus is a registered
Trademark of Vizient, Inc

Sterile, Nonpyrogenic
Preservative Free.
Discard unused portion.
Each mL contains: Protamine sulfate 10 mg;
sodium chloride 9 mg; Water for Injection q.s.
Sulfuric acid and/or dibasic sodium phosphate
(heptahydrate) may have been added for
pH adjustment.
Usual Dosage: See insert.
Each mg of protamine sulfate, calculated on the
dried basis, neutralizes not less than 100 USP
Heparin Units.

NDC 63323-229-15 NP22905


**Protamine
Sulfate
Injection, USP**

**For
Intravenous
Use Only** **50 mg per 5 mL
(10 mg per mL)**

25 x 5 mL
Single Dose Vials
Rx only
Novaplus is a registered trademark of
Vizient, Inc.
novaplus+

**Store at 20° to 25°C (68° to 77°F) [see USP
Controlled Room Temperature].**
Do not permit to freeze.
Mfd. by:
Fresenius Kabi
Lake Zurich, IL 60047

42901B



(01)20363323229157

**PACKAGE LABEL - PRINCIPAL DISPLAY - Protamine 25mL Single Dose Vial
Label**

NDC 63323-229-35 NP22930

Protamine Sulfate

Injection, USP

For Intravenous Use Only

250 mg per 25 mL (10mg per mL)

25 mL

Single Dose Vial

Rx only

Novaplus is a registered
Trademark of Vizient, Inc

NDC 63323-229-35 NP22930

Sterile, Nonpyrogenic
Preservative Free.
Discard unused portion.
Each mL contains: Protamine sulfate 10 mg; sodium chloride 9 mg; Water for Injection q.s. Sulfuric acid and/or dibasic sodium phosphate (heptahydrate) may have been added for pH adjustment.
Usual Dosage: See insert.
Each mg of protamine sulfate, calculated on the dried basis, neutralizes not less than 100 USP Heparin Units.

Protamine Sulfate Injection, USP

For Intravenous Use Only
250 mg per 25 mL (10 mg per mL)

25 mL
Single Dose Vial

Rx only

Novaplus is a registered trademark of Vizient, Inc.



Store at 20° to 25° C (68° to 77° F) [see USP Controlled Room Temperature].
Do not permit to freeze.

Mfd. by:
Fresenius Kabi
Lake Zurich, IL 60047



3 63323-229-35 1
402418D

LOT/EXP

PACKAGE LABEL - PRINCIPAL DISPLAY - Protamine 25mL Single Dose Vial Label

NDC 63323-229-35 NP22930

Protamine Sulfate

Injection, USP

For Intravenous Use Only

250 mg per 25 mL (10mg per mL)

25 mL

Single Dose Vial

Rx only

GTIN: 00363323229351



Sterile, Nonpyrogenic
Preservative Free.
 Discard unused portion.

Each mL contains:
 Protamine sulfate 10 mg;
 sodium chloride 9 mg; Water
 for Injection q.s. Sulfuric acid
 and/or dibasic sodium
 phosphate (heptahydrate)
 may have been added for
 pH adjustment.

Usual Dosage: See insert.

Each mg of protamine sulfate,
 calculated on the dried basis,
 neutralizes not less than
 100 USP Heparin Units.

Novaplust is a registered trademark of
 Vizient, Inc.


Mfd. by:
Fresenius Kabi
 Lake Zurich, IL 60047

Store at 20° to 25°C (68° to
 77°F) [see USP Controlled
 Room Temperature].

Do not permit to freeze.

NDC 63323-229-35 NP22930


**Protamine
 Sulfate**
 Injection, USP



25 mL

Single Dose Vial

Rx only



novaplust

621003D

PROTAMINE SULFATE

protamine sulfate injection, solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:63323-229
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PROTAMINE SULFATE (UNII: 0DE9724IHC) (PROTAMINE - UNII:72G3UY6T4N)	PROTAMINE SULFATE	10 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	9 mg in 1 mL
SULFURIC ACID (UNII: O40UQP6WCF)	
SODIUM PHOSPHATE, DIBASIC (UNII: GR686LBA74)	
WATER (UNII: 059QF0KO0R)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63323-229-15	25 in 1 TRAY	10/18/2000	
1	NDC:63323-229-21	5 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		
2	NDC:63323-229-35	1 in 1 BOX	10/18/2000	
2		25 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		
3	NDC:63323-229-26	10 in 1 TRAY	10/18/2000	11/30/2024
3		25 mL in 1 VIAL, SINGLE-DOSE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA089454	10/18/2000	

Labeler - Fresenius Kabi USA, LLC (608775388)

Establishment			
Name	Address	ID/FEI	Business Operations
Fresenius Kabi USA, LLC		023648251	ANALYSIS(63323-229) , MANUFACTURE(63323-229)

Revised: 3/2024

Fresenius Kabi USA, LLC