

## **CONAZOL ANTIFUNGAL- tolnaftate solution/ drops**

**Marcus USA**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Conazol®  
antifungal**

### ***Drug Facts***

#### **Active Ingredients**

Tolnaftate 1%

#### **Purposes**

Antifungal

#### **Uses**

For effective treatment of athlete's foot, jock itch and ringworm. Not for relief of infected finger nails and toe nails or effective against bacteria or viruses.

#### **Warnings**

##### **For external use only**

- Avoid contact with eyes.
- **Do not use** on children under 2 years of age unless directed by a doctor.

##### **Stop and ask a doctor if**

- If treating athlete's foot and ringworm: If irritation occurs or if there is no improvement within 4 weeks.
- If treating jock itch: If irritation occurs or if there is no improvement within 2 weeks.

##### **Keep out of the reach of children**

- If swallowed, get medical help or contact a Poison Control Center right away.

#### **Directions**

Clean or wash the affected area and dry thoroughly. Apply a thin layer of the product over the affected area twice daily (morning and night) or as directed by a doctor. Supervise children in the use of this product. For athlete's foot: Pay special attention to spaces between toes; wear well-fitting, ventilated shoes, and change socks at least once daily. For athlete's foot and ringworm use daily for 4 weeks. If condition persists longer, consult a doctor. This product is not effective on the scalp or nails.

#### **Other Information**

- Store at room temperature 15°-30°C (59°-86°F)
- Questions or comments, call (800) 428-9489

**Inactive Ingredients**

Purified water, PEG-8, Cetearyl alcohol and Ceteth-20 phosphate and decetyl phosphate, Propylene glycol, Cocamidopropyl betaine, Octoxynol-9, Glyceryl stearate, Stearyl alcohol, Hydroxyethyl cellulose, Imidazolidinyl urea, Cetyl alcohol, Methylparaben, Propylparaben, Triethanolamine, Citric acid

Distributed by: MarcasUSA, LLC  
El Segundo CA, 90245

**PRINCIPAL DISPLAY PANEL - 30 ML Bottle Carton**

***NOTHING MORE EFFECTIVE***

**Conazol<sup>®</sup>**

ANTIFUNGAL

**TOE FUNGUS**

**ELIMINATOR**

***with Tolnaftate 1%***

**Cures &**

**Prevents**

**Fungus**

**1 FL OZ (30 ML)**

NDC: 75940-126-01



**MAXIMUM  
STRENGTH  
WITHOUT A  
PRESCRIPTION**

**BEGINS WORKING  
ON CONTACT**

**NOTHING MORE EFFECTIVE**



**TOE FUNGUS  
ELIMINATOR**

**ELIMINADOR DE HONGOS DEL DEDO DEL PIE  
with Tolnaftate 1%**

**Cures &  
Prevents  
Fungus**

**1 FLOZ (30 ML)**

**JUST 2 SIMPLE STEPS**

*Solo dos simples pasos*

**Cures & prevents fungus on  
the skin around, adjacent to  
and under the nails!**



**STEP 1/PASO 1:**

**Wash affected area with water and  
soap and dry thoroughly.**

*Lave el área afectada  
con agua y jabón y séquelo bien.*



**STEP 2/PASO 2:**

**Apply Conazol Toe Fungus Eliminator  
to affected area allowing it to absorb.**

*Aplique Conazol Toe Fungus Eliminator  
al área afectada permitiendo que se  
absorba.*

**NO-TOUCH APPLICATION**  
TO PREVENT THE  
SPREAD OF INFECTION

**NEW**



**Drug Facts**

**Active Ingredients**

Tolnaftate 1%

**Purposes**

Antifungal

**Uses** For effective treatment of athlete's foot, jock itch and ringworm. Not for relief of infected finger nails and toe nails or effective against bacteria or viruses.

**Warnings**

For external use only

- Avoid contact with eyes. • **Do not use** on children under 2 years of age unless directed by a doctor.

**Stop and ask a doctor if**

- If treating athlete's foot and ringworm: If irritation occurs or if there is no improvement within 4 weeks. • If treating jock itch: If irritation occurs or if there is no improvement within 2 weeks.

**Keep out of the reach of children**

- If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

Clean or wash the affected area and dry thoroughly. Apply a thin layer of the product over the affected area twice daily (morning and night) or as directed by a doctor. Supervise children in the use of this product. For athlete's foot: Pay special attention to spaces between toes; wear well-fitting, ventilated shoes, and change socks at least once daily. For athlete's foot and ringworm use daily for 4 weeks. If condition persists longer, consult a doctor. This product is not effective on the scalp or nails.

**Other Information**

- Store at room temperature 15°-30°C (59°-86°F)
- Questions or comments, call (800) 428-9489

**Inactive Ingredients**

Purified water, PEG-8, Cetearyl alcohol and Ceteth-20 phosphate and decetyl phosphate, Propylene glycol, Cocamidopropyl betaine, Octoxynol-9, Glyceryl stearate, Stearyl alcohol, Hydroxyethyl cellulose, Imidazolidinyl urea, Cetyl alcohol, Methylparaben, Propylparaben, Triethanolamine, Citric acid

**MADE IN U.S.A.**

Distributed by: MarcasUSA, LLC

**TOE FUNGUS  
ELIMINATOR**

**NO-TOUCH APPLICATOR**



**No-contact application prevents spreading of fungus infection to other toes.**

*La aplicación sin contacto previene la propagación de la infección de hongos a otros dedos del pie.*

**1 FL OZ (30 ML)**



El Segundo CA, 90245

**ELIMINATOR**  
ELIMINADOR DE HONGOS DEL DEDO DEL PIE

Under license by Laboratorios Liomont, S.A. de C.V.

CON1002A0407



## CONAZOL ANTIFUNGAL

tolnaftate solution/ drops

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:75940-126
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>Tolnaftate</b> (UNII: 06KB629TKV) (Tolnaftate - UNII:06KB629TKV)	Tolnaftate	10 mg in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>water</b> (UNII: 059QF0K00R)	
<b>POLYETHYLENE GLYCOL 400</b> (UNII: B697894SGQ)	
<b>CETOSTEARYL ALCOHOL</b> (UNII: 2DMT128MIS)	
<b>CETETH-20 PHOSPHATE</b> (UNII: 921FTA1500)	
<b>DIETHANOLAMINE CETYL PHOSPHATE</b> (UNII: 4UG0316V9S)	
<b>propylene glycol</b> (UNII: 6DC9Q167V3)	
<b>cocamidopropyl betaine</b> (UNII: 5OCF3011KX)	
<b>octoxynol-9</b> (UNII: 7JPC6Y25QS)	
<b>GLYCERYL MONOSTEARATE</b> (UNII: 230OU9XXE4)	
<b>Stearyl alcohol</b> (UNII: 2KR89I4HIY)	
<b>HYDROXYETHYL CELLULOSE (3000 MPAS AT 1%)</b> (UNII: 7Q6P4JN1QT)	
<b>IMIDUREA</b> (UNII: M629807ATL)	
<b>CETYL ALCOHOL</b> (UNII: 936JST6JCN)	
<b>methylparaben</b> (UNII: A218C7HI9T)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	
<b>TROLAMINE</b> (UNII: 9O3K93S3TK)	
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:75940-126-01	1 in 1 CARTON	10/29/2017	
1		30 mL in 1 BOTTLE, DROPPER; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333C	10/29/2017	

**Labeler** - MarcusUSA (016139820)

## Establishment

Name	Address	ID/FEI	Business Operations
Private Label Partners		046033481	LABEL(75940-126)

Revised: 1/2020

MarcusUSA