

CHILDRENS DIMETAPP COLD AND COUGH- brompheniramine maleate, dextromethorphan hbr, phenylephrine hcl liquid

GlaxoSmithKline Consumer Healthcare Holdings (US) LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

ACTIVE INGREDIENTS (IN EACH 10 ML)

Brompheniramine maleate, USP 2 mg

Dextromethorphan HBr, USP 10 mg

Phenylephrine HCl , USP 5 mg

PURPOSES

Antihistamine

Cough suppressant

Nasal decongestant

USES

- temporarily relieves cough due to minor throat and bronchial irritation occurring with a cold, and nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves these symptoms due to hay fever (allergic rhinitis):
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat
- temporarily restores freer breathing through the nose

WARNINGS

Do not use

- to sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure

- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or persistent or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

- taking any other oral nasal decongestant or stimulant
- taking sedatives or tranquilizers

When using this product

- **do not use more than directed**
- may cause marked drowsiness
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

DIRECTIONS

- do not take more than 6 doses in any 24-hour period
- measure only with dosage cup provided
- keep dosage cup with product
- ml = milliliter

age	dose
adults and children 12 years and over	20 ml every 4 hours
children 6 to under 12 years	10 ml every 4 hours
children under 6 years	do not use

OTHER INFORMATION

- each 10 ml contains: **sodium 6 mg**
- store at 20-25°C (68-77°F)

INACTIVE INGREDIENTS

anhydrous citric acid, artificial flavor, FD&C blue no. 1, FD&C red no. 40, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

QUESTIONS OR COMMENTS?

Call weekdays from 9 AM to 5 PM EST at **1-800-762-4675**

PRINCIPAL DISPLAY PANEL

NDC 0031-2234-13

**Children's
Dimetapp®**

**BROMPHENIRAMINE MALEATE (Antihistamine)
DEXTROMETHORPHAN HBr (Cough Suppressant)
PHENYLEPHRINE HCl (Nasal Decongestant)**

**Cold
& Cough**

Relieves Nasal Symptoms

- Stuffy Nose**
- Runny Nose**
- Sneezing**

Plus Other Symptoms

- Itchy, Watery Eyes**
- Cough**

**For Ages
6 Yrs.
& Over**

4 FL OZ (118 ml) alcohol-free • grape flavor

Children's Dimetapp®

BROMPHENIRAMINE MALEATE (Antihistamine)
 DEXTROMETHORPHAN HBr (Cough Suppressant)
 PHENYLEPHRINE HCl (Nasal Decongestant)

Cold & Cough

Children's Dimetapp®

BROMPHENIRAMINE MALEATE (Antihistamine)
 DEXTROMETHORPHAN HBr (Cough Suppressant)
 PHENYLEPHRINE HCl (Nasal Decongestant)



Cold & Cough

Relieves Nasal Symptoms

- Stuffy Nose
- Runny Nose
- Sneezing

Plus Other Symptoms

- Itchy, Watery Eyes
- Cough



For Ages
**6 Yrs.
 & Over**

4 FL OZ (118 ml) alcohol-free • grape flavor

Children's Dimetapp®

Drug Facts

Active ingredients (in each 10 ml)

Brompheniramine maleate, USP 2 mg Antihistamine
 Dextromethorphan HBr, USP 10 mg...Cough suppressant
 Phenylephrine HCl, USP 5 mg.....Nasal decongestant

Purposes

Uses

- temporarily relieves cough due to minor throat and bronchial irritation occurring with a cold, and nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves these symptoms due to hay fever (allergic rhinitis):
 - runny nose
 - sneezing
 - itchy, watery eyes
 - itching of the nose or throat
- temporarily restores freer breathing through the nose

Warnings

Do not use

- to sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or persistent or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

- taking any other oral nasal decongestant or stimulant
- taking sedatives or tranquilizers

LOT:
 EXP:

Do Not Use if breakable ring is separated or missing.

Packaged with Tamper-Evident bottle cap.

Children's Dimetapp®

Drug Facts (continued)

When using this product

- do not use more than directed
- may cause marked drowsiness
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosage cup provided
- keep dosage cup with product
- ml = milliliter

age*	dose
adults and children 12 years and over	20 ml every 4 hours
children 6 to under 12 years	10 ml every 4 hours
children under 6 years	do not use

Other information

- each 10 ml contains: sodium 6 mg
- store at 20-25°C (68-77°F)

Inactive ingredients anhydrous citric acid, artificial flavor, FD&C blue no. 1, FD&C red no. 40, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments? Call weekdays from 9 AM to 5 PM EST at 1-800-762-4675

Children's Dimetapp® Cold & Cough

For most recent product information, visit www.dimetapp.com

We pledge to you that Dimetapp® products contain only high quality ingredients and meet strict standards of quality and safety. You can trust Dimetapp® products for your family.

Dosage Cup Provided 

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

Distributed by:
Pfizer, Madison, NJ 07940 USA
© 2018 Pfizer Inc.
U.S. Patent 9,801,941



3 0031 2234 13 2

PAA105829

CHILDRENS DIMETAPP COLD AND COUGH

brompheniramine maleate, dextromethorphan hbr, phenylephrine hcl liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0031-2234
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BROMPHENIRAMINE MALEATE (UNII: IXA7C9ZN03) (BROMPHENIRAMINE - UNII:H57G17P2FN)	BROMPHENIRAMINE MALEATE	1 mg in 5 mL

DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	5 mg in 5 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	2.5 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	PURPLE (purple liquid)	Score	
Shape		Size	
Flavor	GRAPE (grape flavor and odor)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0031-2234-13	1 in 1 CARTON	05/23/2006	
1		118 mL in 1 BOTTLE; Type 1: Convenience Kit of Co-Package		
2	NDC:0031-2234-19	1 in 1 CARTON	05/23/2006	
2		237 mL in 1 BOTTLE; Type 1: Convenience Kit of Co-Package		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	05/23/2006	

Labeler - GlaxoSmithKline Consumer Healthcare Holdings (US) LLC (079944263)