

# **COLD AND ALLERGY- phenylephrine hcl and chlorpheniramine maleate tablet**

## **Allegiant Health**

-----

### **Drug Facts**

#### **Active ingredients (in each tablet)**

Phenylephrine HCl 10 mg  
Chlorpheniramine maleate 4mg

#### **Purpose**

Nasal Decongestant  
Antihistamine

#### **Uses**

temporarily relieves nasal congestion due to the common cold, and these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- sneezing
- itching of the nose or throat
- itchy, watery eyes
- nasal congestion
- sinus congestion and pressure

#### **Warnings**

##### **Do Not Use**

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

##### **Ask a doctor before use if you have**

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland
- diabetes
- heart disease
- high blood pressure
- thyroid disease

##### **Ask a doctor or pharmacist before use if you are**

taking sedatives or tranquilizers

**When using this product**

- **do not use more than directed**
- may cause excitability, especially in children
- may cause drowsiness; alcohol, sedatives or tranquilizers may increase drowsiness
- avoid alcoholic beverages and use caution when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or are accompanied by fever

**If pregnant or breast-feeding**

ask a health professional before use.

**Keep out of reach of children**

In case of accidental overdose, contact a doctor or Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical even if you do not notice any signs or symptoms.

**Directions**

**Adults and children 12 years of age and older:** 1 tablet every 4 to 6 hours. Do not take more than 6 doses in 24 hours

**Children under 12 years of age:** Consult a doctor

**Other Information**

- store at room temperature
- do not use if blister unit is torn or open

**Inactive Ingredients**

croscarmellose sodium, lactose anhydrous, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid

**Package/Label Principal Display Panel**

HealthA2Z® **COLD & ALLERGY**

HealthA2Z®

**PSEUDOEPHEDRINE  
FREE FORMULA**

**COLD & ALLERGY**

*Chlorpheniramine Maleate and Phenylephrine HCl*  
*Antihistamine / Nasal Decongestant*

- Nasal Congestion • Runny Nose
- Sneezing • Itchy, Watery Eyes



**24 TABLETS**

Lot:  
Exp:



**COLD &  
ALLERGY**

HealthA2Z®



Distributed by:

<b>Drug Facts</b> (continued)	
use caution when driving a motor vehicle or operating machinery	
<b>Active ingredients (in each tablet)</b>	
Chlorpheniramine maleate 4mg.....Antihistamine	Phenylephrine HCl 10mg.....Nasal decongestant
<b>Uses</b> temporarily relieves nasal congestion due to the common cold, and these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itching of the nose or throat ■ itchy, watery eyes ■ nasal congestion ■ sinus congestion and pressure	
<b>Warnings</b>	
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.	
Ask a doctor before use if you have ■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma ■ trouble urinating due to an enlarged prostate gland ■ diabetes ■ heart disease ■ high blood pressure ■ thyroid disease	
Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers	
When using this product ■ do not use more than directed ■ may cause excitability, especially in children ■ may cause drowsiness; alcohol, sedatives or tranquilizers may increase drowsiness ■ avoid alcoholic beverages and	
<b>Drug Facts</b> (continued)	
<b>Purpose</b>	
Chlorpheniramine maleate 4mg.....Antihistamine	
Phenylephrine HCl 10mg.....Nasal decongestant	
<b>Uses</b> temporarily relieves nasal congestion due to the common cold, and these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itching of the nose or throat ■ itchy, watery eyes ■ nasal congestion ■ sinus congestion and pressure	
<b>Warnings</b>	
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.	
Ask a doctor before use if you have ■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma ■ trouble urinating due to an enlarged prostate gland ■ diabetes ■ heart disease ■ high blood pressure ■ thyroid disease	
Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers	
When using this product ■ do not use more than directed ■ may cause excitability, especially in children ■ may cause drowsiness; alcohol, sedatives or tranquilizers may increase drowsiness ■ avoid alcoholic beverages and	
<b>Drug Facts</b> (continued)	
<b>Purpose</b>	
Chlorpheniramine maleate 4mg.....Antihistamine	
Phenylephrine HCl 10mg.....Nasal decongestant	
<b>Uses</b> temporarily relieves nasal congestion due to the common cold, and these symptoms due to hay fever or other upper respiratory allergies: ■ runny nose ■ sneezing ■ itching of the nose or throat ■ itchy, watery eyes ■ nasal congestion ■ sinus congestion and pressure	
<b>Warnings</b>	
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.	
Ask a doctor before use if you have ■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma ■ trouble urinating due to an enlarged prostate gland ■ diabetes ■ heart disease ■ high blood pressure ■ thyroid disease	
Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers	
When using this product ■ do not use more than directed ■ may cause excitability, especially in children ■ may cause drowsiness; alcohol, sedatives or tranquilizers may increase drowsiness ■ avoid alcoholic beverages and	
<b>Directions</b>	
Adults and children 12 years of age and older	1 tablet every 4 to 6 hours. Do not take more than 6 tablets in 24 hours.
Children under 12 years of age	Consult a doctor.
<b>Other information</b> ■ store at 20°-25°C (68°-77°F) ■ do not use if blister unit is torn or open	
<b>Inactive ingredients</b> croscarmellose sodium, lactose anhydrous, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid	
Distributed by: Allegiant Health Deer Park, NY 11729	

**Cold and Allergy**

# COLD AND ALLERGY

phenylephrine hcl and chlorpheniramine maleate tablet

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69168-406
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	4 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	7mm
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69168-406-93	1 in 1 CARTON	12/23/2014	
1		24 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/23/2014	

**Labeler** - Allegiant Health (079501930)

