

MOTION SICKNESS RELIEF- dimenhydrinate 50mg tablet
Allegiant Health

Dramamine Original Formula

Active ingredient (in each tablet)

Dimenhydrinate 50 mg

Purpose

Antiemetic

Use

for prevention and treatment of these symptoms associated with motion sickness:

- nausea
- vomiting
- dizziness

Warnings

Do not use

for children under 2 years of age unless directed by a doctor

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers

When using this product

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase dizziness
- be careful when driving a motor vehicle or operating machinery

If pregnant or breast-feeding,

ask a doctor before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-

800-222-1222) right away.

Directions

- to prevent motion sickness, the first dose should be taken $\frac{1}{2}$ to 1 hour before starting activity

Adults and children 12 years and over	<ul style="list-style-type: none">• 1 to 2 tablets every 4-6 hours• do not exceed 8 tablets in 24 hours, or as directed by a doctor
Children 6 to under 12 years	<ul style="list-style-type: none">• $\frac{1}{2}$ to 1 tablet every 6-8 hours• do not exceed 3 tablets in 24 hours, or as directed by a doctor
Children 2 to under 6 years	<ul style="list-style-type: none">• $\frac{1}{2}$ tablet every 6-8 hours• do not exceed 1 $\frac{1}{2}$ tablets in 24 hours, or as directed by a doctor

Other information

- each tablet contains: calcium 35mg
- do not use if blister unit is torn or open
- store at room temperature 68-77°F (20-25°C)
- protect from moisture

Inactive ingredients

croscarmellose sodium, dibasic calcium phosphate dihydrate, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid

Questions or comments?

1-888-952-0050 (Monday through Friday 9AM - 5PM EST)

PRINCIPAL DISPLAY PANEL

AH Distributed by:
Allegiant Health
Deer Park, NY 11729

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS
OPENED OR IF BLISTER UNIT IS TORN, BROKEN
OR SHOWS ANY SIGNS OF TAMPERING**

**NOT FOR HOUSEHOLDS
WITH YOUNG CHILDREN**

LB1712 R0719

Health**AzZ**[®]

*Compare to active ingredient in Dramamine[®]

Motion Sickness Relief



DIMENHYDRINATE 50mg

Antiemetic

- Prevents Nausea, Vomiting & Dizziness
- For Children & Adults



Actual Size

12
Tablets

Lot:
Exp:

Drug Facts (continued)
Inactive Ingredients croscarmellose sodium, dibasic calcium phosphate dihydrate, magnesium stearate, microcrystalline cellulose, silicon dioxide, stearic acid
Questions or comments? 1-888-952-0050 (Monday through Friday 9AM - 5PM EST)

Drug Facts
Active Ingredient (in each tablet) Dimenhydrinate 50 mgAntiemetic
Purpose
Uses for prevention and treatment of these symptoms associated with motion sickness: ■ nausea ■ vomiting ■ dizziness
Warnings
 Do not use for children under 2 years of age unless directed by a doctor
 ■ Ask a doctor before use if you have ■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma
 ■ trouble urinating due to an enlarged prostate gland
 ■ Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers
 When using this product ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ alcohol, sedatives, and tranquilizers may increase dizziness ■ be careful when driving a motor vehicle or operating machinery
 If pregnant or breast-feeding, ask a doctor before use.
 Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.
Directions
 ■ to prevent motion sickness, the first dose should be taken ½ to 1 hour before starting activity
Adults and children 12 years and over ■ 1 to 2 tablets every 4-6 hours ■ do not exceed 8 tablets in 24 hours, or as directed by a doctor
Children 6 to under 12 years ■ ½ to 1 tablet every 6-8 hours ■ do not exceed 3 tablets in 24 hours, or as directed by a doctor
Children 2 to under 6 years ■ ½ tablet every 6-8 hours ■ do not exceed 1 ½ tablets in 24 hours, or as directed by a doctor
Other Information ■ each tablet contains: calcium 35 mg ■ do not use if blister unit is torn or open ■ store at room temperature 68-77°F (20-25°C) ■ protect from moisture
 ▲

*This product is not manufactured or distributed by Prestige Consumer Healthcare Inc., distributor of Dramamine[®].



Motion Sickness Relief

MOTION SICKNESS RELIEF

dimenhydrinate 50mg tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69168-408
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIMENHYDRINATE (UNII: JB937PER5C) (8-CHLOROTHEOPHYLLINE - UNII:GE2UA340FM)	DIMENHYDRINATE	50 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
MAGNESIUM STEARATE (UNII: 70097M6130)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	8mm
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69168-408-86	1 in 1 CARTON	01/15/2012	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M009	01/15/2012	

Labeler - Allegiant Health (079501930)

