

DIMENHYDRINATE- dimenhydrinate tablet
Acme United Corporation

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

1006-Acme

Drug Facts

Active Ingredient (in each tablet)

Dimenhydrinate 50 mg

Purpose

Antiemetic

Uses

for prevention and treatment of nausea, vomiting or dizziness associated with motion sickness

Do not use in children under 2 years of age unless directed by a doctor

Ask doctor before use if you have

- breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland.

Ask doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact Poison Control Center (800-222-1222) right away.

Directions

To prevent motion sickness the first dose should be taken ½ to 1 hour before starting activity.

To prevent or treat motion sickness:

Adults and Children (12 years and older)	take 1-2 tablets every 4-6 hours, not more than 8 tablets in 24 hours, or as directed by doctor.
Children 6 to under 12 years	give 1 tablet every 6-8 hours, not more than 3 tablets in 24 hours or as directed by doctor.
Children 2 to under 6 years	give ¼ to ½ tablet every 6-8 hours, not more than 1½ tablets in 24 hours or as directed by a doctor

Other information

- each tablet contains **calcium 29 mg**
- store at room temperature
- do not use if packet is open or torn

croscarmellose sodium, dicalcium phosphate, magnesium stearate, microcrystalline cellulose

Questions?

1-800-835-2263

Retain carton for complete product information

Manufactured for:

Acme United Corporation

55 Walls Dr, Fairfield, CT 06824

www.FirstAidOnly.com

Made in the USA

©2018 Acme United Corporation

Physician Care®

Motion Sickness

Dimenhydrinate / Antiemetic

Nausea, Vomiting and Dizziness Relief

Tamper-Evident Packets of 2 Tablets

Do not use if packet is open or torn.

Dimenhydrinate 50mg

100 Tablets (50 Packets, 2 Tablets each)

The image displays the front and back of a box for PhysiciansCARE Motion Sickness tablets. The front panel features the PhysiciansCARE logo, the product name 'Motion Sickness', and the active ingredient 'Dimenhydrinate / Antiemetic'. It also lists the benefits: 'Nausea, Vomiting and Dizziness Relief'. The packaging is marked as 'Tamper-Evident' and contains '100 Tablets (50 Packets, 2 Tablets each)'. The back panel provides detailed drug facts, including the active ingredient (Dimenhydrinate 50 mg), purpose (Antiemetic), uses, warnings, directions for adults and children, and other information such as calcium content and inactive ingredients.

Active Ingredient (in each tablet)	Purpose
Dimenhydrinate 50 mg	Antiemetic

Uses
for prevention and treatment of nausea, vomiting or dizziness associated with motion sickness

Warnings
Do not use in children under 2 years of age unless directed by a doctor
Ask doctor before use if you have
• breathing problem such as emphysema or chronic bronchitis
• glaucoma
• trouble urinating due to an enlarged prostate gland.
Ask doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product
• marked drowsiness may occur
• avoid alcoholic drinks
• alcohol, sedatives and tranquilizers may increase drowsiness
• be careful when driving a motor vehicle or operating machinery, if pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children.
In case of overdose, get medical help or contact Poison Control Center (800 222 1222) right away.

Drug Facts (continued)
Directions
To prevent motion sicknesses the first dose should be taken 1/2 to 1 hour before starting activity.
To prevent or treat motion sicknesses:
Adults and children 12 years and older take 1 - 2 tablets every 4 - 6 hours, not more than 8 tablets in 24 hours, or as directed by doctor.
Children 6 to under 12 years give 1 tablet every 8 - 8 hours, not more than 3 tablets in 24 hours or as directed by doctor.
Children 2 to under 6 years give 1/2 to 1 tablet every 8 - 8 hours, not more than 1 1/2 tablets in 24 hours or as directed by a doctor.

Other information
• each tablet contains calcium 29 mg
• store at room temperature
• do not use if packet is open or torn

Inactive ingredients
croscarmellose sodium, dicalcium phosphate, magnesium stearate, microcrystalline cellulose

Questions? 1-800-835-2263

Contact Us 1-800-835-2263 www.FirstAidOnly.com

DIMENHYDRINATE
dimenhydrinate tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0924-1007
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIMENHYDRINATE (UNII: JB937PER5C) (CHLORTHEOPHYLLINE - UNII:GE2UA340 FM)	DIMENHYDRINATE	50 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	9mm
Flavor		Imprint Code	1006;1006
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0924-1007-02	6 in 1 BAG	04/18/2019	
1		2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:0924-1007-01	50 in 1 CARTON	04/18/2019	
2		2 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:0924-1007-03	100 in 1 BOTTLE; Type 0: Not a Combination Product	05/21/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part336	04/18/2019	

Labeler - Acme United Corporation (001180207)