

**ACETAMINOPHEN, DIPHENHYDRAMINE HYDROCHLORIDE- acetaminophen,
diphenhydramine hydrochloride tablet, film coated
Cardinal Health**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

0752-Cardinal

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 500 mg

Diphenhydramine HCl 25 mg

Purpose

Pain reliever

Nighttime sleep-aid

Uses

temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other product containing diphenhydramine, even one used on skin
- in children under 12 years of age
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- a breathing problem such as emphysema or chronic bronchitis

- trouble urinating due to an enlarged prostate gland
- glaucoma

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- avoid alcoholic beverages
- do not drive a motor vehicle or operate machinery
- drowsiness will occur

Stop use and ask a doctor if

- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness.
- new symptoms occur
- redness or swelling is present
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**
- **adults and children 12 years and over:** take 2 caplets at bedtime. Do not take more than 2 caplets of this product in 24 hours.
- **children under 12 years:** do not use

Other information

store at room temperature

croscarmellose sodium, D&C yellow #10 aluminum lake, FD&C blue #1 aluminum lake, hypromellose, magnesium stearate, mineral oil, polyvinylpyrrolidone, pregelatinized starch, silica, sodium starch glycolate, stearic acid, talc, titanium dioxide, triacetin

Questions or comments?

(800) 231-4670

*This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Tylenol® PM Extra Strength.

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS
BROKEN OR MISSING**

DIST BY CAH, DUBLIN, OH 43017

1-800-200-6313

© 2018 Cardinal Health

100% Money Back Guarantee

Return to place of purchase if not satisfied

LEADER TM

NDC 7000-0411-3

Extra Strength Acetaminophen PM

Acetaminophen, 500mg

Diphenhydramine HCl, 25mg

Pain Reliever/Nighttime Sleep-Aid

Non-Habit Forming

COMPARE TO TYLENOL [®] PM EXTRA STRENGTH active ingredients*

100 Caplets



TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING
CIN 5457288 REV. 10/18

12960-10-18

Drug Facts (continued)

■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- glaucoma

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- avoid alcoholic beverages
- do not drive a motor vehicle or operate machinery
- drowsiness will occur

Stop use and ask a doctor if

- sleeplessness persists continuously for more than 2 weeks. Insomnia may be a symptom of a serious underlying medical illness. ■ new symptoms occur
- redness or swelling is present ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Drug Facts

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Active ingredients (in each caplet)

Acetaminophen 500 mg Pain reliever
Diphenhydramine HCl 25 mg Nighttime sleep-aid

Purpose

Uses temporary relief of occasional headaches and minor aches and pains with accompanying sleeplessness

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other product containing diphenhydramine, even one used on skin ■ in children under 12 years of age
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

LEADER²

NDC 70000-0411-3

Extra Strength

Acetaminophen PM

Acetaminophen, 500 mg | Diphenhydramine HCl, 25 mg
Pain Reliever | Nighttime Sleep-Aid

COMPARE TO
TYLENOL[®] PM
EXTRA STRENGTH
active ingredients*

100% Money Back Guarantee

Non-Habit Forming

100 CAPLETS

Actual Size

Drug Facts (continued)

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)
- adults and children 12 years and over: take 2 caplets at bedtime. Do not take more than 2 caplets of this product in 24 hours.
- children under 12 years: do not use

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients croscarmellose sodium, D&C yellow #10 aluminum lake, FD&C blue #1 aluminum lake, hypromellose, magnesium stearate, mineral oil, polyvinylpyrrolidone, pregelatinized starch, silica, sodium starch glycolate, stearic acid, talc, titanium dioxide, triacetin

Questions or comments?
1-800-231-4670



DISTRIBUTED BY CARDINAL HEALTH
DUBLIN, OHIO 43017
www.myleader.com 1-800-200-0315
Essential to Care[™] since 1979



© 2018 Cardinal Health. All Rights Reserved. CARDINAL HEALTH, the Cardinal Health LOGO, LEADER, and the Leader LOGO are trademarks or registered trademarks of Cardinal Health. All other marks are the property of their respective owners.

*This product is not manufactured or distributed by McNeil Consumer Healthcare, owner of the registered trademark Tylenol[®] PM Extra Strength.

ACETAMINOPHEN, DIPHENHYDRAMINE HYDROCHLORIDE

acetaminophen, diphenhydramine hydrochloride tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70000-0411
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg

Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

TALC (UNII: 7SEV7J4R1U)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)
STEARIC ACID (UNII: 4ELV7Z65AP)
LIGHT MINERAL OIL (UNII: N6K5787QVP)
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)
MAGNESIUM STEARATE (UNII: 70097M6I30)
TRIACETIN (UNII: XHX3C3X673)
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)
POVIDONE K30 (UNII: U725QWY32X)
STARCH, CORN (UNII: O8232NY3SJ)
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)
HYPROMELLOSES (UNII: 3NXW29V3WO)
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)

Product Characteristics

Color	blue	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	CPC752
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70000-0411-2	50 in 1 BOTTLE; Type 0: Not a Combination Product	11/29/2018	
2	NDC:70000-0411-1	1 in 1 CARTON	12/10/2018	
2		24 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:70000-0411-3	1 in 1 CARTON	12/10/2018	
3		100 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	11/29/2018	

Labeler - Cardinal Health (063997360)