

DIMENHYDRINATE- dimenhydrinate tablet
Cardinal Health

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

1006-Cardinal

Drug Facts

Active ingredient (in each tablet)

Dimenhydrinate 50 mg

Purpose

Antiemetic

Uses

for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness

Do not give to children under 2 years of age unless directed by a doctor

Ask a doctor before use if you have

- glaucoma
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

■ to prevent motion sickness, the first dose should be taken one-half to one hour before starting activity

■ **adults and children 12 years of age and over:** 1 to 2 tablets every 4-6 hours; not to exceed 8 tablets in 24 hours, or as directed by a doctor

■ **children 6 to under 12 years of age:** 1/2 to 1 tablet every 6-8 hours; not to exceed 3 tablets in 24 hours, or as directed by a doctor

■ **children 2 to under 6 years of age:** 1/2 tablet every 6-8 hours; not to exceed 1 1/2 tablets in 24 hours, or as directed by a doctor

Other information

■ **each tablet contains:** calcium 30 mg

■ store in a dry place at 15° – 30°C (59° – 86°F)

■ protect from moisture

■ see end flap for expiration date and lot number

■ **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**

croscarmellose sodium, dicalcium phosphate, magnesium stearate, microcrystalline cellulose

Questions or comments?

1-800-231-4670

*This product is not manufactured or distributed by Medtech Products Inc., owner of the registered trademark Dramamine®

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

©2021 Cardinal Health. All Rights Reserved.

CARDINAL HEALTH, the Cardinal Health LOGO, LEADER, and the Leader LOGO are trademarks or registered trademarks of Cardinal Health. All other marks are the property of their respective owners

DISTRIBUTED BY CARDINAL HEALTH

DUBLIN, OHIO 43017

www.myleader.com 1-800-200-6313

Essential to Care™ since 1979

All LEADER™ Brand

Products Have A

100%

Money Back

Guarantee

LEADER

NDC 7000-0404-1

Original Formula

Motion Sickness Relief

Dimenhydrinate, 50 mg/ Antiemetic

COMPARE TO DRAMAMINE® ORIGINAL FORMULA active ingredient*

Fast Acting Motion Sickness Relief for Children & Adults

12 Tablets

R50592



NDC 70000-0404-1

Original Formula

Motion Sickness Relief

Dimenhydrinate, 50 mg | Antiemetic



Return to place of purchase if not satisfied.

Fast Acting Motion Sickness Relief for Children & Adults



Actual Size

12 TABLETS

COMPARE TO DRAMAMINE® ORIGINAL FORMULA active ingredient*

100% Money Back Guarantee

©2021 Cardinal Health. All Rights Reserved. CARDINAL HEALTH, the Cardinal Health LOGO, ESSENTIAL TO CARE, LEADER, and the Leader LOGO are trademarks or registered trademarks of Cardinal Health. All other marks are the property of their respective owners.

Cardinal Health
DISTRIBUTED BY CARDINAL HEALTH
DUBLIN, OHIO 43017
www.myleader.com 1-800-200-6575
Essential to Care™ since 1979

Drug Facts (continued)

Inactive ingredients croscarmellose sodium, dicalcium phosphate, magnesium stearate, microcrystalline cellulose

Questions or comments? 1-800-231-4670

CIN 5455548 REV. 8/21



0 96295 13518 3

When using this product

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland

Other information

- children 6 to under 12 years of age: 1/2 to 1 tablet every 6-8 hours; not to exceed 3 tablets in 24 hours, or as directed by a doctor
- children 2 to under 6 years of age: 1/2 tablet every 6-8 hours; not to exceed 1 1/2 tablets in 24 hours, or as directed by a doctor

IS OPENED OR BLISTER IS TORN OR BROKEN

- TAMPER EVIDENT; DO NOT USE IF OUTER PACKAGE
- see end flap for expiration date and lot number
- each tablet contains: calcium 30 mg ■ store in a dry place at 15° - 30°C (59° - 86°F) ■ protect from moisture



DIMENHYDRINATE

dimenhydrinate tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70000-0404
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIMENHYDRINATE (UNII: JB937PER5C) (CHLORTHEOPHYLLINE - UNII:GE2UA340FM)	DIMENHYDRINATE	50 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	9mm
Flavor		Imprint Code	1006;1006
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70000-0404-1	1 in 1 CARTON	01/04/2019	

1	12 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part336	01/04/2019	

Labeler - Cardinal Health (063997360)

Revised: 11/2021

Cardinal Health