

**CIRCLE K NIGHTTIME COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, and doxylamine succinate capsule, liquid filled  
Lil' Drug Store Products, Inc.**

-----  
**Circle K® Nighttime Cold & Flu**

***Drug Facts***

---

<b><i>Active ingredients (in each softgel)</i></b>	<b><i>Purposes</i></b>
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 15 mg	Cough suppressant
Doxylamine succinate 6.25 mg	Antihistamine

---

**Uses**

- temporarily relieves common cold and flu symptoms:
  - sore throat
  - headache
  - minor aches and pains
  - fever
  - runny nose and sneezing
  - cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

**Allergy alert**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

**When using this product**

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning**

Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- **when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing**

### **Other information**

- store between 15-30°C (59-86°F)
- avoid excessive heat

### **Inactive ingredients**

D&C yellow #10, FD&C blue #1, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

### **Questions or comments?**

call toll-free **1-877-507-6516**  
(M-F 8AM-4:30PM CST)

**Proudly distributed by Circle K Stores Inc.**

### **PRINCIPAL DISPLAY PANEL - 16 Softgel Blister Pack Carton**

Compare to the Active Ingredients in  
Vicks® NyQuil® Cold & Flu LiquiCaps®\*\*

CIRCLE K®

Nighttime Cold & Flu

Acetaminophen, Dextromethorphan HBr, and Doxylamine Succinate  
Pain Reliever/Fever Reducer, Cough Suppressant, Antihistamine

Relieves

- Aches - Fever - Sneezing
- Cough - Runny Nose
- Sore Throat

ACTUAL SIZE

16 Softgels  
(Liquid-filled capsules)

**CIRCLE K**

# Nighttime Cold & Flu

**Acetaminophen**, Dextromethorphan HBr, and Doxylamine Succinate  
Pain Reliever/Fever Reducer, Cough Suppressant, Antihistamine

Compare to the Active Ingredients in  
Vicks® NyQuil® Cold & Flu LiquiCaps®\*\*

**CIRCLE K**

# Nighttime Cold & Flu

**Acetaminophen**, Dextromethorphan HBr, and Doxylamine Succinate  
Pain Reliever/Fever Reducer, Cough Suppressant, Antihistamine



ACTUAL SIZE

## Relieves

- Aches - Fever - Sneezing
- Cough - Runny Nose
- Sore Throat

**16** Softgels  
(Liquid-filled capsules)

Product manufactured for:  
Lil' Drug Store Products, Inc.  
9300 Earhart Lane SW  
Cedar Rapids, IA 52404



**16** Softgels  
(Liquid-filled capsules)

**CIRCLE K**

# Nighttime Cold & Flu

**Acetaminophen**, Dextromethorphan HBr, and Doxylamine Succinate  
Pain Reliever/Fever Reducer, Cough Suppressant, Antihistamine

Proudly distributed by Circle K Stores Inc.

PLD-A585B FC006332



Lot No:

Exp Date:

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.**  
**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

**Drug Facts**

**Active ingredients (in each softgel)**      **Purposes**

Acetaminophen 325 mg.....Pain reliever/fever reducer  
Dextromethorphan HBr 15 mg.....Cough suppressant  
Doxylamine succinate 6.25 mg.....Antihistamine

**Uses**

- temporarily relieves common cold and flu symptoms:
• sore throat • headache • minor aches and pains
• fever • runny nose and sneezing
• cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:
• more than 4,000 mg of acetaminophen in 24 hours
• with other drugs containing acetaminophen
• 3 or more alcoholic drinks daily while using this product
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening
• blisters • rash If a skin reaction occurs, stop use and seek medical help right away.
**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains

**Drug Facts (continued)**

an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease • glaucoma
• trouble urinating due to an enlarged prostate gland
• a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis or emphysema
• cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
• taking sedatives or tranquilizers

When using this product

- do not exceed recommended dosage
• excitability may occur, especially in children
• avoid alcoholic drinks • marked drowsiness may occur
• alcohol, sedatives, and tranquilizers may increase drowsiness
• be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
• fever gets worse or lasts more than 3 days
• redness or swelling is present • new symptoms occur
• cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Drug Facts (continued)**

**Directions**

- do not take more than directed (see Overdose warning)
• do not take more than 4 doses in 24 hours
• adults and children 12 years and over: take 2 softgels with water every 6 hours
• swallow whole; do not crush, chew, or dissolve
• children under 12 years: do not use
• when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

**Other information**

- store between 15-30°C (59-86°F)
• avoid excessive heat

**Inactive ingredients**

D&C yellow #10, FD&C blue #1, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

**Questions or comments?**

call toll-free 1-877-507-6516 (M-F 8AM-4:30PM CST)

\*\*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, NyQuil® and LiquiCaps® are registered trademarks of The Procter & Gamble Company.


**PARENTS:**

Learn about teen medicine abuse

www.StopMedicineAbuse.org

PRINT SIDE GLUE

57728C-US-10-19-PU



**PDP**

**RELIEVES ACHES, FEVER, SNEEZING, COUGH, RUNNY NOSE**

Compare to the active ingredients in

**Vicks® NyQuil® Cold & Flu LiquiCaps®\*\***

**CIRCLE K**

**nighttime**

**cold & flu**

**Acetaminophen 325 mg,  
Dextromethorphan HBr 15 mg, Doxylamine Succinate 6.25 mg  
Pain Reliever/Fever Reducer, Cough Suppressant,  
Antihistamine**

[softgel image]

actual size

**16 softgels\***

<p><b>TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.</b></p> <p><b>KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.</b></p>		<p><b>Drug Facts (continued)</b></p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>do not take more than directed (see Overdose warning)</li> <li>do not take more than 4 doses in 24 hours</li> <li>adults and children 12 years and over: take 2 softgels with water every 6 hours</li> <li>swallow whole; do not crush, chew, or dissolve</li> <li>children under 12 years: do not use</li> <li>when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>store between 15°-30° (59°-86°)</li> <li>avoid excessive heat</li> </ul> <p><b>Inactive ingredients</b></p> <p>D&amp;C yellow #10, FD&amp;C blue #1, gelatin, glycerin, lactin, light mineral oil, mannitol, polyethylene glycol, polydioxane, propylene glycol, purified water, sorbitan, sorbitol, white ink</p> <p><b>Questions or comments?</b> call toll-free 1-877-567-6516 (M-F 9AM-4:30PM CST)</p> <p><small>* (Liquid-filled capsules)</small></p> <p><small>**This product is not manufactured or distributed by The Procter &amp; Gamble Company, distributor of Vicks® NyQuil® Cold &amp; Flu LiquiCaps®.</small></p> <p><b>PARENTS:</b> Learn about teen medicine abuse <a href="http://www.StopMedicineAbuse.org">www.StopMedicineAbuse.org</a></p>							
<p><b>Drug Facts</b></p> <table border="1"> <thead> <tr> <th>Active ingredients (in each softgel)</th> <th>Purposes</th> </tr> </thead> <tbody> <tr> <td>Acetaminophen 325 mg.....</td> <td>Pain reliever/fever reducer</td> </tr> <tr> <td>Dextromethorphan HBr 15 mg.....</td> <td>Cough suppressant</td> </tr> <tr> <td>Doxylamine succinate 6.25 mg.....</td> <td>Antihistamine</td> </tr> </tbody> </table> <p><b>Uses</b></p> <ul style="list-style-type: none"> <li>temporarily relieves common cold and flu symptoms:</li> <li>sore throat • headache • minor aches and pains</li> <li>fever • runny nose and sneezing</li> <li>cough due to minor throat and bronchial irritation</li> </ul> <p><b>Warnings</b></p> <p><b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if you take:</p> <ul style="list-style-type: none"> <li>more than 4,000 mg of acetaminophen in 24 hours</li> <li>with other drugs containing acetaminophen</li> <li>3 or more alcoholic drinks daily while using this product</li> </ul> <p><b>Allergy alert:</b> Acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> <li>skin redness</li> <li>hives</li> <li>rash</li> </ul> <p>If a skin reaction occurs, stop use and seek medical help right away.</p> <p><b>Sore throat warning:</b> If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.</li> <li>if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains</li> </ul>	Active ingredients (in each softgel)		Purposes	Acetaminophen 325 mg.....	Pain reliever/fever reducer	Dextromethorphan HBr 15 mg.....	Cough suppressant	Doxylamine succinate 6.25 mg.....	Antihistamine
Active ingredients (in each softgel)	Purposes								
Acetaminophen 325 mg.....	Pain reliever/fever reducer								
Dextromethorphan HBr 15 mg.....	Cough suppressant								
Doxylamine succinate 6.25 mg.....	Antihistamine								

RELIEVES ACHES, FEVER, SNEEZING, COUGH, RUNNY NOSE

Compare to the active ingredients in

**Vicks® NyQuil® Cold & Flu LiquiCaps®\*\***



**16 softgels\***



actual size

**16**

nighttime

**cold & flu**

softgels

**CIRCLE K**

**CIRCLE K NIGHTTIME COLD AND FLU**

acetaminophen, dextromethorphan hydrobromide, and doxylamine succinate capsule, liquid filled

**Product Information**

**Product Type**

HUMAN OTC DRUG

**Item Code (Source)**

NDC:66715-5572

<b>Route of Administration</b>	ORAL
--------------------------------	------

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN, UNSPECIFIED</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	
<b>LIGHT MINERAL OIL</b> (UNII: N6K5787QVP)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	

### Product Characteristics

<b>Color</b>	green	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	PC10
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-5572-8	8 in 1 CARTON	06/16/2021	
1		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/16/2021	

# CIRCLE K NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, and doxylamine succinate capsule, liquid filled

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:66715-5772
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN, UNSPECIFIED</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	
<b>LIGHT MINERAL OIL</b> (UNII: N6K5787QVP)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	

## Product Characteristics

<b>Color</b>	green	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	PC10
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-5772-8	8 in 1 CARTON	03/01/2020	08/10/2024
-		2 in 1 BLISTER PACK; Type 0: Not a Combination		



Product			
<b>Marketing Information</b>			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/05/2014	08/10/2024

**Labeler** - Lil' Drug Store Products, Inc. (093103646)

Revised: 12/2023

Lil' Drug Store Products, Inc.