

LBEL EFFET PARFAIT SIGNS OF AGE APPEARANCE MINIMIZER FOUNDATION SPF 20 FM CLAIRE 1 - BEIGE- octinoxate and titanium dioxide lotion
LBEL EFFET PARFAIT SIGNS OF AGE APPEARANCE MINIMIZER FOUNDATION SPF 20 FM CLAIRE 2 - BEIGE- octinoxate and titanium dioxide lotion
LBEL EFFET PARFAIT SIGNS OF AGE APPEARANCE MINIMIZER FOUNDATION SPF 20 FM CLAIRE 3 - BEIGE- octinoxate and titanium dioxide lotion
LBEL EFFET PARFAIT SIGNS OF AGE APPEARANCE MINIMIZER FOUNDATION SPF 20 FM CLAIRE 4 - BEIGE- octinoxate and titanium dioxide lotion
LBEL EFFET PARFAIT SIGNS OF AGE APPEARANCE MINIMIZER FOUNDATION SPF 20 FM MEDIUM 5 - BEIGE- octinoxate and titanium dioxide lotion
LBEL EFFET PARFAIT SIGNS OF AGE APPEARANCE MINIMIZER FOUNDATION SPF 20 FM MEDIUM 6 - BEIGE- octinoxate and titanium dioxide lotion
LBEL EFFET PARFAIT SIGNS OF AGE APPEARANCE MINIMIZER FOUNDATION SPF 20 FM MEDIUM 7 - BEIGE- octinoxate and titanium dioxide lotion
LBEL EFFET PARFAIT SIGNS OF AGE APPEARANCE MINIMIZER FOUNDATION SPF 20 FM OBSCURE 8 - BROWN- octinoxate and titanium dioxide lotion
LBEL EFFET PARFAIT SIGNS OF AGE APPEARANCE MINIMIZER FOUNDATION SPF 20 FM OBSCURE 9 - BROWN- octinoxate and titanium dioxide lotion
Ventura International LTD

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

LBEL EFFET PARFAIT signs of age appearance minimizer foundation SPF 20 FM

Drug Facts

| <i>Active Ingredients</i> | <i>Purpose</i> |
|----------------------------------|-----------------------|
| OCTINOXATE 7.00%, | Sunscreen |
| TITANIUM DIOXIDE 2.44% | Sunscreen |

Uses

- Helps prevent sunburn

Warnings

- Skin Cancer / Skin Aging Alert; Spending time in the sun increases your risk of skin cancer and early skin aging. This product has been shown only to help prevent sunburn, not skin cancer or early skin aging.
- For external use only.
- Do not use on damage or broken skin.
- When using this product keep out of eyes. Rinse with water to remove.
- Stop use and ask a doctor if rash occurs.
- Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Apply liberally and evenly 15 minutes before sun exposure.
- Children under 6 months of age: Ask a doctor.
- Reapply at least every 2 hours.
- Use a water resistant sunscreen if swimming or sweating.

Other information

- Protect the product in this container from excessive heat and direct sun.

Inactive ingredients

WATER, CYCLOPENTASILOXANE, POLYMETHYL METHACRYLATE, DIMETHICONE, CYCLOHEXASILOXANE, CETYL PEG/PPG-10/1 DIMETHICONE, POLYGLYCERYL-4 ISOSTEARATE, HEXYL LAURATE, TRIMETHYLSILOXYSILICATE, GLYCERIN, SODIUM CHLORIDE, PEG/PPG-19/19 DIMETHICONE, HYDROXYACETOPHENONE, PHENOXYETHANOL, COFFEA ARABICA (COFFEE) SEEDCAKE EXTRACT, PEG/PPG-18/18 DIMETHICONE, ALUMINA, BUTYLENE GLYCOL, OCTYLDODECANOL, CAPRYLYL GLYCOL, TOCOPHERYL ACETATE, CHLORPHENESIN, HYDROGEN DIMETHICONE, BHT, BENZYL ALCOHOL, POTASSIUM SORBATE, CARBOMER, POLYSORBATE 20, PALMITOYL TRIPEPTIDE-1, PALMITOYL TETRAPEPTIDE-7.

MAY CONTAIN:

TITANIUM DIOXIDE, IRON OXIDES, TRIETHOXYCAPRYLYLSILANE.

Dist. by Ventura Corp,
Ltd., San Juan, Puerto Rico 00926.

PRINCIPAL DISPLAY PANEL - 30 ml Bottle Box - CLAIRE 1 - BEIGE

L'BEL

EFFET

PARFAIT

signs of age

appearance

minimizer

foundation

SPF 20 FM

1 fl.oz. e (30 ml)

Drug Facts

Active Ingredients Purpose
 Octinoxate 7.00%..... sunscreen
 Titanium Dioxide 2.44%..... sunscreen

Uses
 ■ Helps prevent sunburn

Warnings
Skin Cancer / Skin Aging Alert:
 Spending time in the sun increases your risk of skin cancer and early skin aging. This product has been shown only to help prevent sunburn, **not** skin cancer or early skin aging.

For external use only.
Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions
 ■ Apply liberally and evenly 15 minutes before sun exposure.
 ■ Children under 6 months of age: Ask a doctor
 ■ Reapply at least every 2 hours
 ■ Use a water resistant sunscreen if swimming or sweating.

Other information
 ■ Protect the product in this container from excessive heat and direct sun.

Inactive ingredients:
 WATER, CYCLOPENTASILOXANE, POLYMETHYL METHACRYLATE, DIMETHICONE, CYCLOHEXASILOXANE, CETYL PEG/PPG-10/1 DIMETHICONE, POLYGLYCERYL-4 ISOSTEARATE, HEXYL LAURATE, TRIMETHYLSILOXYSILICATE, ▶

Drug Facts
 (continued)

GLYCERIN, SODIUM CHLORIDE, PEG/PPG-19/19 DIMETHICONE, HYDROXYACETOPHENONE, PHENOXYETHANOL, COFFEA ARABICA (COFFEE) SEEDCAKE EXTRACT, PEG/PPG-18/18 DIMETHICONE, ALUMINA, BUTYLENE GLYCOL, CAPRYLYL GLYCOL, TOCOPHERYL ACETATE, CHLORPHENESIN, HYDROGEN DIMETHICONE, BHT, BENZYL ALCOHOL, POTASSIUM SORBATE, CARBOMER, POLYSORBATE 20, PALMITOYL TRIPEPTIDE-1, PALMITOYL TETRAPEPTIDE-7.

MAY CONTAIN: TITANIUM DIOXIDE, IRON OXIDES, TRIETHOXYCAPRYLYLSILANE

U.S.A.: Dist. by Ventura Int Ltd. DBA Belcorp USA, Miami, FL 33126. Made in Colombia. PUERTO RICO: Dist. by Ventura Corp, Ltd., San Juan, Puerto Rico 00926. Made in Colombia.
 www.lbel.com

**EFFET
 PARFAIT**

signs of age
 appearance
 minimizer
foundation
 SPF 20 FM

1 fl.oz. e (30 ml)

PRINCIPAL DISPLAY PANEL - 30 ml Bottle Box - CLAIRE 2 - BEIGE

L'BEL
 EFFET
 PARFAIT
 signs of age
 appearance
 minimizer
 foundation

SPF 20 FM

1 fl.oz. e (30 ml)

Drug Facts

| Active Ingredients | Purpose |
|------------------------|-----------|
| Octinoxate 7.00% | sunscreen |
| Titanium Dioxide 2.44% | sunscreen |

Uses

- Helps prevent sunburn

Warnings

Skin Cancer / Skin Aging Alert: Spending time in the sun increases your risk of skin cancer and early skin aging. This product has been shown only to help prevent sunburn, **not** skin cancer or early skin aging.

For external use only.

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Apply liberally and evenly 15 minutes before sun exposure.
- Children under 6 months of age: Ask a doctor
- Reapply at least every 2 hours
- Use a water resistant sunscreen if swimming or sweating.

Other information

- Protect the product in this container from excessive heat and direct sun.

Inactive ingredients:
 WATER, CYCLOPENTASILOXANE, POLYMETHYL METHACRYLATE, DIMETHICONE, CYCLOHEXASILOXANE, CETYL PEG/PPG-10/1 DIMETHICONE, POLYGLYCERYL-4 ISOSTEARATE, HEXYL LAURATE, TRIMETHYLSILOXYSILICATE

Drug Facts
(continued)

GLYCERIN, SODIUM CHLORIDE, PEG/PPG-19/19 DIMETHICONE, HYDROXYACETOPHENONE, PHENOXYETHANOL, COFFEA ARABICA (COFFEE) SEEDCAKE EXTRACT, PEG/PPG-18/18 DIMETHICONE, ALUMINA, BUTYLENE GLYCOL, CAPRYLYL GLYCOL, TOCOPHERYL ACETATE, CHLORPHENESIN, HYDROGEN DIMETHICONE, BHT, BENZYL ALCOHOL, POTASSIUM SORBATE, CARBOMER, POLYSORBATE 20, PALMITOYL TRIPEPTIDE-1, PALMITOYL TETRAPEPTIDE-7.

MAY CONTAIN: TITANIUM DIOXIDE, IRON OXIDES, TRIETHOXYCAPRYLYLSILANE

U.S.A.: Dist. by Ventura Int Ltd. DBA Belcorp USA, Miami, FL 33126. Made in Colombia. PUERTO RICO: Dist. by Ventura Corp. Ltd., San Juan, Puerto Rico 00926. Made in Colombia.
www.lbel.com

EFFET
PARFAIT

signs of age
appearance
minimizer
foundation
SPF 20 FM

1 fl.oz. e (30 ml)

PRINCIPAL DISPLAY PANEL - 30 ml Bottle Box - CLAIRE 3 - BEIGE

L'BEL

EFFET
PARFAIT

signs of age

appearance
 minimizer
 foundation
 SPF 20 FM
 1 fl.oz. e (30 ml)

Drug Facts

| Active Ingredients | Purpose |
|------------------------|-----------------|
| Octinoxate 7.00% | sunscreen |
| Titanium Dioxide 2.44% | sunscreen |

Uses

- Helps prevent sunburn

Warnings

Skin Cancer / Skin Aging Alert:
 Spending time in the sun increases your risk of skin cancer and early skin aging. This product has been shown only to help prevent sunburn, **not** skin cancer or early skin aging.

For external use only.

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Apply liberally and evenly 15 minutes before sun exposure.
- Children under 6 months of age: Ask a doctor
- Reapply at least every 2 hours
- Use a water resistant sunscreen if swimming or sweating.

Other information

- Protect the product in this container from excessive heat and direct sun.

Inactive ingredients:
 WATER, CYCLOPENTASILOXANE, POLYMETHYL METHACRYLATE, DIMETHICONE, CYCLOHEXASILOXANE, CETYL PEG/PPG-10/1 DIMETHICONE, POLYGLYCERYL-4 ISOSTEARATE, HEXYL LAURATE, TRIMETHYLSILOXYSILICATE,

Drug Facts
 (continued)

GLYCERIN, SODIUM CHLORIDE, PEG/PPG-19/19 DIMETHICONE, HYDROXYACETOPHENONE, PHENOXYETHANOL, COFFEA ARABICA (COFFEE) SEEDCAKE EXTRACT, PEG/PPG-18/18 DIMETHICONE, ALUMINA, BUTYLENE GLYCOL, CAPRYLYL GLYCOL, TOCOPHERYL ACETATE, CHLORPHENESIN, HYDROGEN DIMETHICONE, BHT, BENZYL ALCOHOL, POTASSIUM SORBATE, CARBOMER, POLYSORBATE 20, PALMITOYL TRIPEPTIDE-1, PALMITOYL TETRAPEPTIDE-7.

MAY CONTAIN: TITANIUM DIOXIDE, IRON OXIDES, TRIETHOXYCAPRYLYLSILANE

U.S.A.: Dist. by Ventura Int Ltd. DBA Belcorp USA, Miami, FL 33126. Made in Colombia. PUERTO RICO: Dist. by Ventura Corp, Ltd., San Juan, Puerto Rico 00926. Made in Colombia.
 www.lbel.com

EFFET
 PARFAIT

signs of age
 appearance
 minimizer
foundation
 SPF 20 FM

1 fl.oz. e (30 ml)

PRINCIPAL DISPLAY PANEL - 30 ml Bottle Box - CLAIRE 4 - BEIGE

L'BEL
 EFFET

PARFAIT

signs of age
appearance
minimizer
foundation
SPF 20 FM

1 fl.oz. e (30 ml)

Drug Facts

| Active Ingredients | Purpose |
|---------------------------|-----------------|
| Octinoxate 7.00% | sunscreen |
| Titanium Dioxide 2.44% | sunscreen |

Uses

- Helps prevent sunburn

Warnings

Skin Cancer / Skin Aging Alert:
Spending time in the sun increases your risk of skin cancer and early skin aging. This product has been shown only to help prevent sunburn, **not** skin cancer or early skin aging.

For external use only.

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Apply liberally and evenly 15 minutes before sun exposure.
- Children under 6 months of age: Ask a doctor
- Reapply at least every 2 hours
- Use a water resistant sunscreen if swimming or sweating.

Other information

- Protect the product in this container from excessive heat and direct sun.

Inactive ingredients:

WATER, CYCLOPENTASILOXANE, POLYMETHYL METHACRYLATE, DIMETHICONE, CYCLOHEXASILOXANE, CETYL PEG/PPG-10/1 DIMETHICONE, POLYGLYCERYL-4 ISOSTEARATE, HEXYL LAURATE, TRIMETHYLSILOXYSILICATE, ▶

Drug Facts
(continued)

GLYCERIN, SODIUM CHLORIDE, PEG/PPG-19/19 DIMETHICONE, HYDROXYACETOPHENONE, PHENOXYETHANOL, COFFEA ARABICA (COFFEE) SEEDCAKE EXTRACT, PEG/PPG-18/18 DIMETHICONE, ALUMINA, BUTYLENE GLYCOL, CAPRYLYL GLYCOL, TOCOPHERYL ACETATE, CHLORPHENESIN, HYDROGEN DIMETHICONE, BHT, BENZYL ALCOHOL, POTASSIUM SORBATE, CARBOMER, POLYSORBATE 20, PALMITOYL TRIPEPTIDE-1, PALMITOYL TETRAPEPTIDE-7.

MAY CONTAIN: TITANIUM DIOXIDE, IRON OXIDES, TRIETHOXYCAPRYLYLSILANE

U.S.A.: Dist. by Ventura Int Ltd. DBA Belcorp USA, Miami, FL 33126. Made in Colombia. PUERTO RICO: Dist. by Ventura Corp, Ltd., San Juan, Puerto Rico 00926. Made in Colombia.

www.lbel.com

EFFET
PARFAIT

signs of age
appearance
minimizer
foundation
SPF 20 FM

1 fl.oz. e (30 ml)

L'BEL

EFFET
PARFAIT

signs of age
appearance
minimizer
foundation
SPF 20 FM

1 fl.oz. e (30 ml)

Drug Facts

| Active Ingredients | Purpose |
|------------------------|-----------------|
| Octinoxate 7.00% | sunscreen |
| Titanium Dioxide 2.44% | sunscreen |

Uses

- Helps prevent sunburn

Warnings

Skin Cancer / Skin Aging Alert:
Spending time in the sun increases your risk of skin cancer and early skin aging. This product has been shown only to help prevent sunburn, **not** skin cancer or early skin aging.

For external use only.

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Apply liberally and evenly 15 minutes before sun exposure.
- Children under 6 months of age: Ask a doctor
- Reapply at least every 2 hours
- Use a water resistant sunscreen if swimming or sweating.

Other information

- Protect the product in this container from excessive heat and direct sun.

Inactive ingredients:

WATER, CYCLOPENTASILOXANE, POLYMETHYL METHACRYLATE, DIMETHICONE, CYCLOHEXASILOXANE, CETYL PEG/PPG-10/1 DIMETHICONE, POLYGLYCERYL-4 ISOSTEARATE, HEXYL LAURATE, TRIMETHYLSILOXYSILICATE, ▶

Drug Facts

(continued)

GLYCERIN, SODIUM CHLORIDE, PEG/PPG-19/19 DIMETHICONE, HYDROXYACETOPHENONE, PHENOXYETHANOL, COFFEA ARABICA (COFFEE) SEEDCAKE EXTRACT, PEG/PPG-18/18 DIMETHICONE, ALUMINA, BUTYLENE GLYCOL, CAPRYLYL GLYCOL, TOCOPHERYL ACETATE, CHLORPHENESIN, HYDROGEN DIMETHICONE, BHT, BENZYL ALCOHOL, POTASSIUM SORBATE, CARBOMER, POLYSORBATE 20, PALMITOYL TRIPEPTIDE-1, PALMITOYL TETRAPEPTIDE-7.

MAY CONTAIN: TITANIUM DIOXIDE, IRON OXIDES, TRIETHOXYCAPRYLYLSILANE

U.S.A.: Dist. by Ventura Int Ltd. DBA Belcorp USA, Miami, FL 33126. Made in Colombia. PUERTO RICO: Dist. by Ventura Corp, Ltd., San Juan, Puerto Rico 00926. Made in Colombia.

www.lbel.com

EFFET
PARFAIT

signs of age
appearance
minimizer
foundation
SPF 20 FM

1 fl.oz. e (30 ml)

PRINCIPAL DISPLAY PANEL - 30 ml Bottle Box - MEDIUM 6 - BEIGE

L'BEL

EFFET

PARFAIT

signs of age

appearance

minimizer

foundation

SPF 20 FM

1 fl.oz. e (30 ml)

Drug Facts

Active Ingredients Purpose

Octinoxate 7.00%..... sunscreen
Titanium Dioxide 2.44%..... sunscreen

Uses

- Helps prevent sunburn

Warnings

Skin Cancer / Skin Aging Alert:
Spending time in the sun increases your risk of skin cancer and early skin aging. This product has been shown only to help prevent sunburn, **not** skin cancer or early skin aging.

For external use only.

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Apply liberally and evenly 15 minutes before sun exposure.
- Children under 6 months of age: Ask a doctor
- Reapply at least every 2 hours
- Use a water resistant sunscreen if swimming or sweating.

Other information

- Protect the product in this container from excessive heat and direct sun.

Inactive ingredients:

WATER, CYCLOPENTASILOXANE, POLYMETHYL METHACRYLATE, DIMETHICONE, CYCLOHEXASILOXANE, CETYL PEG/PPG-10/1 DIMETHICONE, POLYGLYCERYL-4 ISOSTEARATE, HEXYL LAURATE, TRIMETHYLSILOXYSILICATE, ▶

Drug Facts

(continued)

GLYCERIN, SODIUM CHLORIDE, PEG/PPG-19/19 DIMETHICONE, HYDROXYACETOPHENONE, PHENOXYETHANOL, COFFEA ARABICA (COFFEE) SEEDCAKE EXTRACT, PEG/PPG-18/18 DIMETHICONE, ALUMINA, BUTYLENE GLYCOL, CAPRYLYL GLYCOL, TOCOPHERYL ACETATE, CHLORPHENESIN, HYDROGEN DIMETHICONE, BHT, BENZYL ALCOHOL, POTASSIUM SORBATE, CARBOMER, POLYSORBATE 20, PALMITOYL TRIPEPTIDE-1, PALMITOYL TETRAPEPTIDE-7.

MAY CONTAIN: TITANIUM DIOXIDE, IRON OXIDES, TRIETHOXYCAPRYLYLSILANE

U.S.A.: Dist. by Ventura Int Ltd. DBA Belcorp USA, Miami, FL 33126. Made in Colombia. PUERTO RICO: Dist. by Ventura Corp, Ltd., San Juan, Puerto Rico 00926. Made in Colombia.

www.lbel.com

EFFET
PARFAIT

signs of age
appearance
minimizer
foundation
SPF 20 FM

1 fl.oz. e (30 ml)

PRINCIPAL DISPLAY PANEL - 30 ml Bottle Box - MEDIUM 7 - BEIGE

L'BEL

EFFET
PARFAIT

signs of age
appearance
minimizer
foundation

SPF 20 FM

1 fl.oz. e (30 ml)

Drug Facts

| Active Ingredients | Purpose |
|------------------------|-----------|
| Octinoxate 7.00% | sunscreen |
| Titanium Dioxide 2.44% | sunscreen |

Uses

- Helps prevent sunburn

Warnings

Skin Cancer / Skin Aging Alert: Spending time in the sun increases your risk of skin cancer and early skin aging. This product has been shown only to help prevent sunburn, **not** skin cancer or early skin aging.

For external use only.

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Apply liberally and evenly 15 minutes before sun exposure.
- Children under 6 months of age: Ask a doctor
- Reapply at least every 2 hours
- Use a water resistant sunscreen if swimming or sweating.

Other information

- Protect the product in this container from excessive heat and direct sun.

Inactive ingredients:
 WATER, CYCLOPENTASILOXANE, POLYMETHYL METHACRYLATE, DIMETHICONE, CYCLOHEXASILOXANE, CETYL PEG/PPG-10/1 DIMETHICONE, POLYGLYCERYL-4 ISOSTEARATE, HEXYL LAURATE, TRIMETHYLSILOXYSILICATE

Drug Facts
(continued)

GLYCERIN, SODIUM CHLORIDE, PEG/PPG-19/19 DIMETHICONE, HYDROXYACETOPHENONE, PHENOXYETHANOL, COFFEA ARABICA (COFFEE) SEEDCAKE EXTRACT, PEG/PPG-18/18 DIMETHICONE, ALUMINA, BUTYLENE GLYCOL, CAPRYLYL GLYCOL, TOCOPHERYL ACETATE, CHLORPHENESIN, HYDROGEN DIMETHICONE, BHT, BENZYL ALCOHOL, POTASSIUM SORBATE, CARBOMER, POLYSORBATE 20, PALMITOYL TRIPEPTIDE-1, PALMITOYL TETRAPEPTIDE-7.

MAY CONTAIN: TITANIUM DIOXIDE, IRON OXIDES, TRIETHOXYCAPRYLYLSILANE

U.S.A.: Dist. by Ventura Int Ltd. DBA Belcorp USA, Miami, FL 33126. Made in Colombia. PUERTO RICO: Dist. by Ventura Corp. Ltd., San Juan, Puerto Rico 00926. Made in Colombia.
www.lbel.com

EFFET
PARFAIT

signs of age
appearance
minimizer
foundation
SPF 20 FM

1 fl.oz. e (30 ml)

PRINCIPAL DISPLAY PANEL - 30 ml Bottle Box - OBSCURE 8 - BROWN

L'BEL

EFFET
PARFAIT

signs of age

appearance
 minimizer
 foundation
 SPF 20 FM
 1 fl.oz. e (30 ml)

Drug Facts

| Active Ingredients | Purpose |
|------------------------|-----------------|
| Octinoxate 7.00% | sunscreen |
| Titanium Dioxide 2.44% | sunscreen |

Uses

- Helps prevent sunburn

Warnings

Skin Cancer / Skin Aging Alert:
 Spending time in the sun increases your risk of skin cancer and early skin aging. This product has been shown only to help prevent sunburn, **not** skin cancer or early skin aging.

For external use only.

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Apply liberally and evenly 15 minutes before sun exposure.
- Children under 6 months of age: Ask a doctor
- Reapply at least every 2 hours
- Use a water resistant sunscreen if swimming or sweating.

Other information

- Protect the product in this container from excessive heat and direct sun.

Inactive ingredients:

WATER, CYCLOPENTASILOXANE, POLYMETHYL METHACRYLATE, DIMETHICONE, CYCLOHEXASILOXANE, CETYL PEG/PPG-10/1 DIMETHICONE, POLYGLYCERYL-4 ISOSTEARATE, HEXYL LAURATE, TRIMETHYLSILOXYSILICATE,

Drug Facts
(continued)

GLYCERIN, SODIUM CHLORIDE, PEG/PPG-19/19 DIMETHICONE, HYDROXYACETOPHENONE, PHENOXYETHANOL, COFFEA ARABICA (COFFEE) SEEDCAKE EXTRACT, PEG/PPG-18/18 DIMETHICONE, ALUMINA, BUTYLENE GLYCOL, CAPRYLYL GLYCOL, TOCOPHERYL ACETATE, CHLORPHENESIN, HYDROGEN DIMETHICONE, BHT, BENZYL ALCOHOL, POTASSIUM SORBATE, CARBOMER, POLYSORBATE 20, PALMITOYL TRIPEPTIDE-1, PALMITOYL TETRAPEPTIDE-7.

MAY CONTAIN: TITANIUM DIOXIDE, IRON OXIDES, TRIETHOXYCAPRYLYLSILANE

U.S.A.: Dist. by Ventura Int Ltd. DBA Belcorp USA, Miami, FL 33126. Made in Colombia. PUERTO RICO: Dist. by Ventura Corp, Ltd., San Juan, Puerto Rico 00926. Made in Colombia.
 www.lbel.com

EFFET
 PARFAIT

signs of age
 appearance
 minimizer
foundation
 SPF 20 FM

1 fl.oz. e (30 ml)

PRINCIPAL DISPLAY PANEL - 30 ml Bottle Box - OBSCURE 9 - BROWN

L'BEL
 EFFET

PARFAIT

signs of age
appearance
minimizer
foundation
SPF 20 FM

1 fl.oz. e (30 ml)

Drug Facts

| Active Ingredients | Purpose |
|---------------------------|-----------------|
| Octinoxate 7.00% | sunscreen |
| Titanium Dioxide 2.44% | sunscreen |

Uses

- Helps prevent sunburn

Warnings

Skin Cancer / Skin Aging Alert:
Spending time in the sun increases your risk of skin cancer and early skin aging. This product has been shown only to help prevent sunburn, **not** skin cancer or early skin aging.

For external use only.

Do not use on damaged or broken skin.

When using this product keep out of eyes. Rinse with water to remove.

Stop use and ask a doctor if rash occurs.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Apply liberally and evenly 15 minutes before sun exposure.
- Children under 6 months of age: Ask a doctor
- Reapply at least every 2 hours
- Use a water resistant sunscreen if swimming or sweating.

Other information

- Protect the product in this container from excessive heat and direct sun.

Inactive ingredients:

WATER, CYCLOPENTASILOXANE, POLYMETHYL METHACRYLATE, DIMETHICONE, CYCLOHEXASILOXANE, CETYL PEG/PPG-10/1 DIMETHICONE, POLYGLYCERYL-4 ISOSTEARATE, HEXYL LAURATE, TRIMETHYLSILOXYSILICATE, ▶

Drug Facts
(continued)

GLYCERIN, SODIUM CHLORIDE, PEG/PPG-19/19 DIMETHICONE, HYDROXYACETOPHENONE, PHENOXYETHANOL, COFFEA ARABICA (COFFEE) SEEDCAKE EXTRACT, PEG/PPG-18/18 DIMETHICONE, ALUMINA, BUTYLENE GLYCOL, CAPRYLYL GLYCOL, TOCOPHERYL ACETATE, CHLORPHENESIN, HYDROGEN DIMETHICONE, BHT, BENZYL ALCOHOL, POTASSIUM SORBATE, CARBOMER, POLYSORBATE 20, PALMITOYL TRIPEPTIDE-1, PALMITOYL TETRAPEPTIDE-7.

MAY CONTAIN: TITANIUM DIOXIDE, IRON OXIDES, TRIETHOXYCAPRYLYLSILANE

U.S.A.: Dist. by Ventura Int Ltd. DBA Belcorp USA, Miami, FL 33126. Made in Colombia. PUERTO RICO: Dist. by Ventura Corp, Ltd., San Juan, Puerto Rico 00926. Made in Colombia.

www.lbel.com

EFFET
PARFAIT

signs of age
appearance
minimizer
foundation
SPF 20 FM

1 fl.oz. e (30 ml)

FOUNDATION SPF 20 FM CLAIRE 1 - BEIGE

octinoxate and titanium dioxide lotion

Product Information

| | | | |
|-------------------------|----------------|--------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:14783-916 |
| Route of Administration | TOPICAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--------------------------------------------------------------------------|-------------------|------------------|
| Octinoxate (UNII: 4Y5P7MUD51) (Octinoxate - UNII:4Y5P7MUD51) | Octinoxate | 0.07 g in 1 mL |
| Titanium Dioxide (UNII: 15FIX9V2JP) (Titanium Dioxide - UNII:15FIX9V2JP) | Titanium Dioxide | 0.0244 g in 1 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|--------------------------------------------------|----------|
| WATER (UNII: 059QF0KO0R) | |
| CYCLOMETHICONE 5 (UNII: 0THT5PCI0R) | |
| DIMETHICONE (UNII: 92RU3N3Y1O) | |
| CYCLOMETHICONE 6 (UNII: XHK3U310BA) | |
| POLYGLYCERYL-4 ISO STEARATE (UNII: 820DPX33S7) | |
| HEXYL LAURATE (UNII: 4CG9F9W01Q) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| SODIUM CHLORIDE (UNII: 451W47IQ8X) | |
| PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL) | |
| PHENOXYETHANOL (UNII: HIE492ZZ3T) | |
| COFFEE BEAN (UNII: JFH385Y744) | |
| PEG/PPG-18/18 DIMETHICONE (UNII: 9H0AO7T794) | |
| ALUMINUM OXIDE (UNII: LM26O6933) | |
| BUTYLENE GLYCOL (UNII: 3XUS85K0RA) | |
| OCTYLDODECANOL (UNII: 461N1O614Y) | |
| CAPRYLYL GLYCOL (UNII: 00YIU5438U) | |
| .ALPHA.-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0) | |
| CHLORPHENESIN (UNII: I670DAL4SZ) | |
| BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K) | |
| BENZYL ALCOHOL (UNII: LKG8494WBH) | |
| POTASSIUM SORBATE (UNII: 1VPU26JZZ4) | |
| CARBOXYPOLYMETHYLENE (UNII: 0A5MM307FC) | |
| POLYSORBATE 20 (UNII: 7T1F30V5YH) | |
| PALMITOYL TRIPEPTIDE-1 (UNII: RV743D216M) | |
| PALMITOYL TETRAPEPTIDE-7 (UNII: Q41S464P1R) | |
| FERRIC OXIDE RED (UNII: 1K09F3G675) | |
| TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E) | |
| HYDROGEN DIMETHICONE (13 CST) (UNII: 4QGR4P2YOI) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|-----------|---------------------|----------------------|--------------------|
|---|-----------|---------------------|----------------------|--------------------|

| | | | |
|---|------------------|------------------------------------------------------|------------|
| 1 | NDC:14783-916-02 | 1 in 1 BOX | 12/02/2019 |
| 1 | NDC:14783-916-01 | 30 mL in 1 BOTTLE; Type 0: Not a Combination Product | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|------------------------------------------|----------------------|--------------------|
| OTC MONOGRAPH FINAL | part352 | 12/02/2019 | |

LBEL EFFET PARFAIT SIGNS OF AGE APPEARANCE MINIMIZER FOUNDATION SPF 20 FM CLAIRE 2 - BEIGE

octinoxate and titanium dioxide lotion

Product Information

| | | | |
|-------------------------|----------------|--------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:14783-917 |
| Route of Administration | TOPICAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--------------------------------------------------------------------------|-------------------|------------------|
| Octinoxate (UNII: 4Y5P7MUD51) (Octinoxate - UNII:4Y5P7MUD51) | Octinoxate | 0.07 g in 1 mL |
| Titanium Dioxide (UNII: 15FIX9V2JP) (Titanium Dioxide - UNII:15FIX9V2JP) | Titanium Dioxide | 0.0244 g in 1 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|-----------------------------------------------|----------|
| WATER (UNII: 059QF0K00R) | |
| CYCLOMETHICONE 5 (UNII: 0THF5PC10R) | |
| DIMETHICONE (UNII: 92RU3N3Y1O) | |
| CYCLOMETHICONE 6 (UNII: XHK3U310BA) | |
| POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7) | |
| HEXYL LAURATE (UNII: 4CG9F9W01Q) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| SODIUM CHLORIDE (UNII: 451W47IQ8X) | |
| PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL) | |
| PHENOXYETHANOL (UNII: HIE492ZZ3T) | |
| COFFEE BEAN (UNII: JFH385Y744) | |
| PEG/PPG-18/18 DIMETHICONE (UNII: 9H0AO7T794) | |
| ALUMINUM OXIDE (UNII: LM26O6933) | |
| BUTYLENE GLYCOL (UNII: 3XUS85K0RA) | |
| OCTYLDODECANOL (UNII: 461N1O614Y) | |
| CAPRYLYL GLYCOL (UNII: 00YIU5438U) | |
| .ALPHA.-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0) | |
| CHLORPHENESIN (UNII: I670DAL4SZ) | |
| BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K) | |
| BENZYL ALCOHOL (UNII: LKG8494WBH) | |
| POTASSIUM SORBATE (UNII: 1VPU26JZZ4) | |

| | |
|--------------------------------------------------|--|
| CARBOXYPOLYMETHYLENE (UNII: 0A5MM307FC) | |
| POLYSORBATE 20 (UNII: 7T1F30V5YH) | |
| PALMITOYL TRIPEPTIDE-1 (UNII: RV743D216M) | |
| PALMITOYL TETRAPEPTIDE-7 (UNII: Q41S464P1R) | |
| FERRIC OXIDE RED (UNII: 1K09F3G675) | |
| TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E) | |
| HYDROGEN DIMETHICONE (13 CST) (UNII: 4QGR4P2YOI) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|------------------------------------------------------|----------------------|--------------------|
| 1 | NDC:14783-917-02 | 1 in 1 BOX | 12/02/2019 | |
| 1 | NDC:14783-917-01 | 30 mL in 1 BOTTLE; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|------------------------------------------|----------------------|--------------------|
| OTC MONOGRAPH FINAL | part352 | 12/02/2019 | |

LBEL EFFET PARFAIT SIGNS OF AGE APPEARANCE MINIMIZER FOUNDATION SPF 20 FM CLAIRE 3 - BEIGE

octinoxate and titanium dioxide lotion

Product Information

| | | | |
|-------------------------|----------------|--------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:14783-918 |
| Route of Administration | TOPICAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--------------------------------------------------------------------------|-------------------|------------------|
| Octinoxate (UNII: 4Y5P7MUD51) (Octinoxate - UNII:4Y5P7MUD51) | Octinoxate | 0.07 g in 1 mL |
| Titanium Dioxide (UNII: 15FIX9V2JP) (Titanium Dioxide - UNII:15FIX9V2JP) | Titanium Dioxide | 0.0244 g in 1 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|-----------------------------------------------|----------|
| WATER (UNII: 059QF0K00R) | |
| CYCLOMETHICONE 5 (UNII: 0THT5PC10R) | |
| DIMETHICONE (UNII: 92RU3N3Y10) | |
| CYCLOMETHICONE 6 (UNII: XHK3U310BA) | |
| POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7) | |
| HEXYL LAURATE (UNII: 4CG9F9W01Q) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| SODIUM CHLORIDE (UNII: 451W47IQ8X) | |
| PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL) | |
| PHENOXYETHANOL (UNII: HIE49ZZZ3T) | |

| |
|--------------------------------------------------|
| COFFEE BEAN (UNII: JFH385Y744) |
| PEG/PPG-18/18 DIMETHICONE (UNII: 9H0AO7T794) |
| ALUMINUM OXIDE (UNII: LM26O6933) |
| BUTYLENE GLYCOL (UNII: 3XUS85K0RA) |
| OCTYLDODECANOL (UNII: 461N1O614Y) |
| CAPRYLYL GLYCOL (UNII: 00YIU5438U) |
| .ALPHA.-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0) |
| CHLORPHENESIN (UNII: I670DAL4SZ) |
| BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K) |
| BENZYL ALCOHOL (UNII: LKG8494WBH) |
| POTASSIUM SORBATE (UNII: 1VPU26JZZ4) |
| CARBOXYPOLYMETHYLENE (UNII: 0A5MM307FC) |
| POLYSORBATE 20 (UNII: 7T1F30V5YH) |
| PALMITOYL TRIPEPTIDE-1 (UNII: RV743D216M) |
| PALMITOYL TETRAPEPTIDE-7 (UNII: Q41S464P1R) |
| FERRIC OXIDE RED (UNII: 1K09F3G675) |
| TRIETHOXYCAPRYLSILANE (UNII: LDC331P08E) |
| HYDROGEN DIMETHICONE (13 CST) (UNII: 4QGR4P2YO1) |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|------------------------------------------------------|----------------------|--------------------|
| 1 | NDC:14783-918-02 | 1 in 1 BOX | 12/02/2019 | |
| 1 | NDC:14783-918-01 | 30 mL in 1 BOTTLE; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|------------------------------------------|----------------------|--------------------|
| OTC MONOGRAPH FINAL | part352 | 12/02/2019 | |

LBEL EFFET PARFAIT SIGNS OF AGE APPEARANCE MINIMIZER FOUNDATION SPF 20 FM CLAIRE 4 - BEIGE

octinoxate and titanium dioxide lotion

Product Information

| | | | |
|-------------------------|----------------|--------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:14783-919 |
| Route of Administration | TOPICAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--------------------------------------------------------------------------|-------------------|------------------|
| Octinoxate (UNII: 4Y5P7MUD51) (Octinoxate - UNII:4Y5P7MUD51) | Octinoxate | 0.07 g in 1 mL |
| Titanium Dioxide (UNII: 15FIX9V2JP) (Titanium Dioxide - UNII:15FIX9V2JP) | Titanium Dioxide | 0.0244 g in 1 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|--------------------------------------------------|----------|
| WATER (UNII: 059QF0KO0R) | |
| CYCLOMETHICONE 5 (UNII: 0THT5PC10R) | |
| DIMETHICONE (UNII: 92RU3N3Y1O) | |
| CYCLOMETHICONE 6 (UNII: XHK3U310BA) | |
| POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7) | |
| HEXYL LAURATE (UNII: 4CG9F9W01Q) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| SODIUM CHLORIDE (UNII: 451W47IQ8X) | |
| PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL) | |
| PHENOXYETHANOL (UNII: HIE492ZZ3T) | |
| COFFEE BEAN (UNII: JFH385Y744) | |
| PEG/PPG-18/18 DIMETHICONE (UNII: 9H0AO7T794) | |
| ALUMINUM OXIDE (UNII: LM26O6933) | |
| BUTYLENE GLYCOL (UNII: 3XUS85K0RA) | |
| OCTYLDODECANOL (UNII: 461N1O614Y) | |
| CAPRYLYL GLYCOL (UNII: 00YIU5438U) | |
| .ALPHA.-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0) | |
| CHLORPHENESIN (UNII: I670DAL4SZ) | |
| BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K) | |
| BENZYL ALCOHOL (UNII: LKG8494WBH) | |
| POTASSIUM SORBATE (UNII: 1VPU26JZZ4) | |
| CARBOXYPOLYMETHYLENE (UNII: 0A5MM307FC) | |
| POLYSORBATE 20 (UNII: 7T1F30V5YH) | |
| PALMITOYL TRIPEPTIDE-1 (UNII: RV743D216M) | |
| PALMITOYL TETRAPEPTIDE-7 (UNII: Q41S464P1R) | |
| FERRIC OXIDE RED (UNII: 1K09F3G675) | |
| TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E) | |
| HYDROGEN DIMETHICONE (13 CST) (UNII: 4QGR4P2YO1) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|------------------------------------------------------|----------------------|--------------------|
| 1 | NDC:14783-919-02 | 1 in 1 BOX | 12/02/2019 | |
| 1 | NDC:14783-919-01 | 30 mL in 1 BOTTLE; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|------------------------------------------|----------------------|--------------------|
| OTC MONOGRAPH FINAL | part352 | 12/02/2019 | |

LBEL EFFET PARFAIT SIGNS OF AGE APPEARANCE MINIMIZER FOUNDATION SPF 20 FM MEDIUM 5 - BEIGE

octinoxate and titanium dioxide lotion

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:14783-920 |
| Route of Administration | TOPICAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---------------------------------------------------------------------------------|-------------------|------------------|
| Octinoxate (UNII: 4Y5P7MUD51) (Octinoxate - UNII:4Y5P7MUD51) | Octinoxate | 0.07 g in 1 mL |
| Titanium Dioxide (UNII: 15FIX9V2JP) (Titanium Dioxide - UNII:15FIX9V2JP) | Titanium Dioxide | 0.0244 g in 1 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|---------------------------------------------------------|----------|
| WATER (UNII: 059QF0KO0R) | |
| CYCLOMETHICONE 5 (UNII: 0THT5PCI0R) | |
| DIMETHICONE (UNII: 92RU3N3Y1O) | |
| CYCLOMETHICONE 6 (UNII: XHK3U310BA) | |
| POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7) | |
| HEXYL LAURATE (UNII: 4CG9F9W01Q) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| SODIUM CHLORIDE (UNII: 451W47IQ8X) | |
| PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL) | |
| PHENOXYETHANOL (UNII: HIE492ZZ3T) | |
| COFFEE BEAN (UNII: JFH385Y744) | |
| PEG/PPG-18/18 DIMETHICONE (UNII: 9H0AO7T794) | |
| ALUMINUM OXIDE (UNII: LM26O6933) | |
| BUTYLENE GLYCOL (UNII: 3XUS85K0RA) | |
| OCTYLDODECANOL (UNII: 461N1O614Y) | |
| CAPRYLYL GLYCOL (UNII: 00YIU5438U) | |
| .ALPHA.-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0) | |
| CHLORPHENESIN (UNII: I670DAL4SZ) | |
| BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K) | |
| BENZYL ALCOHOL (UNII: LKG8494WBH) | |
| POTASSIUM SORBATE (UNII: 1VPU26JZZ4) | |
| CARBOXYPOLYMETHYLENE (UNII: 0A5MM307FC) | |
| POLYSORBATE 20 (UNII: 7T1F30V5YH) | |
| PALMITOYL TRIPEPTIDE-1 (UNII: RV743D216M) | |
| PALMITOYL TETRAPEPTIDE-7 (UNII: Q41S464P1R) | |
| FERRIC OXIDE RED (UNII: 1K09F3G675) | |
| TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E) | |
| HYDROGEN DIMETHICONE (13 CST) (UNII: 4QGR4P2YOI) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|------------------------------------------------------|----------------------|--------------------|
| 1 | NDC:14783-920-02 | 1 in 1 BOX | 12/02/2019 | |
| 1 | NDC:14783-920-01 | 30 mL in 1 BOTTLE; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|------------------------------------------|----------------------|--------------------|
| OTC MONOGRAPH FINAL | part352 | 12/02/2019 | |

LBEL EFFET PARFAIT SIGNS OF AGE APPEARANCE MINIMIZER FOUNDATION SPF 20 FM MEDIUM 6 - BEIGE

octinoxate and titanium dioxide lotion

Product Information

| | | | |
|-------------------------|----------------|--------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:14783-921 |
| Route of Administration | TOPICAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--------------------------------------------------------------------------|-------------------|------------------|
| Octinoxate (UNII: 4Y5P7MUD51) (Octinoxate - UNII:4Y5P7MUD51) | Octinoxate | 0.07 g in 1 mL |
| Titanium Dioxide (UNII: 15FIX9V2JP) (Titanium Dioxide - UNII:15FIX9V2JP) | Titanium Dioxide | 0.0244 g in 1 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|-----------------------------------------------|----------|
| WATER (UNII: 059QF0KO0R) | |
| CYCLOMETHICONE 5 (UNII: 0TH5PC10R) | |
| DIMETHICONE (UNII: 92RU3N3Y1O) | |
| CYCLOMETHICONE 6 (UNII: XHK3U310BA) | |
| POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7) | |
| HEXYL LAURATE (UNII: 4CG9F9W01Q) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| SODIUM CHLORIDE (UNII: 451W47IQ8X) | |
| PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL) | |
| PHENOXYETHANOL (UNII: HIE492ZZ3T) | |
| COFFEE BEAN (UNII: JFH385Y744) | |
| PEG/PPG-18/18 DIMETHICONE (UNII: 9H0AO7T794) | |
| ALUMINUM OXIDE (UNII: LM26O6933) | |
| BUTYLENE GLYCOL (UNII: 3XUS85K0RA) | |
| OCTYLDODECANOL (UNII: 461N1O614Y) | |
| CAPRYLYL GLYCOL (UNII: 00YIU5438U) | |
| .ALPHA.-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0) | |
| CHLORPHENESIN (UNII: I670DAL4SZ) | |
| BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K) | |
| BENZYL ALCOHOL (UNII: LKG8494WBH) | |
| POTASSIUM SORBATE (UNII: 1VPU26JZZ4) | |
| CARBOXYPOLYMETHYLENE (UNII: 0A5MM307FC) | |
| POLYSORBATE 20 (UNII: 7T1F30V5YH) | |
| PALMITOYL TRIPEPTIDE-1 (UNII: RV743D216M) | |
| PALMITOYL TETRAPEPTIDE-7 (UNII: Q41S464P1R) | |
| FERRIC OXIDE RED (UNII: 1K09F3G675) | |

TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)
HYDROGEN DIMETHICONE (13 CST) (UNII: 4QGR4P2YO1)

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|------------------------------------------------------|----------------------|--------------------|
| 1 | NDC:14783-921-02 | 1 in 1 BOX | 12/02/2019 | |
| 1 | NDC:14783-921-01 | 30 mL in 1 BOTTLE; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|------------------------------------------|----------------------|--------------------|
| OTC MONOGRAPH FINAL | part352 | 12/02/2019 | |

LBEL EFFET PARFAIT SIGNS OF AGE APPEARANCE MINIMIZER FOUNDATION SPF 20 FM MEDIUM 7 - BEIGE

octinoxate and titanium dioxide lotion

Product Information

| | | | |
|-------------------------|----------------|--------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:14783-922 |
| Route of Administration | TOPICAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--------------------------------------------------------------------------|-------------------|------------------|
| Octinoxate (UNII: 4Y5P7MUD51) (Octinoxate - UNII:4Y5P7MUD51) | Octinoxate | 0.07 g in 1 mL |
| Titanium Dioxide (UNII: 15FIX9V2JP) (Titanium Dioxide - UNII:15FIX9V2JP) | Titanium Dioxide | 0.0244 g in 1 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|-----------------------------------------------|----------|
| WATER (UNII: 059QF0KO0R) | |
| CYCLOMETHICONE 5 (UNII: 0THT5PC10R) | |
| DIMETHICONE (UNII: 92RU3N3Y1O) | |
| CYCLOMETHICONE 6 (UNII: XHK3U310BA) | |
| POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7) | |
| HEXYL LAURATE (UNII: 4CG9F9W01Q) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| SODIUM CHLORIDE (UNII: 451W47IQ8X) | |
| PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL) | |
| PHENOXYETHANOL (UNII: HIE492ZZ3T) | |
| COFFEE BEAN (UNII: JFH385Y744) | |
| PEG/PPG-18/18 DIMETHICONE (UNII: 9H0AO7T794) | |
| ALUMINUM OXIDE (UNII: LM26O6933) | |
| BUTYLENE GLYCOL (UNII: 3XUS85K0RA) | |

| | |
|--------------------------------------------------|--|
| OCTYLDODECANOL (UNII: 461N1O614Y) | |
| CAPRYLYL GLYCOL (UNII: 00YIU5438U) | |
| .ALPHA.-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0) | |
| CHLORPHENESIN (UNII: I670DAL4SZ) | |
| BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K) | |
| BENZYL ALCOHOL (UNII: LKG8494WBH) | |
| POTASSIUM SORBATE (UNII: 1VPU26JZZ4) | |
| CARBOXYPOLYMETHYLENE (UNII: 0A5MM307FC) | |
| POLYSORBATE 20 (UNII: 7T1F30V5YH) | |
| PALMITOYL TRIPEPTIDE-1 (UNII: RV743D216M) | |
| PALMITOYL TETRAPEPTIDE-7 (UNII: Q41S464P1R) | |
| FERRIC OXIDE RED (UNII: 1K09F3G675) | |
| TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E) | |
| HYDROGEN DIMETHICONE (13 CST) (UNII: 4QGR4P2YOI) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|------------------------------------------------------|----------------------|--------------------|
| 1 | NDC:14783-922-02 | 1 in 1 BOX | 12/02/2019 | |
| 1 | NDC:14783-922-01 | 30 mL in 1 BOTTLE; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|------------------------------------------|----------------------|--------------------|
| OTC MONOGRAPH FINAL | part352 | 12/02/2019 | |

LBEL EFFET PARFAIT SIGNS OF AGE APPEARANCE MINIMIZER FOUNDATION SPF 20 FM OBSCURE 8 - BROWN

octinoxate and titanium dioxide lotion

Product Information

| | | | |
|-------------------------|----------------|--------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:14783-923 |
| Route of Administration | TOPICAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--------------------------------------------------------------------------|-------------------|------------------|
| Octinoxate (UNII: 4Y5P7MUD51) (Octinoxate - UNII:4Y5P7MUD51) | Octinoxate | 0.07 g in 1 mL |
| Titanium Dioxide (UNII: 15FIX9V2JP) (Titanium Dioxide - UNII:15FIX9V2JP) | Titanium Dioxide | 0.0244 g in 1 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|-------------------------------------|----------|
| WATER (UNII: 059QF0KO0R) | |
| CYCLOMETHICONE 5 (UNII: 0THT5PC10R) | |
| DIMETHICONE (UNII: 92RU3N3Y1O) | |

| |
|--------------------------------------------------|
| CYCLOMETHICONE 6 (UNII: XHK3U310BA) |
| POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7) |
| HEXYL LAURATE (UNII: 4CG9F9W01Q) |
| GLYCERIN (UNII: PDC6A3C0OX) |
| SODIUM CHLORIDE (UNII: 451W47IQ8X) |
| PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL) |
| PHENOXYETHANOL (UNII: HIE492ZZ3T) |
| COFFEE BEAN (UNII: JFH385Y744) |
| PEG/PPG-18/18 DIMETHICONE (UNII: 9H0AO7T794) |
| ALUMINUM OXIDE (UNII: LM26O6933) |
| BUTYLENE GLYCOL (UNII: 3XUS85K0RA) |
| OCTYLDODECANOL (UNII: 461N1O614Y) |
| CAPRYLYL GLYCOL (UNII: 00YIU5438U) |
| .ALPHA.-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0) |
| CHLORPHENESIN (UNII: I670DAL4SZ) |
| BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K) |
| BENZYL ALCOHOL (UNII: LKG8494WBH) |
| POTASSIUM SORBATE (UNII: 1VPU26JZZ4) |
| CARBOXYPOLYMETHYLENE (UNII: 0A5MM307FC) |
| POLYSORBATE 20 (UNII: 7T1F30V5YH) |
| PALMITOYL TRIPEPTIDE-1 (UNII: RV743D216M) |
| PALMITOYL TETRAPEPTIDE-7 (UNII: Q41S464P1R) |
| FERRIC OXIDE RED (UNII: 1K09F3G675) |
| TRIETHOXYCAPRYLSILANE (UNII: LDC331P08E) |
| HYDROGEN DIMETHICONE (13 CST) (UNII: 4QGR4P2YOI) |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|------------------------------------------------------|----------------------|--------------------|
| 1 | NDC:14783-923-02 | 1 in 1 BOX | 12/02/2019 | |
| 1 | NDC:14783-923-01 | 30 mL in 1 BOTTLE; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|------------------------------------------|----------------------|--------------------|
| OTC MONOGRAPH FINAL | part352 | 12/02/2019 | |

LBEL EFFET PARFAIT SIGNS OF AGE APPEARANCE MINIMIZER FOUNDATION SPF 20 FM OBSCURE 9 - BROWN

octinoxate and titanium dioxide lotion

Product Information

| | | | |
|-------------------------|----------------|--------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:14783-924 |
| Route of Administration | TOPICAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--------------------------------------------------------------------------|-------------------|------------------|
| Octinoxate (UNII: 4Y5P7MUD51) (Octinoxate - UNII:4Y5P7MUD51) | Octinoxate | 0.07 g in 1 mL |
| Titanium Dioxide (UNII: 15FIX9V2JP) (Titanium Dioxide - UNII:15FIX9V2JP) | Titanium Dioxide | 0.0244 g in 1 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|--------------------------------------------------|----------|
| WATER (UNII: 059QF0KO0R) | |
| CYCLOMETHICONE 5 (UNII: 0THT5PCI0R) | |
| DIMETHICONE (UNII: 92RU3N3Y1O) | |
| CYCLOMETHICONE 6 (UNII: XHK3U310BA) | |
| POLYGLYCERYL-4 ISOSTEARATE (UNII: 820DPX33S7) | |
| HEXYL LAURATE (UNII: 4CG9F9W01Q) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| SODIUM CHLORIDE (UNII: 451W47IQ8X) | |
| PEG/PPG-19/19 DIMETHICONE (UNII: EHH90CO7TL) | |
| PHENOXYETHANOL (UNII: HIE49ZZZ3T) | |
| COFFEE BEAN (UNII: JFH385Y744) | |
| PEG/PPG-18/18 DIMETHICONE (UNII: 9H0AO7T794) | |
| ALUMINUM OXIDE (UNII: LM26O6933) | |
| BUTYLENE GLYCOL (UNII: 3XUS85K0RA) | |
| OCTYLDODECANOL (UNII: 461N1O614Y) | |
| CAPRYLYL GLYCOL (UNII: 00YIU5438U) | |
| .ALPHA.-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0) | |
| CHLORPHENESIN (UNII: I670DAL4SZ) | |
| BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K) | |
| BENZYL ALCOHOL (UNII: LKG8494WBH) | |
| POTASSIUM SORBATE (UNII: 1VPU26JZZ4) | |
| CARBOXYPOLYMETHYLENE (UNII: 0A5MMB07FC) | |
| POLYSORBATE 20 (UNII: 7T1F30V5YH) | |
| PALMITOYL TRIPEPTIDE-1 (UNII: RV743D216M) | |
| PALMITOYL TETRAPEPTIDE-7 (UNII: Q41S464P1R) | |
| FERRIC OXIDE RED (UNII: 1K09F3G675) | |
| TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E) | |
| HYDROGEN DIMETHICONE (13 CST) (UNII: 4QGR4P2YOI) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|------------------------------------------------------|----------------------|--------------------|
| 1 | NDC:14783-924-02 | 1 in 1 BOX | 12/02/2019 | |
| 1 | NDC:14783-924-01 | 30 mL in 1 BOTTLE; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|------------------------------------------|----------------------|--------------------|
| OTC MONOGRAPH FINAL | part352 | 12/02/2019 | |

Labeler - Ventura International LTD (603192787)

Establishment

| Name | Address | ID/FEI | Business Operations |
|-----------------------------|---------|-----------|----------------------------------------------------------------------------------------------------------------|
| Bel Star S.A. (Colombia) | | 880160197 | MANUFACTURE(14783-916, 14783-917, 14783-918, 14783-919, 14783-920, 14783-921, 14783-922, 14783-923, 14783-924) |

Revised: 12/2019

Ventura International LTD