

MEDIQUE ALCALAK- calcium carbonate tablet, chewable
MEDI FIRST ANTACID- calcium carbonate tablet, chewable
MEDI FIRST PLUS ANTACID- calcium carbonate tablet, chewable
Unifirst First Aid Corporation

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Unifirst First Aid Antacid

Drug Facts

Active ingredient (in each tablet)

Calcium Carbonate 420 mg

Purpose

Antacid

Uses

relieves

- acid indigestion
- sour stomach
- heartburn
- upset stomach associated with these symptoms

Warnings

Do not use

■ the maximum dosage of this product for more than 2 weeks except under the advice and supervision of a doctor

Ask a doctor or pharmacist before use if you

- are taking a prescription drug. Antacids may interact with certain prescription drugs.
- have kidney disease

When using this product

- do not take more than 19 tablets in a 24 hour period. If symptoms persist for more than two weeks, stop using this product and see a doctor.

If pregnant or breast feeding ,ask a health professional before use.

Keep out of the reach of children.

Directions

Adults and children: (12 years and older) Chew 2 tablets every 2 - 3 hours or as symptoms occur, repeat hourly if needed. Do not exceed 19 tablets in 24 hours.

Children under 12 years: Ask a doctor

Other information

- calcium content per tablet: 168 mg
- **phenylketonurics:** each tablet may contain 1.5 mg phenylalanine
- store at room temperature 59-86°F (15-30°C) in a dry place
- tamper-evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

aspartame*, croscarmellose sodium*, gum acacia*, magnesium stearate, maltodextrin, mineral oil*, mint flavor, sorbitol*, sucrose*

**may contain*

Questions or comments Call 1-800-634-7680

Medi-First Plus Antacid Label

100 Tablets

(50 x 2's)

MEDI-FIRST® Plus

Antacid

Calcium Carbonate 420mg

Carbonato de Calcio 420mg

Pull to Open

Tire Para Abrir

Calcium Rich

Chewable Tablets

Tabletas Masticables, Ricas en Calcio

Compare active ingredient to:

TUMS®


Registered Trademark of Smithkline-Beecham

Tamper Evident

Unit Dose Packets

Empaquetado con sellado

evidente en dosis unitarias



Medi-First Antacid Label

100
Tablets
(50 x 2)

MEDI-FIRST®

Antacid Calcium Carbonate

Compare active ingredient to:

Compare el ingrediente activo con:

Tums®

Registered Trademark of Smithkline-Beecham

Marca Registrada de Smithkline-Beecham

Pull to Open

Tire Para Abrir

Heartburn, Indigestion • Calcium Carbonate 420mg

las agruras, la indigestion • Antiácido Carbonato de Calcio 420mg

Antiácido Carbonato de Calcio

Tamper Evident Unit Dose Packets

Empaquetado con Sellado Evidente en Dosis Unitarias

MEDI-FIRST®
Antacid Calcium Carbonate

Compare active ingredient to:
Compare el ingrediente activo con:

TUMS®
Compare active ingredient to:
Compare el ingrediente activo con:

Heartburn, Indigestion - Calcium Carbonate 420 mg
Los gases, la indigestión - Carbonato de Calcio 420 mg

Antiácido Carbonato de Calcio

Tamper Evident Unit Dose Packets
Empaquetado con Sellado Evidente en Dosis Unitarias

8023380X
Rev. 4/13/21
Mfg. 1,14

| Drug Facts | Drug Facts (continued) |
|--|---|
| <p>Active ingredient (in each tablet) Purpose Calcium Carbonate 420 mg Antacid</p> <p>Uses relieves ■ acid indigestion ■ sour stomach ■ heartburn ■ upset stomach associated with these symptoms</p> <p>Warnings Do not use ■ the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a doctor Ask a doctor or pharmacist before use if you ■ are taking a prescription drug. Antacids may interact with certain prescription drugs. ■ have kidney disease</p> <p>When using this product ■ do not take more than 19 tablets in a 24 hour period. If symptoms persist for more than two weeks, stop using this product and see a doctor. If pregnant or breastfeeding, ask a health professional before use. Keep out of the reach of children.</p> | <p>Directions Adults and children (12 years and older) Chew 2 tablets every 2 - 3 hours or as symptoms occur, repeat hourly if needed. Do not exceed 19 tablets in 24 hours. Children under 12 years Ask a doctor</p> <p>Other information ■ calcium content per tablet: 168 mg ■ phenylethanolamines: each tablet may contain 1.5 mg phenylethanolamine ■ store at room temperature 59°-86°F (15°-30°C) in a dry place ■ do not use any opened or torn packets</p> <p>Inactive ingredients aspartame*, croscarmellose sodium*, gum acacia*, magnesium stearate, maltodextrin, mineral oil*, mint flavor, sorbitol*, sucrose* *may contain</p> <p>Questions or comments Call 1-800-634-7680</p> |

MEDI-FIRST®

Antacid

Calcium Carbonate

Compare active ingredient to:
Compare el ingrediente activo con:

TUMS®

Registered Trademark of Takeda Pharmaceuticals North America
Marca Registrada de Takeda Pharmaceuticals North America

Heartburn, Indigestion - Calcium Carbonate 420 mg
Los gases, la indigestión - Carbonato de Calcio 420 mg

Antiácido Carbonato de Calcio

Tamper Evident Unit Dose Packets
Empaquetado con Sellado Evidente en Dosis Unitarias

| Información del Medicamento | Información del Medicamento (continuación) |
|---|--|
| <p>Ingrediente activo (en cada tableta) Propiedad Carbonato de Calcio - 420 mg Antácido</p> <p>Usos alivia ■ la indigestión por acidez ■ la acidez estomacal ■ los gases ■ malestar estomacal asociado a estos síntomas</p> <p>Advertencias No consuma ■ la dosis máxima de este producto por más de 2 semanas, excepto bajo indicaciones y supervisión de un médico Consulte con el doctor o un farmacéutico antes de consumir este medicamento si usted ■ está actualmente tomando un medicamento recetado. Los medicamentos pueden interactuar con ciertos medicamentos recetados. ■ tiene enfermedad renal Cuando está usando este producto ■ no tomar más de 19 tabletas en un periodo de 24 horas. Si los síntomas persisten por más de dos semanas, deje de tomar este producto y consulte a un médico. Si está embarazada o en período de lactancia, consulte la opinión de un médico antes de consumir este producto.</p> | <p>Mantenga este producto fuera del alcance de los niños.</p> <p>Instrucciones Adultos y niños (para mayores de 12 años) Masticar 2 tabletas cada 2 o 3 horas o como se presenten los síntomas, repetir cada una hora si es necesario. No consuma más de 19 tabletas en 24 horas. Niños menores de 12 años: Consultar al médico</p> <p>Otra información ■ cada tableta contiene 168 mg de calcio elemental ■ fenilethanolaminas: puede contener 1.5 mg de fenilethanolamina por tableta ■ almacenamiento: consérvese a temperatura ambiente 59°-86°F (15°-30°C) y en un lugar seco ■ empaquetado con sellado evidente ■ no use ningún paquete abierto o roto</p> <p>Ingredientes inactivos aspartame*, croscarmellose sodica*, goma de acacia*, estearato de magnesio, maltodextrina, aceite mineral*, sabor de la menta, sorbitol*, sacarosa* *puede contener</p> <p>¿Tiene alguna pregunta o comentario? Lláme al 1-800-634-7680</p> |

Retain carton for complete product information

Medique Alcalak Label

Medique ®

Alcalak

Calcium-Rich Antacid

Mint Flavor

Heartburn, Indigestion • Calcium Carbonate 420 mg

6

Tablets

(3 x 2)

Tamper Evident Unit Dose Packets

| Drug Facts | Purpose |
|---|---------|
| <p>Active ingredients (in each tablet) Calcium Carbonate 420 mg</p> <p>Uses relieves ■ acid indigestion ■ sour stomach ■ heartburn ■ upset stomach associated with these symptoms</p> <p>Warnings Do not use ■ the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a doctor Ask a doctor or pharmacist before use if you ■ are taking a prescription drug. Antacids may interact with certain prescription drugs. ■ have kidney disease</p> <p>When using this product ■ do not take more than 19 tablets in a 24 hour period. If symptoms persist for more than two weeks, stop using this product and see a doctor. If pregnant or breastfeeding, ask a health professional before use. Keep out of the reach of children.</p> | Antacid |

Alcalak

Medique®

Alcalak

Calcium-Rich Antacid

Mint Flavor

Heartburn, Indigestion • Calcium Carbonate 420 mg

6
Tablets
(3 x 2)

Manufactured for
Medique Products
Fort Myers, FL 33967
1-800-634-7680
www.mediqueproducts.com

1016980X
Rev. 4/13/21
Mfg. 1,14

Reorder #10169

3 47682 10169 5

RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION

Tamper Evident Unit Dose Packets

| Drug Facts (continued) | |
|------------------------------|--|
| Directions | Chew 2 tablets every 2 - 3 hours or as symptoms occur, repeat hourly if needed. Do not exceed 19 tablets in 24 hours. |
| Children under 12 years: | Ask a doctor |
| Other information | ■ calcium content per tablet: 168 mg ■ phenylethanolamines: each tablet may contain 1.5 mg phenylethanolamine ■ store at room temperature 59°-86°F (15°-30°C) in a dry place ■ tamper-evident sealed packets ■ do not use any opened or torn packets |
| Inactive ingredients | aspartame*, croscarmellose sodium*, gum acacia*, magnesium stearate, maltodextrin, mineral oil*, mint flavor, sorbitol*, sucrose* *may contain |
| Questions or comments | Call 1-800-634-7680 |

MEDIQUE ALCALAK

calcium carbonate tablet, chewable

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:47682-088 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|--------------------------|-----------------|
| CALCIUM CARBONATE (UNII: H0G9379FGK) (CALCIUM CATION - UNII:2M83C4R6ZB, CARBONATE ION - UNII:7UJQ5OPE7D) | CALCIUM CARBONATE | 420 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|--|-----------------|
| ACACIA (UNII: 5C5403N26O) | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| MALTODEXTRIN (UNII: 7CVR7L4A2D) | |
| SUCROSE (UNII: C151H8M554) | |

Product Characteristics

| | | | |
|-----------------|---------------|---------------------|----------|
| Color | white (white) | Score | no score |
| Shape | ROUND (ROUND) | Size | 12mm |
| Flavor | MINT (MINT) | Imprint Code | FR;8 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|----------|------------------|--|-----------------------------|---------------------------|
| 1 | NDC:47682-088-69 | 3 in 1 BOX | 12/30/2008 | |
| 1 | | 2 in 1 PACKET; Type 0: Not a Combination Product | | |
| 2 | NDC:47682-088-64 | 12 in 1 BOX | 12/30/2008 | |
| 2 | | 2 in 1 PACKET; Type 0: Not a Combination Product | | |
| 3 | NDC:47682-088-33 | 50 in 1 BOX | 12/30/2008 | |
| 3 | | 2 in 1 PACKET; Type 0: Not a Combination Product | | |
| 4 | NDC:47682-088-47 | 100 in 1 BOX | 12/30/2008 | |
| 4 | NDC:47682-088-99 | 2 in 1 PACKET; Type 0: Not a Combination Product | | |
| 5 | NDC:47682-088-13 | 250 in 1 BOX | 12/30/2008 | |
| 5 | | 2 in 1 PACKET; Type 0: Not a Combination Product | | |
| 6 | NDC:47682-088-99 | 2 in 1 PACKET; Type 0: Not a Combination Product | 12/30/2008 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC monograph final | part331 | 12/30/2008 | |

MEDI FIRST ANTACID

calcium carbonate tablet, chewable

Product Information

| | | | |
|-------------------------|----------------|--------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:47682-089 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------|----------|
| CALCIUM CARBONATE (UNII: H0G9379FGK) (CALCIUM CATION - UNII:2M83C4R6ZB, CARBONATE ION - UNII:7UJQ5OPE7D) | CALCIUM CARBONATE | 420 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| ACACIA (UNII: 5C5403N26O) | |
| MAGNESIUM STEARATE (UNII: 70097M6I3O) | |
| MALTODEXTRIN (UNII: 7CVR7L4A2D) | |
| SUCROSE (UNII: C151H8M554) | |

Product Characteristics

| | | | |
|----------|---------------|--------------|----------|
| Color | white (white) | Score | no score |
| Shape | ROUND (ROUND) | Size | 12mm |
| Flavor | MINT (MINT) | Imprint Code | FR;8 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:47682-089-13 | 250 in 1 BOX | 12/30/2008 | |
| 1 | | 2 in 1 PACKET; Type 0: Not a Combination Product | | |
| 2 | NDC:47682-089-48 | 125 in 1 BOX | 12/30/2008 | |
| 2 | | 2 in 1 PACKET; Type 0: Not a Combination Product | | |

| | | | | |
|---|------------------|--|------------|--|
| 3 | NDC:47682-089-33 | 50 in 1 BOX | 12/30/2008 | |
| 3 | NDC:47682-089-99 | 2 in 1 PACKET; Type 0: Not a Combination Product | | |
| 4 | NDC:47682-089-99 | 2 in 1 PACKET; Type 0: Not a Combination Product | 12/30/2008 | |
| 5 | NDC:47682-089-50 | 25 in 1 BOX | 04/01/2019 | |
| 5 | | 2 in 1 PACKET; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC monograph final | part331 | 12/30/2008 | |

MEDI FIRST PLUS ANTACID

calcium carbonate tablet, chewable

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:47682-092 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------|----------|
| CALCIUM CARBONATE (UNII: H0G9379FGK) (CALCIUM CATION - UNII:2M83C4R6ZB, CARBONATE ION - UNII:7UJQ5OPE7D) | CALCIUM CARBONATE | 420 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| ACACIA (UNII: 5C5403N260) | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| MALTODEXTRIN (UNII: 7CVR7L4A2D) | |
| SUCROSE (UNII: C151H8M554) | |

Product Characteristics

| | | | |
|-----------------|---------------|---------------------|----------|
| Color | white (white) | Score | no score |
| Shape | ROUND (ROUND) | Size | 12mm |
| Flavor | MINT (MINT) | Imprint Code | FR;8 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:47682-092-33 | 50 in 1 BOX | 12/30/2008 | |
| 1 | | 2 in 1 PACKET; Type 0: Not a Combination Product | | |
| 2 | NDC:47682-092-48 | 125 in 1 BOX | 12/30/2008 | |
| 2 | | 2 in 1 PACKET; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------|--|----------------------|--------------------|
| OTC monograph final | part331 | 12/30/2008 | |

Labeler - Unifirst First Aid Corporation (832947092)

Revised: 3/2022

Unifirst First Aid Corporation