

ARONAMIN GOLD- fursultiamine, riboflavin tetrabutryate, pyridoxal phosphate, hydroxocobalamin acetate, ascorbic acid, tocopheryl acetate tablet, film coated
OASIS TRADING

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Fursultiamine(Active vitamin B1) 50.0mg
Riboflavin tetrabutryate(Active vitamin B2) 2.5mg
Pyridoxal phosphate(Active vitamin B6) 2.5mg
Hydroxocobalamin acetate(Active vitamin B12) 5.22µg
Ascorbic acid(Vitamin C) 70mg
Tocopheryl acetate(Vitamin E) 20.0mg

Vitamines, Mineral & nutrients

Keep out of reach of children

Adults: take 1 tablet twice a day.

1) Do not take this medicine.

(1) Patients with hypersensitivity reactions to NSAIDs and the components contained in NSAIDs

(2) Infants under three months of age

(3) Because this drug contains lactose, patients with genetic problems such as galactose intolerance, Lapp lactase deficiency, or glucose-galactose malabsorption Do not administer.

2) Do not take the following medicine while taking this medicine.

(1) levodopa

3) Consult a doctor, dentist or pharmacist before taking this medicine.

(1) Patients receiving medical treatment

(2) patients with hyperoxaluria (excessive urinary excretion of urine)

(3) Pregnant women and possibly pregnant women, lactating women, premature infants, infants

(4) Patients with gout or kidney stones

4) Stop taking this drug immediately and consult a doctor, dentist, or pharmacist if you: Whenever possible, bring this attached document with you.

(1) If you have any of the following symptoms

Stomach discomfort, diarrhea, constipation, rash, fever, nausea, vomiting, dilated stool, stomatitis (mouth salt), anorexia, abdominal bloating

(2) Administration of this drug may result in faster or more frequent menstruation, and bleeding may last for a long time.

(3) The risk of thrombosis may be increased if a woman taking an oral contraceptive containing estrogen or a patient with thrombotic placenta is taking vitamin E.

(4) Long-term administration of high doses may result in resistance.

(5) Prolonged use of pyridoxine at a dose of 500 mg to 2 g / day may result in neuropathy or neuropathy (functional disorder or pathologic changes) in the peripheral nervous system.

(6) Hematologic adverse events may occur when vitamin B12 is administered at a dose of 10 µg or more per day in patients with insufficient folic acid.

5) Other Precautions for Taking

- (1) Keep the prescribed dosage and dosage.
- (2) it may interfere with the detection of blood glucose during various urine tests
- (3) The urine may turn yellow, which may affect the clinical examination

6) Storage Precautions

- (1) Keep out of the reach of children.
- (2) Avoid direct sunlight. Store in a cool, dry place.
- (3) To prevent misuse (misuse) and to preserve quality, please do not put in another container.

Microcrystalline Cellulose, Magnesium Stearate, Crospovidone, Copovidone, Colloidal Silicon Dioxide, OY-25014, OY-S-29019

For oral use only



ARONAMIN GOLD

fursultiamine, riboflavin tetrabutryate, pyridoxal phosphate, hydroxocobalamin acetate, ascorbic acid, tocopheryl acetate tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72689-0024
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
RIBOFLAVIN TETRABUTYRATE (UNII: F211C9MSGY) (RIBOFLAVIN TETRABUTYRATE - UNII:F211C9MSGY)	RIBOFLAVIN TETRABUTYRATE	2.5 mg
FURSULTIAMINE (UNII: 05J61265PX) (FURSULTIAMINE - UNII:05J61265PX)	FURSULTIAMINE	50 mg
HYDROXOCOBALAMIN ACETATE (UNII: S535M27N3Q) (HYDROXOCOBALAMIN - UNII:Q40X8H422O)	HYDROXOCOBALAMIN	5.22 ug
.ALPHA.-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0) (.ALPHA.-TOCOPHEROL - UNII:H4N855PNZ1)	.ALPHA.-TOCOPHEROL ACETATE	20 mg
ASCORBIC ACID (UNII: PQ6CK8PD0R) (ASCORBIC ACID - UNII:PQ6CK8PD0R)	ASCORBIC ACID	70 mg

Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
COPOVIDONE K25-31 (UNII: D9C330MD8B)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSPVIDONE (UNII: 2S7830E561)	

Product Characteristics

Color	red	Score	no score
Shape	OVAL	Size	16mm
Flavor		Imprint Code	AroG
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72689-0024-1	100 in 1 BLISTER PACK; Type 0: Not a Combination Product	11/20/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		11/20/2018	

Labeler - OASIS TRADING (689991468)

Registrant - OASIS TRADING (689991468)

Establishment

Name	Address	ID/FEI	Business Operations
OASIS TRADING		689991468	manufacture(72689-0024) , label(72689-0024)

Revised: 12/2018

OASIS TRADING