

IBUPROFEN- ibuprofen tablet
FAMILY DOLLAR

SEE NEW WARNINGS

Ibuprofen

Tablets USP, 200 mg

Pain Reliever/Fever Reducer (NSAID)**

Active ingredient (For Tablet)

(in each tablet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Active ingredient (For Caplet)

(in each caplet)

Ibuprofen 200 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- minor pain of arthritis
- toothache
- backache
- the common cold
- menstrual cramps
- temporarily reduces fever

Allergy alert:

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning:

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of the reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions (For Tablets)

- **do not take more than directed**
- **the smallest effective dose should be used**

Adults and children 12 years and older

- take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor

Children under 12 years

- ask a doctor

Directions (For Caplets)

- **do not take more than directed**
- **the smallest effective dose should be used**

Adults and children 12 years and older

- take 1 caplet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 caplet, 2 caplets may be used
- do not exceed 6 caplets in 24 hours, unless directed by a doctor

Children under 12 years

- ask a doctor

Other information

- store between 20-25°C (68-77°F). [See USP Controlled Temperature]
- avoid excessive heat above 40°C (104°F)

Inactive ingredients

colloidal silicon dioxide, ferric oxide red, maize starch, polyethylene glycol, povidone k-30, pregelatinized starch, sodium starch glycolate, stearic acid, talc, titanium dioxide.

Questions or comments?

call **1-877-770-3183** weekdays 9:00 AM to 4:30PM EST.

Tablets & Caplets

code # :
 size : 2 X 2 X 4
 ref # : PE190811A r/w : PE19T 8TD
 material : 016 BSS date : 10/10/201

0 52251 26457 5
 581 082828
 For a refund (with receipt) or assistance
 within 90 days of any Family Dollar store
 return package and unexpired product
 to: Family Dollar, Inc., 1101
 500 ONYX Parkway, Chesapeake, VA 23020
 Distributed By: Millwood Brands, LLC

R-01
 53200008
 300000 111

Ibuprofen
 Tablets USP, 200 mg
 Pain Reliever/Fever Reducer (NSAID) #
 100 Coated Caplets®
 FAMILY wellness™
 COMPARE TO ASPIRIN®
 OTC
 Actual Size

Important: Read all product information before using.
 Keep this box for important information.
 Do not use if full lower seal is broken or missing.

Drug Facts
Active Ingredient **Purpose**
 (in each caplet)
 ibuprofen 200 mg..... Pain reliever/fever reducer
 (NSAID)
 *nonsteroidal anti-inflammatory drug

Uses
 temporarily relieve minor aches and pains due to:
 headache muscular aches minor pain of
 arthritis toothache backache in the common cold
 menstrual cramps
 temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause severe allergic reaction,
 especially in people allergic to aspirin.
 Symptoms may include:
 hives facial swelling asthma (wheezing)
 shock skin redness rash blisters
 If an allergic reaction occurs, stop use and seek medical
 help right away.
Stomach bleeding warning: This product contains an
 NSAID, which may cause severe stomach bleeding. The
 chance is higher if you:
 are age 60 or older
 have had stomach ulcers or bleeding problems
 take a blood thinning (anticoagulant) or steroid drug
 take other drugs containing prescription or
 nonprescription NSAIDs (aspirin, ibuprofen, naproxen,
 or others)
 have 3 or more alcoholic drinks every day while using
 this product
 take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin
 increase the risk of heart attack, heart failure, and stroke.
 These can be fatal. The risk is higher if you use more than
 directed or for longer than directed.

Drug Facts (continued)
 Do not use
 if you have ever had an allergic reaction to ibuprofen or
 any other pain reliever/fever reducer
 in right before or after heart surgery
 Ask a doctor before use if
 you have problems or serious side effects from taking
 pain relievers or fever reducers
 the stomach bleeding warning applies to you
 you have a history of stomach problems, such as
 heartburn
 you have high blood pressure, heart disease, liver
 disease, kidney disease, asthma or had a stroke
 you are taking a diuretic
 Ask a doctor or pharmacist before use if you are
 taking aspirin for heart attack or stroke, because
 ibuprofen may decrease the benefit of aspirin
 under a doctor's care for any serious condition
 taking any other drug
 When using this product:
 do not take with food or milk if stomach upset occurs
 Stop use and ask a doctor if
 you experience any of the following signs of stomach
 bleeding:
 • black stool • vomit blood • loose bloody or black stools
 • have stomach pain that does not get better
 if you have symptoms of heart problems or stroke • chest
 pain • trouble breathing • weakness in one part or side
 of body • slurred speech • big swelling
 if pain gets worse or lasts more than 10 days
 if fever gets worse or lasts more than 3 days
 if redness or swelling is present in the painful area
 if any new symptoms appear

If pregnant or breast-feeding, ask a health professional
 before use. It is especially important not to use ibuprofen
 during the last 3 months of pregnancy unless definitely
 directed to do so by a doctor because it may cause
 problems in the unborn child or complications during
 delivery.
 Keep out of the reach of children. In case of overdose, ...

Drug Facts (continued)
 get medical help or contact a Poison Control Center right
 away; 1-800-222-1222

Directions
 do not take more than directed
 in the smallest effective dose should be used
 adults and children 12 years and older
 take 1 caplet every 4 to 6 hours
 while symptoms persist
 if pain or fever does not
 respond to 1 caplet, 2 caplets
 may be used
 do not exceed 6 caplets in
 24 hours, unless directed by
 a doctor
 children under 12 years
 ask a doctor

Other Information
 store between 20-25°C (68-77°F). (See USP Controlled
 Room Temperature).
 avoid excessive heat above 40°C (104°F)

Inactive Ingredients calcium stearate, croscellose, iron
 oxide red, maize starch, polyethylene glycol, polydextrose
 K-30, pregelatinized starch, sodium starch glycolate,
 stearic acid, talc, titanium dioxide

Questions or comments?
 call 1-877-770-3183
 weekdays 9:00 AM to 4:30 PM EST

"This product is not manufactured or distributed by Wyeth Consumer
 Healthcare, distributor of Advil."
MADE IN INDIA
 Contains no ingredient made from a gluten-containing grain
 (wheat, barley, or rye)

300000 111
 53200008
 R-01

Lot
 Exp.

item	: 1780 B200N 50CP	view	: 100000
code #	:	date	: 10/27/11
size	: 1x3/4 X 1x3/4 X 3x1/4		
ref #	: PPF160911C		
material	: C15 08S		



MADE IN INDIA
 0171 9046 8118/8178
 500 VINDA Parkway, Chesapeake, VA 23020
 Distributed By: Millimed Pharmaceuticals, LLC
 Dominion Healthcare, a Division of Amgen
 This product is not manufactured or distributed by Amgen

R-01
 532000014
 300000159

50 Coated Caplets®
Ibuprofen
 Tablets USP, 200 mg
 Pain Reliever/Fever Reducer (NSAID)
 OTC
 FAMILY wellness™
 CONTAINS 10 TABLETS

Important: Read all product information before using. Keep this box for important information. Do not use if all boxes are broken or missing.

Drug Facts
Active Ingredient (in each caplet): Purpose
 Ibuprofen 200 mg.....Pain reliever/fever reducer (NSAID)
 *Nonsteroidal anti-inflammatory drug

Uses
 to temporarily relieve minor aches and pains due to:
 in headache in muscle aches in minor pain of arthritis in toothache or backache in the common cold
 in menstrual cramps
 to temporarily reduce fever

Warnings
 Allergy alert: Ibuprofen may cause severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives; facial swelling; difficulty breathing
 in shock or dizziness
 If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause stomach bleeding. The chance is higher if you are age 60 or older
 in have had stomach ulcers or bleeding problems
 in take a blood thinner (anticoagulant) or alcohol drug
 in take other drugs containing aspirin or nonsteroidal NSAIDs (aspirin, naproxen, or others)
 in have 3 or more alcoholic drinks every day while using this product
 in take more or for a longer time than directed

Drug Facts (continued)
Heart attack and stroke warnings: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
 Do not use
 in if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
 in right before or after heart surgery
Ask a doctor before use if
 in you have problems or serious side effects from taking pain relievers or fever reducers
 in the stomach bleeding warning applies to you
 in you have a history of stomach problems, such as heartburn
 in you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
 in you are taking a diuretic
Ask a doctor or pharmacist before use if you are
 in taking aspirin for heart attack or stroke, because ibuprofen may decrease the benefit of aspirin
 in under a doctor's care for any serious condition
 in taking any other drug
When using this product
 in take with food or milk if stomach upset occurs
 in take with food or milk if stomach upset occurs
Stop use and seek medical help if
 in you experience any of the following signs of stomach bleeding: • red, black or maroon stools • dark, tarry stools • vomit that looks like coffee grounds • black stools • have stomach pain that does not get better
 in you have symptoms of heart problems or stroke • chest pain • trouble breathing • weakness in one part or side of body • slurred speech • leg swelling or pain • pale, warm or blue more than 10 days in lower leg or foot more than 3 days in women or swelling is present in the painful area

Drug Facts (continued)
 in any new symptoms appear
 in pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
 Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions
 in do not take more than directed
 in the smallest effective dose should be used
 adults and children
 12 years and older
 in take 1 caplet every 4 to 6 hours
 while symptoms persist
 in if pain or fever does not respond to 1 caplet, 2 caplets may be used
 in do not exceed 6 caplets in 24 hours, unless directed by a doctor
 children under 12 years
 in ask a doctor

Other Information
 in store between 20-25°C (68-77°F). [See USP Controlled Room Temperature], in avoid excessive heat above 40°C (104°F)
 Ibuprofen Ingestible® contains silicon dioxide, iron oxide red, microcrystalline cellulose, polyethylene glycol, polydioxane K-30, pregelatinized starch, sodium starch glycolate

300000159
 532000014
 R-01

Drug Facts (continued)
 glycolate, stearic acid, talc, titanium dioxide
Questions or comments?
 call 1-877-776-3110
 weekdays 8:00 AM to 4:30 PM EST
 Contains 10 caplets made from a gluten-containing grain (wheat, barley, or rye)

Lot
 Exp.

3

← 1 3/4 → ← 1 3/4 →

PIONEER PKG	
end user :	GRANDTAS
item :	120 ORANGE 100CT
code # :	
size :	1 3/4 X 1 3/4 X 3 1/2
ref # :	ZPL006113
material :	.016 SBS
view :	FR
date :	08



MADE IN INDIA
 NOT 100% SUSTAINABLE
 508 YONK Partners, Chesapeake, VA 23820
 Distributed By: Mawood Brands, LLC
 Consumer Healthcare, distributor of Advil®
 This product is not manufactured or distributed by Mawood Brands, LLC

300000474
 532000019
 R-00

Ibuprofen
 Tablets USP, 200 mg
 Pain Reliever/Fever Reducer (NSAID)**

FAMILY
wellness
 CONTAINS NO ADDED SUGAR

24 Coated Tablets

OTC
 ADVIL

Actual Size

Important: Read all the product information before using. Keep this box for important information. Do not use if full inner seal is broken or missing.

Drug Facts
Active ingredient (in each tablet): Purpose
 Ibuprofen 200 mg.....Pain reliever/fever reducer (NSAID)
 *Nonsteroidal anti-inflammatory drug

Uses
 It temporarily relieves minor aches and pains due to:
 • headache or muscular aches or minor pain of arthritis
 • toothache or backache or the common cold
 • menstrual cramps
 • it temporarily reduces fever

Warnings
 Allergy alert: Ibuprofen may cause severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives or facial swelling or rash or wheezing or shock or skin redness or rash or blisters
 If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 • are age 60 or older
 • have had stomach ulcers or bleeding problems
 • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 • have 3 or more alcoholic drinks every day while using this product
 • take more or for a longer time than directed

Drug Facts (continued)

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use
 • if you have ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
 • right before or after heart surgery

Ask a doctor before use if
 • you have problems or serious side effects from taking pain relievers or fever reducers
 • the stomach bleeding warning applies to you
 • you have a history of stomach problems, such as heartburn or you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
 • you are taking a diuretic

Ask a doctor or pharmacist before use if you are
 • taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
 • under a doctor's care for any serious condition
 • taking any other drug

When using this product
 • take with food or milk, if stomach upset occurs

Stop use and ask a doctor if
 • you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better
 • you have symptoms of heart problems or stroke: • chest pain • trouble breathing • weakness in one part or side of body • altered speech • leg swelling or pain gets worse or lasts more than 10 days or lower leg swells or lasts more than 3 days if redness or swelling is present in the painful area

Drug Facts (continued)

• in any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

• do not take more than directed
 • in the smallest effective dose should be used

adults and children 12 years and older	• take 1 tablet every 4 to 6 hours while symptoms persist • if pain or fever does not respond to 1 tablet, 2 tablets may be used • do not exceed 6 tablets in 24 hours, unless directed by a doctor
children under 12 years	• ask a doctor

Other information

• store between 20-25°C (68-77°F). [See USP Controlled Room Temperature].
 • avoid excessive heat above 40°C (104°F)

Inactive ingredients: colloidal silicon dioxide, iron oxide red, maize starch, polyethylene glycol, polyvinyl K-30, pregelatinized starch, sodium starch glycolate, stearic acid, talc, titanium dioxide

Drug Facts (continued)

glycolate, stearic acid, talc, titanium dioxide


Questions or comments?
 call 1-877-770-9188 weekdays 8:00 AM to 4:30 PM EST

Contains no ingredient made from a gluten-containing grain (wheat, barley, or rye)

Lot
 Exp.

300000474
 532000019
 R-00

SIDE 2




Don't modifications to side 2 including artwork:

- 1) Updated adhesive decenter pattern
- 2) Centered drug facts top to bottom



SIDE 1



Don't modifications to frontside artwork (1):

Don't modifications to frontside artwork (2):

↖ .125" or

ibuprofen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55319-396
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
TALC (UNII: 7SEV7J4R1U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POVIDONE (UNII: FZ989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

Product Characteristics

Color	brown (Reddish Brown color Tablets)	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	G2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-396-05	50 in 1 BOTTLE; Type 0: Not a Combination Product	10/29/2018	
2	NDC:55319-396-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	10/29/2018	
3	NDC:55319-396-25	250 in 1 BOTTLE; Type 0: Not a Combination Product	10/29/2018	
4	NDC:55319-396-24	24 in 1 BOTTLE; Type 0: Not a Combination Product	10/31/2018	
5	NDC:55319-396-20	200 in 1 BOTTLE; Type 0: Not a Combination Product	04/01/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079174	10/29/2018	

IBUPROFEN

ibuprofen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55319-397
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POVIDONE (UNII: FZ989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
STARCH, CORN (UNII: O8232NY3SJ)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

Product Characteristics

Color	brown (Reddish Brown tablets)	Score	no score
Shape	OVAL (Caplet shaped Tablets)	Size	14mm
Flavor		Imprint Code	G2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55319-397-05	50 in 1 BOTTLE; Type 0: Not a Combination Product	10/29/2018	
2	NDC:55319-397-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	10/29/2018	
3	NDC:55319-397-25	250 in 1 BOTTLE; Type 0: Not a Combination Product	10/29/2018	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079174	10/29/2018	

Labeler - FAMILY DOLLAR (024472631)

