

## **WALGREENS PAIN RELIEF- benzocaine gel**

### **Walgreens Company**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

### **Drug Facts**

#### **Active Ingredient**

Benzocaine 20%

#### **Purpose**

Oral Pain Reliever

#### **Uses**

- temporary relief of occasional minor irritation, pain and sore mouth

#### **Warnings**

**Allergy alert:** Do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

#### **Do not use**

- more than directed
- for more than 7 days unless told to do so by a dentist or doctor

#### **Stop use and ask a doctor if**

- swelling, rash or fever develops
- irritation, pain, or redness persists or worsens

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### **Directions**

- do not use tube if it is cut prior to opening
- cut open tip of tube on score mark
- use your fingertip or cotton applicator to apply a small pea-size amount of Oral Pain Relief Gel
- apply to affected area up to four times daily or as directed by a dentist or physician
- Adults and children 2 years of age and older: Apply to affected area
- Children under 12 years of age should be supervised in the use of this product
- Children under 2 years of age: Consult a doctor

**Other information**

- store at a controlled room temperature 59°-86°F (15°-30°C)

**Inactive ingredients**

Benzyl Alcohol, Carbomer, D&C YellowNo.10, FD&C Blue No1, FD&C Red No 40, Glycerin, Methylparaben, Propylparaben, Polyethylene Glycol, Propylene Glycol, Sodium Saccharin

**Principal Display Panel - 0.33oz Carton Label**

**Walgreens                      NDC 0363-5061-33**

***Oral Pain Relief***

***Maxium Strength GEL***

***Fast Toothache Pain Reliever***

***20% Benzocaine Topical Gel***

**NET WT. 0.33 OZ (9.35g)**

**Made in the USA**

Walgreens



Compare to the active ingredient in Anbesol® Maximum Strength Gel\*

# Instant Oral Pain Relief

BENZOCAINE 20% / ORAL ANESTHETIC

Maximum Strength

Gel

Helps relieve oral pain

# Instant Oral Pain Relief

BENZOCAINE 20% / ORAL ANESTHETIC

Maximum Strength

Gel

NET WT 0.33 OZ (9.35 g)

\*Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.  
\*\*This product is not manufactured or distributed by © 2018 Pfizer Consumer Healthcare, owner of the registered trademark Anbesol®.

W3ORG0921-F

DISTRIBUTED BY: WALGREEN CO.  
200 WILMOT RD., DEERFIELD, IL 60015  
100% SATISFACTION GUARANTEED  
Walgreens.com 1-800-925-4733  
©2021 Walgreen Co. MADE IN U.S.A.

ITEM 219529 W10439-1221-L



<b>Drug Facts</b>	<b>Active ingredient</b> Benzocaine 20% ..... Oral anesthetic
<b>Purpose</b> temporarily relieves pain associated with the following mouth and gum irritations: • toothache • sore gums • canker sores or ulcers • minor dental procedures • dentures	<b>Uses</b> Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: • pale, gray, or blue colored skin (cyanosis) • headache • rapid heart rate • shortness of breath • dizziness or lightheadedness • fatigue or lack of energy
<b>Warnings</b> • braces • minor dental procedures • dentures and gum irritations: • toothache • sore gums • canker sores or ulcers • minor dental procedures • dentures	<b>Do not use</b> • for teething • in children under 2 years of age occurs, stop use and seek medical help right away. Benzocaine, or other "caine" anesthetics. If a skin reaction of allergy to local anesthetics such as procaine, butacaine, or tetracaine occurs, stop use and seek medical help right away.
<b>When using this product</b> • avoid contact with the eyes • do not exceed recommended dosage • do not use for more than 7 days unless told to do so by a dentist or doctor	<b>Other information</b> • do not use tube if it is cut prior to opening • to open tube, cut open tip of tube on score mark with scissors • store at a controlled room temperature 68°-77°F (20°-25°C) • children under 2 years of age: do not use • children between 2 and 12 years of age: adult supervision should be given in use of this product • adults and children 2 years of age or older: apply to affected area up to 4 times daily or as directed by doctor/dentist
<b>Stop use and ask a doctor if</b> • sore mouth symptoms do not improve in 7 days • irritation, pain, or redness persists or worsens • swelling, rash or fever develops	<b>Directions</b> • adults and children 2 years of age or older: apply to affected area up to 4 times daily or as directed by doctor/dentist • children between 2 and 12 years of age: adult supervision should be given in use of this product • children under 2 years of age: do not use
<b>Keep out of the reach of children.</b> In case of overdose, get medical help or contact a Poison Control Center right away.	<b>Inactive ingredients</b> Benzyl Alcohol, Carbomer, D&C Yellow No. 10, FD&C Blue No. 1, FD&C Red No. 40, Flavor, Glycerin, Methylparaben, Polyethylene Glycol, Propylene Glycol, Sodium Saccharin
<b>Drug Facts (continued)</b>	<b>Questions or comments?</b> 1-800-925-4733



Principal Display Panel - 0.33oz Tube Label

Walgreens

NDC 0363-5061-33

## Oral Pain Relief

### Maximum Strength GEL

### Fast Toothache Pain Reliever

### 20 % Benzocaine Topical Gel

Made in the USA

NET WT. 0.33 OZ (9.35g)



Maximum Strength **Gel**

# Instant Oral Pain Relief

BENZOCAINE 20% / ORAL ANESTHETIC

NET WT 0.33 OZ (9.35 g)

Walgreens Pharmacist Recommended. Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.

DISTRIBUTED BY: WALGREEN CO., 200 WILMOT RD., DEERFIELD, IL 60015 100% SATISFACTION GUARANTEED walgreens.com ©2021 Walgreen Co. MADE IN U.S.A.

**Active ingredient** Benzocaine 20% **Purpose** Oral anesthetic **Uses** temporarily relieves pain associated with the following mouth and gum irritations: • toothache • sore gums • canker sores • braces • minor dental procedures • dentures.

**Warnings** **METHEMOGLOBINEMIA WARNING:** Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: • pale, gray, or blue colored skin (cyanosis) • headache • rapid heart rate • shortness of breath • dizziness or lightheadedness • fatigue or lack of energy. **Allergy alert:** do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics. If a skin reaction occurs, stop use and seek medical help right away. **Do not use** • for teething • in children under 2 years of age. **When using this product** • avoid contact with the eyes • do not exceed recommended dosage • do not use for more than 7 days unless told to do so by a dentist or doctor. **Stop use and ask a doctor if** • sore mouth symptoms do not improve in 7 days • irritation, pain, or redness persists or worsens • swelling, rash or fever develops. **Keep out of the reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. **Directions** • adults and children 2 years of age or older: apply to affected area up to 4 times daily or as directed by doctor/dentist. • children between 2 and 12 years of age: adult supervision should be given in use of this product. • children under 2 years of age: do not use.

**Other information** • do not use tube if it is cut prior to opening • to open tube, cut open tip of tube on score mark with scissors • store at a controlled room temperature 68°-77°F (20°-25°C)

W3ORG0921-F ITEM 219529 W10491-0000-0 **Questions?** 1-800-925-4733

## WALGREENS PAIN RELIEF

benzocaine gel

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-5061
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	200 mg in 1 g

## Inactive Ingredients

Ingredient Name	Strength
<b>BENZYL ALCOHOL</b> (UNII: LKG8494WBH)	
<b>CARBOMER 940</b> (UNII: 4Q93RCW27E)	
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	
<b>POLYETHYLENE GLYCOL 300</b> (UNII: 5655G9Y8AQ)	
<b>POLYETHYLENE GLYCOL 400</b> (UNII: B697894SGQ)	
<b>Saccharin Sodium</b> (UNII: SB8ZUX40TY)	

## Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	SPEARMINT	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-5061-33	1 in 1 CARTON	03/19/2013	
1		9.35 g in 1 TUBE; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part356	03/19/2013	

**Labeler** - Walgreens Company (008965063)

Revised: 12/2021

Walgreens Company