

**FLU HBP MAXIMUM STRENGTH- acetaminophen, chlorpheniramine maleate,
dextromethorphan hbr tablet, film coated
CHAIN DRUG MARKETING ASSOCIATION INC**

Quality Choice 44-738

Active ingredients (in each caplet)

Acetaminophen 325 mg
Chlorpheniramine maleate 2 mg
Dextromethorphan HBr 10 mg

Purpose

Pain reliever/fever reducer
Antihistamine
Cough suppressant

Uses

- temporarily relieves these common cold and flu symptoms:
 - minor aches and pains
 - sore throat
 - sneezing
 - headache
 - runny nose
 - cough
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a

doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- difficulty in urination due to enlargement of the prostate gland
- liver disease
- cough that occurs with too much phlegm (mucus)
- glaucoma
- a breathing problem or persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- avoid alcoholic beverages
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- excitability may occur, especially in children
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
- adults and children 12 years and over: take 2 caplets every 4 hours while symptoms persist. Do not take more than 10 caplets in 24 hours.
- children under 12 years: ask a doctor

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number
- protect from excessive moisture

Inactive ingredients

corn starch, crospovidone, FD&C red #40 aluminum lake, lecithin, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments?

1-800-426-9391

Principal Display Panel

QC®
QUALITY
CHOICE

NDC 63868-567-20

*Compare to the active ingredients in
Coricidin® HBP Maximum Strength
Multi-Symptom Flu

Maximum Strength

Flu HBP

For People with High Blood Pressure

Acetaminophen 325 mg
Chlorpheniramine maleate 2 mg
Dextromethorphan HBr 10 mg

Pain Reliever/Fever Reducer,
Antihistamine & Cough Suppressant

Decongestant-Free

Relieves: Body Aches,
Pains and Headache,

Fever, Cough, Runny Nose,
Sore Throat, and Sneezing

20 Caplets

actual size

**TAMPER EVIDENT: DO NOT USE
IF PACKAGE IS OPENED OR IF
BLISTER UNIT IS TORN, BROKEN
OR SHOWS ANY SIGNS OF TAMPERING**

*This product is not manufactured or distributed by Bayer
HealthCare LLC, owner of the registered trademark
Coricidin® HBP Maximum Strength Multi-Symptom Flu.

50844 REV1019A73809

Distributed by CDMA, Inc.

43157 W Nine Mile

Novi, MI 48375

www.qualitychoice.com

Questions: 800-935-2362

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

SATISFACTION GUARANTEED

100% QC

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

100% SATISFACTION GUARANTEED Distributed by CDMA, Inc. 43157 W Nine Mile Novi, MI 48375 www.qualitychoice.com Questions: 800-935-2362

PARENTS: Learn about teen medicine abuse www.StopMedicineAbuse.org 50844 REV1019A73809

NDC 63868-567-20

QC QUALITY CHOICE

*Compare to the active ingredients in Coricidin® HBP Maximum Strength Multi-Symptom Flu

Maximum Strength Flu HBP

For People with High Blood Pressure

**Acetaminophen 325 mg
Chlorpheniramine maleate 2 mg
Dextromethorphan HBr 10 mg**

**Pain Reliever/Fever Reducer,
Antihistamine & Cough Suppressant
Decongestant-Free**

Relieves: Body Aches, Pains and Headache, Fever, Cough, Runny Nose, Sore Throat, and Sneezing



20 Caplets actual size

*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Coricidin® HBP Maximum Strength Multi-Symptom Flu.

No print/No varnish area
Lot no/Exp date

B-0220-738-09-R
REV1019A73809

6 35515 99792 9

Drug Facts (continued)

microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments? 1-800-426-9391

Drug Facts (continued)

store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F). Protect from excessive moisture. See and flap for expiration date and lot number.

Inactive ingredients: corn starch, croscarmellose, FD&C red #40, aluminum lake, lecithin, magnesium stearate, titanium dioxide

Drug Facts (continued)

Uses

- temporarily relieves these common cold and flu symptoms:
 - minor aches and pains
 - sore throat
 - sneezing
 - headache
 - runny nose
 - cough
 - temporarily reduces fever

Drug Facts (continued)

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash
- skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- difficulty in urination due to enlargement of the prostate gland
- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or persistent or chronic cough as occurs with smoking, asthma, chronic bronchitis, or emphysema

Drug Facts (continued)

Warnings (continued)

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- avoid alcoholic beverages
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- excitability may occur, especially in children
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed
- adults and children 12 years and over: take 2 caplets every 4 hours while symptoms persist. Do not take more than 10 caplets in 24 hours.
- children under 12 years: ask a doctor

Other information

- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN

Quality Choice 44-738

FLU HBP MAXIMUM STRENGTH

acetaminophen, chlorpheniramine maleate, dextromethorphan hbr tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63868-567
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY35J)	

CROSPROVIDONE, UNSPECIFIED (UNII: 2S7830E561)
FD&C RED NO. 40 (UNII: WZB9127XOA)
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)
MAGNESIUM STEARATE (UNII: 70097M6I30)
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)
STEARIC ACID (UNII: 4ELV7Z65AP)
TALC (UNII: 7SEV7J4R1U)
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

Product Characteristics			
Color	red	Score	no score
Shape	OVAL	Size	16mm
Flavor		Imprint Code	44;738
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-567-20	2 in 1 CARTON	03/25/2020	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/25/2020	

Labeler - CHAIN DRUG MARKETING ASSOCIATION INC (011920774)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(63868-567) , pack(63868-567)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(63868-567)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(63868-567)

Revised: 8/2023

CHAIN DRUG MARKETING ASSOCIATION INC