

S.T.37 - hexylresorcinol solution
NUMARK BRANDS, INC

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Drug Facts

Active Ingredient	Purposes
Hexylresorinol 0.1%	Oral Pain Reliever
	First aid antiseptic

Uses temporarily relieves:

- minor irritations of the mouth
- minor pain in the mouth
- sore mouth
- sore throat
- first aid to help prevent skin infection in minor cuts, scrapes, and burns

Warnings

For external use only

Sore throat warning if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly.

Do not use

- In the eyes
- over large areas of the body

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

When using this product

- **do not exceed recommended dosage**

Stop use and ask a doctor if

- condition persists
- condition worsens
- you need to use product longer than 1 week

Stop use and ask a doctor or dentist if

- sore mouth symptoms do not improve in 7 days
- irritation, pain, and redness persist or worsen

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions (as a First Aid antiseptic)

- use full strength
- clean affected area
- apply a small amount of product on the area
- use 1-3 times daily
- may be covered with a sterile bandage
- if bandaged let dry first

Directions (as an Oral pain reliever)

- apply to affected area
- gargle, swish around, or allow to remain in place for at least one minute and then spit out

adults and children 2 years old and over	use up to 4 times daily or as directed by a dentist or doctor
children under 2 years old	ask a dentist or doctor

Other Information

- **store at controlled room temperature 59° - 86°F (15° - 30°C)**

Inactive Ingredients

citric acid, edetate disodium, glycerin, propylene glycol, sodium bisulfite, sodium citrate, purified water.

Questions call 1-800-214-2379

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

8 oz SIZE

2.5 more ounces at the same low price

S.T.37®

**FIRST AID ANTISEPTIC/ORAL PAIN RELIEVER
SOLUTION**

FOR BURNS, CUTS, ABRASIONS & HYGIENIC CARE OF THE MOUTH.

Do not use if foil seal under cap is broken or missing at time of purchase

8 FL OZ (236.5 mL)

WON'T STING — WON'T STAIN

S.T.37 kills common pathogenic bacteria quickly — on contact!

Laboratory tests have established S.T.37 antiseptic solution as a general antiseptic for household use.

S.T.37 is pleasant to use because it produces little or no stinging or burning. In fact, it soothes irritated areas.

S.T.37 does not stain the skin or leave an order.

Everyday Uses for S.T.37

Cuts & Abrasions: Apply full strength S.T.37 into the cut or over the injured area. Since it is germicidal for the common pathogenic bacteria, S.T.37 may minimize the development of infection. Excellent for cleansing cuts and scrapes.

Burns: Apply full strength S.T.37 for treatment of minor burns. S.T.37 combats the cause of infections.

Hygienic Care of Mouth: Use undiluted and gargle or rinse to help keep mouth clean and fresh. S.T.37 temporarily relieves minor pain and protects irritated areas of a sore mouth or throat.

Distributed by Numark Brands, Inc.
27070 Miles Rd, Suite A, Solon Ohio 44139
www.numarkbrands.com

B10082
Rev. 3/18

NOITNOS

FIRST AID ANTISEPTIC/ORAL PAIN RELIEVER

S.T.37

WON'T STING—WON'T STAIN

S.T.37

FIRST AID ANTISEPTIC/ORAL PAIN RELIEVER
SOLUTION

Drug Facts

Active ingredient Hexyresorcinol 0.1%
Purposes Oral pain reliever
First aid antiseptic

Uses temporarily relieves:
■ minor irritations of the mouth ■ sore mouth
■ minor pain in the mouth ■ sore throat
■ first aid to help prevent skin infection in minor cuts, scrapes, and burns

Warnings

For external use only

Sore throat warning If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly.

Do not use

■ in the eyes ■ over large areas of the body

Ask a doctor before use if you have

■ deep or puncture wounds ■ animal bites
■ serious burns

When using this product ■ do not exceed recommended dosage

Stop use and ask a doctor if

■ condition persists ■ condition worsens
■ you need to use product longer than 1 week

Stop use and ask a doctor or dentist if

■ sore mouth symptoms do not improve in 7 days
■ irritation, pain, and redness persist or worsen

If pregnant or breast feeding, ask a health professional before use.

WON'T STING—WON'T STAIN

S.T.37

FIRST AID ANTISEPTIC/ORAL PAIN RELIEVER
SOLUTION

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions (as a First Aid antiseptic)
■ use full strength ■ clean affected area
■ apply a small amount of product on the area
■ use 1-3 times daily
■ may be covered with a sterile bandage
■ if bandaged, let dry first

Directions (as an Oral pain reliever)
■ apply to affected area
■ gargle, swish around, or allow to remain in place for at least one minute and then spit out

adults and children
2 years old and over

use up to 4 times
daily or as directed
by a dentist or doctor
children under 2
years old ask a dentist or
doctor

Other information ■ store at controlled room temperature 59°-86°F (15°-30°C)

Inactive ingredients citric acid, edetate disodium, glycerin, propylene glycol, sodium bisulfite, sodium citrate, purified water.

Questions call 1-800-214-2379

Distributed by Numark Brands, Inc.
27070 Miles Rd, Suite A, Solon Ohio 44139
www.numarkbrands.com

8 oz SIZE
2.5 more ounces
at the same low price

S.T.37

FIRST AID ANTISEPTIC/ORAL PAIN RELIEVER
SOLUTION

FOR BURNS,
CUTS, ABRASIONS
& HYGIENIC CARE
OF THE MOUTH.

Do not use if foil seal under cap is
broken or missing at time of purchase

8 FL OZ (236.5 mL)

WON'T STING—WON'T STAIN

S.T.37

FIRST AID ANTISEPTIC/ORAL PAIN RELIEVER
SOLUTION

S.T.37 kills common pathogenic bacteria quickly —on contact!

Laboratory tests have established S.T.37 antiseptic solution as a general antiseptic for household use.

S.T.37 is pleasant to use because it produces little or no stinging or burning. In fact, it soothes irritated areas.

S.T.37 does not stain the skin or leave an odor.

Everyday Uses for S.T.37

Cuts & Abrasions: Apply full strength S.T.37 into the cut or over the injured area. Since it is germicidal for the common pathogenic bacteria, S.T.37 may minimize the development of infection. Excellent for cleansing cuts and scrapes.

Burns: Apply full strength S.T.37 for treatment of minor burns. S.T.37 combats the cause of infections.

Hygienic Care of Mouth: Use undiluted and gargle or rinse to help keep mouth clean and fresh. S.T.37 temporarily relieves minor pain and protects irritated areas of a sore mouth or throat.



B10082
Rev. 3/18

S.T.37®
FIRST AID ANTISEPTIC/ORAL PAIN RELIEVER
SOLUTION

FOR BURNS, CUTS, ABRASIONS & HYGIENIC CARE OF THE MOUTH

8 FL OZ (236.5 mL)

Do not use if foil seal under cap is broken or missing at time of purchase

S.T.37 kills common pathogenic bacteria quickly - on contact!

Laboratory tests have established S.T.37 antiseptic solution as a general antiseptic for household use.

S.T.37 is pleasant to use because it produces little or no stinging or burning. In fact, it soothes irritated areas.

S.T.37 does not stain the skin or leave an odor.

Everyday Uses for S.T.37

Cuts & Abrasions: Apply full strength S.T.37 into the cut or over the injured area. Since it is germicidal for the common pathogenic bacteria, S.T.37 may minimize the development of infection. Excellent for cleansing cuts and scrapes.

Burns: Apply full strength S.T.37 for treatment of minor burns. S.T.37 combats the cause of infections.

Hygienic Care of Mouth: Use undiluted and gargle or rinse to help keep mouth clean and fresh. S.T.37 temporarily relieves minor pain and protects irritated areas of a sore mouth or throat.

Distributed by Numark Brands, Inc.

27070 Miles Rd, Suite A, Solon Ohio 44139 1-800-214-2379

www.numarkbrands.com Rev. 3/18 J12008

REFER TO OUTER CARTON FOR FULL INFORMATION

USES: temporarily relieves: • minor irritations of the mouth • minor pain in the mouth • sore mouth • sore throat • first aid to help prevent skin infection in minor cuts, scrapes, and burns.

DIRECTIONS: (as a First Aid antiseptic) • use full strength • clean affected area • apply a small amount of product on the area • use 1-3 times daily • may be covered with a sterile bandage • if bandaged let dry first.

DIRECTIONS: (as an Oral pain reliever) • apply to affected area • gargle, swish around, or allow to remain in place for at least one minute and then spit out. For adults and children 2 years old and over: use up to 4 times daily or as directed by a dentist or doctor. For children under 2 years old: ask a dentist or doctor.

WARNINGS: For external use only. **Sore throat warning** if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. **Do not use** • in the eyes • over large areas of the body. **Ask a doctor before use if you have** • deep or puncture wounds • animal bites • serious burns. **When using this product** • do not exceed recommended dosage. **Stop use and ask a doctor if** • condition persists • condition worsens • you need to use product longer than 1 week. **Stop use and ask a doctor or dentist if** • sore mouth symptoms do not improve in 7 days • irritation, pain, and redness persist or worsen. **If pregnant or breast feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

ACTIVE INGREDIENT: Hexylresorcinol 0.1%.

INACTIVE INGREDIENTS: Citric acid, Edetate disodium, Glycerin, Propylene glycol, Sodium bisulfite, Sodium citrate, Purified water.



ST.37
FIRST AID ANTISEPTIC/ORAL PAIN RELIEVER
SOLUTION

**FOR BURNS,
CUTS, ABRASIONS
& HYGIENIC CARE
OF THE MOUTH.**

8 FL OZ (236.5 mL)

Do not use if foil seal under cap is broken or missing at time of purchase

S.T.37 kills common pathogenic bacteria quickly - on contact! Laboratory tests have established S.T.37 antiseptic solution as a general antiseptic for household use.

S.T.37 is pleasant to use because it produces little or no stinging or burning. In fact, it soothes irritated areas.

S.T.37 does not stain the skin or leave an odor.

Everyday Uses for S.T.37

Cuts & Abrasions: Apply full strength S.T.37 into the cut or over the injured area. Since it is germicidal for the common pathogenic bacteria, S.T.37 may minimize the development of infection. Excellent for cleansing cuts and scrapes.

Burns: Apply full strength S.T.37 for treatment of minor burns. S.T.37 combats the cause of infections.

Hygienic Care of Mouth: Use undiluted and gargle or rinse to help keep mouth clean and fresh. S.T.37 temporarily relieves minor pain and protects irritated areas of a sore mouth or throat.

Distributed by Numark Brands, Inc.
27070 Miles Rd, Suite A, Solon Ohio 44139 1-800-214-2379
www.numarkbrands.com Rev. 3/18 J12008

S.T.37

hexylresorcinol solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69846-860
Route of Administration	ORAL, TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Hexylresorcinol (UNII: R9QTB5E82N) (Hexylresorcinol - UNII:R9QTB5E82N)	Hexylresorcinol	1.1 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
Citric Acid Monohydrate (UNII: 2968PHW8QP)	
Edetate Disodium (UNII: 7FLD91C86K)	
Glycerin (UNII: PDC6A3C0OX)	
Propylene Glycol (UNII: 6DC9Q167V3)	
Sodium Bisulfite (UNII: TZX5469Z6I)	
Sodium Citrate (UNII: 1Q73Q2JULR)	
Water (UNII: 059QF0KOOR)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69846-860-00	1 in 1 AMPULE	11/20/1929	
1		236.5 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:69846-860-01	1 in 1 CARTON	11/20/1929	
2		473 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		11/20/1929	

Labeler - NUMARK BRANDS, INC (080184668)