

## **EXTRA STRENGTH HEADACHE RELIEF- acetaminophen, aspirin, caffeine tablet WALGREENS**

### **EXTRA STRENGTH HEADACHE RELIEF**

#### ***Drug Facts***

<b><i>Active ingredient (in each caplet)</i></b>	<b><i>Purpose</i></b>
Acetaminophen 250 mg	Pain reliever
Aspirin 250 mg (NSAID) *	Pain reliever
Caffeine 65 mg	Pain reliever aid

\* nonsteroidal anti-inflammatory drug

#### **Uses**

temporarily relieves minor aches and pains due to:

- headache
- a cold
- arthritis
- muscular aches
- sinusitis
- toothache
- premenstrual & menstrual cramps

#### **Warnings**

##### **Reye's syndrome**

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

##### **Allergy alert**

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening

- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

### **Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

### **Stomach bleeding warning**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Caffeine warning**

The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

### **Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

### **Ask doctor before use if**

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you are taking a diuretic
- you have asthma

### **Ask a doctor or pharmacist before use if you are taking**

- any other drug containing an NSAID (prescription or nonprescription)
- a prescription drug for diabetes, gout, or arthritis
- any other drug, or are under a doctor's care for any serious condition
- a blood thinning (anticoagulant) or steroid drug

**Stop use and ask doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- painful area is red or swollen
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- any new symptoms appear

These could be signs of a serious condition

**If pregnant or breast-feeding** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children****Overdose Warning**

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. 1(800) 222-1222

**Directions**

- **do not use more than directed (see Overdose Warning)**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours
- children under 12 years: ask a doctor

**Other information**

- **Each caplet contains:** Magnesium 0.58 mg
- store at controlled room temperature 20° - 25° C (68° - 77° F)
- read all product information before using. Keep this box for important information

**Inactive ingredients**

colloidal silicon dioxide, hypromellose, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, stearic acid, talc

**Questions or comments**

1-888-333-9792

PRINCIPAL DISPLAY PANEL

COMPARE TO THE ACTIVE

INGREDIENTS OF EXCEDRIN®

EXTRA STRENGTH\*

EXTRA STRENGHT

HEADACHE RELIEF

ACETAMINOPHE, ASPIRIN (NSAID) AND CAFFEINE

250 Caplets

VALUE SIZE

NDC 0363-7214-25

Walgreens

Compare to the active ingredients in Excedrin® Extra Strength†

Headache Relief

ACETAMINOPHEN / ASPIRIN (NSAID) / PAIN RELIEVER / CAFFEINE / PAIN RELIEVER AID

EXTRA STRENGTH

CAPLETS

250 CAPLETS

ACTUAL SIZE

TAMPER EVIDENT: DO NOT USE IF SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredients (in each caplet)

Acetaminophen 250 mg.....Pain reliever

Aspirin 250 mg (NSAID).....Pain reliever

Caffeine 65 mg.....Pain reliever aid

\*nonsteroidal anti-inflammatory drug

Uses

temporarily relieves minor aches and pains due to:

headache    a cold    arthritis

toothache    premenstrual & menstrual cramps

Warnings

Reye's syndrome:

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert:

Acetaminophen may cause severe skin reactions. Symptoms may include:    skin reddening    blisters

If a skin reaction occurs, stop use and seek medical help right away.

Allergy alert:

Aspirin may cause a severe allergic reaction which may include:    hives    facial swelling    asthma (wheezing)    stomach pain

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if you take    more than 8 caplets in 24 hours, which is the maximum daily amount    with other drugs containing acetaminophen

Stomach bleeding warning:

This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you    are age 60 or older    have had stomach ulcers or bleeding problems    take a blood thinning (anticoagulant) or steroid drug    take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Drug Facts (continued under label)

DISTRIBUTED BY: WALGREEN CO.  
200 W. MORT RD., DEERFIELD, IL 60015  
100% SATISFACTION GUARANTEED  
walgreens.com ©2020 Walgreen Co.

ITEM 342812 W10076-0221-F

3 11917 100358 0

ORG120-F Made in India

LOT:    EXP:

PEEL HERE

Drug Facts (continued)

Do not use

if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer

with any other drug containing acetaminophen (prescription or nonprescription).

If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

you have liver disease

stomach bleeding warning applies to you

you have a history of stomach problems, such as heartburn

you have high blood pressure, heart disease, liver cirrhosis or kidney disease

you are taking a diuretic

you have asthma

Ask a doctor or pharmacist before use if you are

taking a prescription drug for diabetes, gout, or arthritis

taking any other drug or are under a doctor's care for any serious condition

Stop use and ask a doctor if

an allergic reaction occurs. Seek medical help right away.

you experience any of the following signs of stomach bleeding

feel faint    vomit blood    have bloody or black stools

have stomach pain that does not get better

ringing in the ears or loss of hearing occurs

painful area is red or swollen

pain gets worse or lasts for more than 10 days

fever gets worse or lasts for more than 3 days

any new symptoms appear

These could be signs of a serious condition.

If pregnant or breast-feeding

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not use more than directed

drink a full glass of water with each dose

adults and children 12 years and over: take 2 caplets every 6 hours; not more than 8 caplets in 24 hours

children under 12 years: ask a doctor

Other information

store between 20°-25°C (68°-77°F)

Inactive ingredients

colloidal silicon dioxide, hypromellose, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, stearic acid, talc

Questions or comments?

Call 1-888-333-9792

EXTRA STRENGTH HEADACHE RELIEF

acetaminophen, aspirin, caffeine tablet

Product Information

Product Type		HUMAN OTC DRUG	Item Code (Source)	NDC:0363-7214
Route of Administration		ORAL		
Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)			ACETAMINOPHEN	250 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)			ASPIRIN	250 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)			CAFFEINE	65 mg
Inactive Ingredients				
Ingredient Name				Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
POVIDONE K30 (UNII: U725QWY32X)				
STARCH, CORN (UNII: O8232NY3SJ)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
TALC (UNII: 7SEV7J4R1U)				
Product Characteristics				
Color	white		Score	no score
Shape	CAPSULE (capsule shaped, biconvex)		Size	18mm
Flavor			Imprint Code	S53
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-7214-25	250 in 1 BOTTLE; Type 0: Not a Combination Product	11/17/2020	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M013	11/17/2020	

**Labeler -** WALGREENS (008965063)