

# TAZVERIK- tazemetostat tablet, film coated

## Epizyme, Inc.

### HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use TAZVERIK® safely and effectively. See full prescribing information for TAZVERIK.

TAZVERIK (tazemetostat) tablets, for oral use

Initial U.S. Approval: 2020

### RECENT MAJOR CHANGES

Warnings and Precautions, Secondary Malignancies (5.1)

11/2023

### INDICATIONS AND USAGE

TAZVERIK is a methyltransferase inhibitor indicated for the treatment of:

- Adults and pediatric patients aged 16 years and older with metastatic or locally advanced epithelioid sarcoma not eligible for complete resection. (1.1)
- Adult patients with relapsed or refractory follicular lymphoma whose tumors are positive for an EZH2 mutation as detected by an FDA-approved test and who have received at least 2 prior systemic therapies. (1.2)
- Adult patients with relapsed or refractory follicular lymphoma who have no satisfactory alternative treatment options. (1.2)

These indications are approved under accelerated approval based on overall response rate and duration of response. Continued approval for these indications may be contingent upon verification and description of clinical benefit in a confirmatory trial(s).

### DOSAGE AND ADMINISTRATION

- Recommended dosage is 800 mg taken orally twice daily with or without food. (2.2)

### DOSAGE FORMS AND STRENGTHS

Tablets: 200 mg (3)

### CONTRAINDICATIONS

None. (4)

### WARNINGS AND PRECAUTIONS

- Secondary Malignancies:** TAZVERIK increases the risk of developing secondary malignancies, including T-cell lymphoblastic lymphoma, myelodysplastic syndrome, acute myeloid leukemia, and B-cell acute lymphoblastic leukemia. Monitor patients long-term for the development of secondary malignancies. (5.1)
- Embryo-Fetal Toxicity:** Can cause fetal harm. Advise patients of potential risk to a fetus and to use effective non-hormonal contraception. (5.2)

### ADVERSE REACTIONS

- The most common ( $\geq 20\%$ ) adverse reactions in patients with epithelioid sarcoma are pain, fatigue, nausea, decreased appetite, vomiting, and constipation. (6.1)
- The most common ( $\geq 20\%$ ) adverse reactions in patients with follicular lymphoma are fatigue, upper respiratory tract infection, musculoskeletal pain, nausea, and abdominal pain. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Epizyme at 855-463-5127 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

### DRUG INTERACTIONS

- Strong and Moderate Cytochrome P450 (CYP)3A Inhibitors:** Avoid coadministration of strong and moderate CYP3A inhibitors with TAZVERIK. Reduce the dose of TAZVERIK if coadministration of moderate CYP3A inhibitors cannot be avoided. (2.3, 7.1)
- Strong and Moderate CYP3A Inducers:** Avoid coadministration with TAZVERIK. (7.1)

### USE IN SPECIFIC POPULATIONS

- Lactation:** Advise not to breastfeed. (8.2)

---

## **FULL PRESCRIBING INFORMATION: CONTENTS\***

### **1 INDICATIONS AND USAGE**

- 1.1 Epithelioid Sarcoma
- 1.2 Relapsed or Refractory Follicular Lymphoma

### **2 DOSAGE AND ADMINISTRATION**

- 2.1 Patient Selection
- 2.2 Recommended Dosage
- 2.3 Dosage Modifications for Adverse Reactions
- 2.4 Dosage Modifications for Drug Interactions

### **3 DOSAGE FORMS AND STRENGTHS**

### **4 CONTRAINDICATIONS**

### **5 WARNINGS AND PRECAUTIONS**

- 5.1 Secondary Malignancies
- 5.2 Embryo-Fetal Toxicity

### **6 ADVERSE REACTIONS**

- 6.1 Clinical Trial Experience

### **7 DRUG INTERACTIONS**

- 7.1 Effect of Other Drugs on TAZVERIK
- 7.2 Effect of TAZVERIK on Other Drugs

### **8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy
- 8.2 Lactation
- 8.3 Females and Males of Reproductive Potential
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment
- 8.7 Hepatic Impairment

### **11 DESCRIPTION**

### **12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

### **13 NONCLINICAL TOXICOLOGY**

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

### **14 CLINICAL STUDIES**

- 14.1 Epithelioid Sarcoma
- 14.2 Relapsed or Refractory Follicular Lymphoma

### **15 REFERENCES**

### **16 HOW SUPPLIED/STORAGE AND HANDLING**

### **17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

---

## **FULL PRESCRIBING INFORMATION**

### **1 INDICATIONS AND USAGE**

#### **1.1 Epithelioid Sarcoma**

TAZVERIK is indicated for the treatment of adults and pediatric patients aged 16 years and older with metastatic or locally advanced epithelioid sarcoma not eligible for complete resection.

This indication is approved under accelerated approval based on overall response rate and duration of response [see *Clinical Studies (14.1)*]. Continued approval for this indication may be contingent upon verification and description of clinical benefit in a confirmatory trial(s).

#### **1.2 Relapsed or Refractory Follicular Lymphoma**

- TAZVERIK is indicated for the treatment of adult patients with relapsed or refractory (R/R) follicular lymphoma (FL) whose tumors are positive for an EZH2 mutation as detected by an FDA-approved test and who have received at least 2 prior systemic therapies.
- TAZVERIK is indicated for the treatment of adult patients with R/R FL who have no satisfactory alternative treatment options.

These indications are approved under accelerated approval based on overall response rate and duration of response [see *Clinical Studies (14.2)*]. Continued approval for these indications may be contingent upon verification and description of clinical benefit in a confirmatory trial(s).

### **2 DOSAGE AND ADMINISTRATION**

#### **2.1 Patient Selection**

Select patients with R/R FL for treatment with TAZVERIK based on the presence of EZH2 mutation of codons Y646, A682, or A692 in tumor specimens [see *Clinical Studies (14.2)*]. Information on FDA-approved tests for the detection of EZH2 mutation in relapsed or refractory follicular lymphoma is available at: <http://www.fda.gov/CompanionDiagnostics>.

#### **2.2 Recommended Dosage**

The recommended dosage of TAZVERIK is 800 mg orally twice daily with or without food until disease progression or unacceptable toxicity.

Swallow tablets whole. Do not cut, crush, or chew tablets.

Do not take an additional dose if a dose is missed or vomiting occurs after TAZVERIK, but continue with the next scheduled dose.

#### **2.3 Dosage Modifications for Adverse Reactions**

Table 1 summarizes the recommended dose reductions, and Table 2 summarizes the recommended dosage modifications of TAZVERIK for adverse reactions.

**Table 1. Recommended Dose Reductions of TAZVERIK for Adverse Reactions**

Dose Reduction	Dosage
First	600 mg orally twice daily
Second	400 mg orally twice daily*

\*Permanently discontinue TAZVERIK in patients who are unable to tolerate 400 mg orally twice daily.

**Table 2. Recommended Dosage Modifications of TAZVERIK for Adverse Reactions**

Adverse Reaction	Severity	Dosage Modification
Neutropenia [see Adverse Reactions (6.1)]	Neutrophil count less than $1 \times 10^9/L$	<ul style="list-style-type: none"> <li>Withhold until neutrophil count is greater than or equal to <math>1 \times 10^9/L</math> or baseline.</li> <li>For first occurrence, resume at same dose.</li> <li>For second and third occurrence, resume at reduced dose.</li> <li>Permanently discontinue after fourth occurrence.</li> </ul>
Thrombocytopenia [see Adverse Reactions (6.1)]	Platelet count less than $50 \times 10^9/L$	<ul style="list-style-type: none"> <li>Withhold until platelet count is greater than or equal to <math>75 \times 10^9/L</math> or baseline.</li> <li>For first and second occurrence, resume at reduced dose.</li> <li>Permanently discontinue after third occurrence.</li> </ul>
Anemia [see Adverse Reactions (6.1)]	Hemoglobin less than 8 g/dL	<ul style="list-style-type: none"> <li>Withhold until improvement to at least Grade 1 or baseline, then resume at same or reduced dose.</li> </ul>
Other adverse reactions [see Adverse Reactions (6.1)]	Grade 3	<ul style="list-style-type: none"> <li>Withhold until improvement to at least Grade 1 or baseline.</li> <li>For first and second occurrence, resume at reduced dose.</li> <li>Permanently discontinue after third occurrence.</li> </ul>
	Grade 4	<ul style="list-style-type: none"> <li>Withhold until improvement to at least Grade 1 or baseline.</li> <li>For first occurrence, resume at reduced dose.</li> <li>Permanently discontinue after second occurrence.</li> </ul>

## 2.4 Dosage Modifications for Drug Interactions

### Strong and Moderate CYP3A Inhibitors

Avoid coadministration of TAZVERIK with strong or moderate CYP3A inhibitors. If coadministration with a moderate CYP3A inhibitor cannot be avoided, reduce the TAZVERIK dose as shown in Table 3 below. After discontinuation of the moderate CYP3A inhibitor for 3 elimination half-lives, resume the TAZVERIK dose that was taken prior to initiating the inhibitor [see *Drug Interactions (7.1)*, *Clinical Pharmacology (12.3)*].

**Table 3. Recommended Dose Reductions of TAZVERIK for Moderate CYP3A Inhibitors**

<b>Current Dosage</b>	<b>Adjusted Dosage</b>
800 mg orally twice daily	400 mg orally twice daily
600 mg orally twice daily	400 mg for first dose and 200 mg for second dose
400 mg orally twice daily	200 mg orally twice daily

## 3 DOSAGE FORMS AND STRENGTHS

Tablets: 200 mg film-coated, red, round, biconvex shape and debossed with "EZM 200" on one side and plain on the other.

## 4 CONTRAINDICATIONS

None.

## 5 WARNINGS AND PRECAUTIONS

### 5.1 Secondary Malignancies

The risk of developing secondary malignancies is increased following treatment with TAZVERIK. Across clinical trials of 758 adults who received TAZVERIK 800 mg twice daily as monotherapy, myelodysplastic syndrome (MDS), acute myeloid leukemia (AML), or B-cell acute lymphoblastic leukemia (B-ALL) occurred in 1.7% of patients. One pediatric patient developed T-cell lymphoblastic lymphoma (T-LBL). Monitor patients long-term for the development of secondary malignancies.

### 5.2 Embryo-Fetal Toxicity

Based on findings from animal studies and its mechanism of action, TAZVERIK can cause fetal harm when administered to pregnant women. There are no available data on TAZVERIK use in pregnant women to inform the drug-associated risk. Administration of tazemetostat to pregnant rats and rabbits during organogenesis resulted in dose-dependent increases in skeletal developmental abnormalities in both species beginning at maternal exposures approximately 1.5 times the adult human exposure (area under the plasma concentration time curve [AUC<sub>0-45h</sub>]) at the 800 mg twice daily dose.

Advise pregnant women of the potential risk to a fetus. Advise females of reproductive potential to use effective contraception during treatment with TAZVERIK and for 6 months after the final dose. Advise males with female partners of reproductive potential to use effective contraception during treatment with TAZVERIK and for 3 months after the final dose [see *Use in Specific Populations* (8.1, 8.3)].

6 ADVERSE REACTIONS

The following clinically significant adverse reactions are described elsewhere in labeling:

- Secondary Malignancies [see *Warnings and Precautions* (5.1)].

6.1 Clinical Trial Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared with rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Epithelioid Sarcoma

The safety of TAZVERIK was evaluated in a cohort (Cohort 5) of Study EZH-202 that enrolled patients with epithelioid sarcoma [see *Clinical Studies* (14.1)]. Patients received TAZVERIK 800 mg orally twice daily (n=62). Among patients who received TAZVERIK, 44% were exposed for 6 months or longer and 24% were exposed for greater than one year.

Serious adverse reactions occurred in 37% of patients who received TAZVERIK. Serious adverse reactions in ≥3% of patients who received TAZVERIK were hemorrhage, pleural effusion, skin infection, dyspnea, pain, and respiratory distress.

One patient (2%) permanently discontinued TAZVERIK due to an adverse reaction of altered mood.

Dosage interruptions due to an adverse reaction occurred in 34% of patients who received TAZVERIK. The most frequent adverse reactions requiring dosage interruptions in ≥3% were hemorrhage, increased alanine aminotransferase (ALT), and increased aspartate aminotransferase (AST).

Dose reduction due to an adverse reaction occurred in one (2%) patient who received TAZVERIK; the dose was reduced in this patient for decreased appetite.

The most common adverse reactions (≥20%) were pain, fatigue, nausea, decreased appetite, vomiting, and constipation.

Table 4 presents adverse reactions in patients with epithelioid sarcoma in Cohort 5 of Study EZH-202.

**Table 4. Adverse Reactions (≥10%) in Patients with Epithelioid Sarcoma Who Received TAZVERIK in Cohort 5 of Study EZH-202**

Adverse Reaction	TAZVERIK N=62	
	All Grades (%)	Grade 3 or 4 (%)
General		

Pain <sup>a</sup>	52	7
Fatigue <sup>b</sup>	47	1.6
Gastrointestinal		
Nausea	36	0
Vomiting	24	0
Constipation	21	0
Diarrhea	16	0
Abdominal pain <sup>c</sup>	13	1.6
Metabolism and nutrition		
Decreased appetite	26	4.8
Respiratory, thoracic and mediastinal		
Cough	18	0
Dyspnea <sup>d</sup>	16	4.8
Vascular		
Hemorrhage <sup>e</sup>	18	4.8
Nervous system		
Headache	18	0
Investigations		
Weight decreased	16	7

a Includes tumor pain, pain in extremity, non-cardiac chest pain, flank pain, back pain, arthralgia, bone pain, cancer pain, musculoskeletal pain, myalgia, neck pain

b Includes fatigue and asthenia

c Includes abdominal pain, gastrointestinal pain, abdominal pain lower

d Includes dyspnea and dyspnea exertional

e Includes wound hemorrhage, rectal hemorrhage, pulmonary hemorrhage, hemorrhage intracranial, cerebral hemorrhage, hemoptysis

Table 5 summarizes select laboratory abnormalities in patients with epithelioid sarcoma in Cohort 5 of Study EZH-202.

**Table 5. Select Laboratory Abnormalities (≥ 10%) Worsening from Baseline in Patients with Epithelioid Sarcoma Who Received TAZVERIK in Cohort 5 of Study EZH-202**

Laboratory Abnormality	TAZVERIK*	
	All Grades (%)	Grade 3 or 4 (%)
Hematology		
Decreased hemoglobin	49	15
Decreased lymphocytes	36	13
Decreased white blood cell count	19	0
Chemistry		
Increased triglycerides	36	3.3
Increased glucose	33	1.6
Decreased sodium	30	1.7
Decreased phosphate	28	1.7
Decreased albumin	23	0
Increased alkaline phosphatase	23	1.7

Decreased potassium	20	1.7
Increased aspartate aminotransferase	18	3.5
Decreased calcium	16	0
Decreased glucose	16	0
Increased partial thromboplastin time	15	5
Increased alanine aminotransferase	14	3.4
Increased creatinine	12	0
Increased potassium	12	0

\*The denominator used to calculate the rate varied from 39 to 61 based on the number of patients with a baseline value and at least one post-treatment value.

### Relapsed or Refractory Follicular Lymphoma

The safety of TAZVERIK was evaluated in two cohorts (Cohorts 4 and 5) of Study E7438-G000-101 that enrolled patients with relapsed or refractory follicular lymphoma [see *Clinical Studies (14.2)*]. Patients received TAZVERIK 800 mg orally twice daily (n=99). Among patients who received TAZVERIK, 68% were exposed for 6 months or longer, 39% were exposed for 12 months or longer, and 21% were exposed for 18 months or longer.

The median age was 62 years (range 36 to 87 years), 54% were male, and 95% had an Eastern Cooperative Oncology Group (ECOG) performance status of 0-1. The median number of prior therapies was 3 (range 1 to 11). Patients were required have a creatinine clearance  $\geq 40$  mL/min per the Cockcroft and Gault formula.

Serious adverse reactions occurred in 30% of patients who received TAZVERIK. Serious adverse reactions in  $\geq 2\%$  of patients who received TAZVERIK were general physical health deterioration, abdominal pain, pneumonia, sepsis, and anemia.

Permanent discontinuation due to an adverse reaction occurred in 8% of patients who received TAZVERIK. Adverse reaction resulting in permanent discontinuation in  $\geq 2\%$  of patients was second primary malignancy.

Dosage interruptions due to an adverse reaction occurred in 28% of patients who received TAZVERIK. Adverse reactions requiring dosage interruptions in  $\geq 3\%$  of patients were thrombocytopenia and fatigue.

Dose reduction due to an adverse reaction occurred in 9% of patients who received TAZVERIK.

The most common adverse reactions ( $\geq 20\%$ ) were fatigue, upper respiratory tract infection, musculoskeletal pain, nausea, and abdominal pain.

Table 6 presents adverse reactions in patients with relapsed or refractory follicular lymphoma in Cohorts 4 and 5 of Study E7438-G000-101.

**Table 6. Adverse Reactions ( $\geq 10\%$ ) in Patients with Relapsed or Refractory Follicular Lymphoma Who Received TAZVERIK in Cohorts 4 and 5 of Study E7438-G000-101**

Adverse Reaction	TAZVERIK N=99	
	All Grades (%)	Grade 3 or 4 (%)



General		
Fatigue <sup>a</sup>	36	5
Pyrexia	10	0
Infections		
Upper respiratory tract infection <sup>b</sup>	30	0
Lower respiratory tract infection <sup>c</sup>	17	0
Urinary tract infection <sup>d</sup>	11	2
Gastrointestinal		
Nausea	24	1
Abdominal pain <sup>e</sup>	20	3
Diarrhea	18	0
Vomiting	12	1
Musculoskeletal and connective tissue		
Musculoskeletal pain <sup>f</sup>	22	1
Skin and subcutaneous tissue		
Alopecia	17	0
Rash <sup>g</sup>	15	0
Respiratory and mediastinal system		
Cough <sup>h</sup>	17	0
Nervous system		
Headache <sup>i</sup>	13	0

a Includes fatigue and asthenia

b Includes laryngitis, nasopharyngitis, pharyngitis, rhinitis, sinusitis, upper respiratory tract infection, viral upper respiratory tract infection

c Includes bronchitis, lower respiratory tract infection, tracheobronchitis

d Includes cystitis, urinary tract infection, urinary tract infection staphylococcal

e Includes abdominal discomfort, abdominal pain, abdominal pain lower, abdominal pain upper

f Includes back pain, limb discomfort, musculoskeletal chest pain, musculoskeletal discomfort, musculoskeletal pain, myalgia, neck pain, non-cardiac chest pain, pain in extremity, pain in jaw, spinal pain

g Includes erythema, rash, rash erythematous, rash generalized, rash maculo-papular, rash pruritic, rash pustular, skin exfoliation

h Includes cough and productive cough

i Includes headache, migraine, sinus headache

Clinically relevant adverse reactions occurring in <10% of patients who received TAZVERIK included:

- Infection: sepsis (2%), pneumonia (2%), and herpes zoster (2%)

Table 7 summarizes select laboratory abnormalities in patients with follicular lymphoma in Cohorts 4 and 5 of Study E7438-G000-101.

**Table 7. Select Laboratory Abnormalities (≥10%) Worsening from Baseline in Patients with Relapsed or Refractory Follicular Lymphoma Who Received TAZVERIK in Cohorts 4 and 5 of Study E7438-G000-101**

Laboratory Abnormality	TAZVERIK*	
	All Grades (%)	Grade 3 or 4 (%)

<b>Hematology</b>		
Decreased lymphocytes	57	18
Decreased hemoglobin	50	8
Decreased platelets	50	7
Decreased white blood cells	41	9
Decreased neutrophils	20	7
<b>Chemistry</b>		
Increased glucose	53	10
Increased aspartate aminotransferase	24	0
Increased alanine aminotransferase	21	2.3
Increased alkaline phosphatase	18	1.0
Increased creatinine	17	0

\*The denominator used to calculate the rate varied from 88 to 96 based on the number of patients with a baseline value and at least one post-treatment value.

## 7 DRUG INTERACTIONS

### 7.1 Effect of Other Drugs on TAZVERIK

#### Strong and Moderate CYP3A Inhibitors

Coadministration of TAZVERIK with a strong or moderate CYP3A inhibitor increases tazemetostat plasma concentrations [see *Clinical Pharmacology (12.3)*], which may increase the frequency or severity of adverse reactions. Avoid coadministration of strong or moderate CYP3A inhibitors with TAZVERIK. If coadministration of moderate CYP3A inhibitors cannot be avoided, reduce TAZVERIK dose [see *Dosage and Administration (2.3)*].

#### Strong and Moderate CYP3A Inducers

Coadministration of TAZVERIK with a strong or moderate CYP3A inducer may decrease tazemetostat plasma concentrations [see *Clinical Pharmacology (12.3)*], which may decrease the efficacy of TAZVERIK. Avoid coadministration of moderate and strong CYP3A inducers with TAZVERIK.

### 7.2 Effect of TAZVERIK on Other Drugs

#### CYP3A Substrates

Coadministration of TAZVERIK with CYP3A substrates, including hormonal contraceptives, can result in decreased concentrations and reduced efficacy of CYP3A substrates [see *Use in Specific Populations (8.3)*, *Clinical Pharmacology (12.3)*].

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### Risk Summary

Based on findings from animal studies and its mechanism of action [see *Clinical Pharmacology* (12.1)], TAZVERIK can cause fetal harm when administered to pregnant women. There are no available data on TAZVERIK use in pregnant women to inform the drug-associated risk. Administration of tazemetostat to pregnant rats and rabbits during organogenesis resulted in dose-dependent increases in skeletal developmental abnormalities in both species beginning at maternal exposures approximately 1.5 times the adult human exposure [ $AUC_{0-45h}$ ] at the 800 mg twice daily dose (see *Data*). Advise pregnant women of the potential risk to a fetus.

In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

## Data

### *Animal Data*

In pregnant rats, once daily oral administration of tazemetostat during the period of organogenesis from gestation day (GD) 7 through 17 resulted in no maternal adverse effects at doses up to 100 mg/kg/day (approximately 6 times the adult human exposure at 800 mg twice daily). Skeletal malformations and variations occurred in fetuses at doses of  $\geq 50$  mg/kg (approximately 2 times the adult human exposure at the 800 mg twice daily dose). At 200 mg/kg (approximately 14 times the adult human exposure at the 800 mg twice daily dose), major findings included increased post implantation loss, missing digits, fused vertebrae, domed heads and fused bones of the skull, and reduced fetal body weights.

In pregnant rabbits, no adverse maternal effects were observed after once daily oral administration of 400 mg/kg/day tazemetostat (approximately 7 times the adult human exposure at the 800 mg twice daily dose) from GD 7 through 19. Skeletal variations were present at doses  $\geq 100$  mg/kg/day (approximately 1.5 times the adult human exposure at the 800 mg twice daily dose), with skeletal malformations at  $\geq 200$  mg/kg/day (approximately 5.6 times the adult human exposure at the 800 mg twice daily dose). At 400 mg/kg (approximately 7 times the adult human exposure at the 800 mg twice daily dose), major findings included increased post implantation loss and cleft palate and snout.

## **8.2 Lactation**

### Risk Summary

There are no animal or human data on the presence of tazemetostat in human milk or on its effects on the breastfed child or milk production. Because of the potential risk for serious adverse reactions from TAZVERIK in the breastfed child, advise women not to breastfeed during treatment with TAZVERIK and for one week after the final dose.

## **8.3 Females and Males of Reproductive Potential**

### Pregnancy Testing

Verify the pregnancy status of females of reproductive potential prior to initiating TAZVERIK [see *Use in Specific Populations* (8.1)].

### Risk Summary

TAZVERIK can cause fetal harm when administered to pregnant women [see *Use in Specific Populations (8.1)*].

## Contraception

### *Females*

Advise females of reproductive potential to use effective non-hormonal contraception during treatment with TAZVERIK and for 6 months after the final dose. TAZVERIK can render some hormonal contraceptives ineffective [see *Drug Interactions (7.2)*].

### *Males*

Advise males with female partners of reproductive potential to use effective contraception during treatment with TAZVERIK and for at least 3 months after the final dose.

## **8.4 Pediatric Use**

The safety and effectiveness of TAZVERIK have been established in pediatric patients aged 16 years and older (adolescents) with metastatic or locally advanced epithelioid sarcoma. Use of TAZVERIK for this indication is supported by evidence from adequate and well-controlled studies in adults (including 3 adolescent patients aged 16 years) [see *Adverse Reactions (6.1)*, *Clinical Pharmacology (12.3)*, *Clinical Studies (14.1)*].

The safety and effectiveness of TAZVERIK in pediatric patients aged less than 16 years have not been established.

## Juvenile Animal Toxicity Data

In a 13-week juvenile rat toxicology study, animals were dosed daily from post-natal day 7 to day 97 (approximately equivalent to neonate to adulthood). Tazemetostat resulted in:

- T-LBL at doses  $\geq 50$  mg/kg (approximately 2.8 times the adult human exposure at the 800 mg twice daily dose)
- Increased trabecular bone at doses  $\geq 100$  mg/kg (approximately 10 times the adult human exposure at the 800 mg twice daily dose)
- Increased body weight at doses  $\geq 50$  mg/kg (approximately equal to the adult human exposure at the 800 mg twice daily dose)
- Distended testicles in males at doses  $\geq 50$  mg/kg (approximately equal to the adult human exposure at the 800 mg twice daily dose)

## **8.5 Geriatric Use**

Clinical studies of TAZVERIK did not include sufficient numbers of patients with epithelioid sarcoma or relapsed or refractory follicular lymphoma aged 65 and over to determine whether they respond differently from younger subjects.

## **8.6 Renal Impairment**

No dose adjustment of TAZVERIK is recommended for patients with mild to severe renal impairment or end stage renal disease [see *Clinical Pharmacology (12.3)*].


## **8.7 Hepatic Impairment**

No dose adjustment of TAZVERIK is recommended for patients with mild hepatic

impairment (total bilirubin > 1 to 1.5 times upper limit of normal [ULN] or AST > ULN). TAZVERIK has not been studied in patients with moderate (total bilirubin > 1.5 to 3 times ULN) or severe (total bilirubin > 3 times ULN) hepatic impairment [see *Clinical Pharmacology* (12.3)].

## 11 DESCRIPTION

Tazemetostat is a methyltransferase inhibitor. Tazemetostat hydrobromide has the following chemical name: [1,1'-Biphenyl]-3-carboxamide, *N*-[(1,2-dihydro-4,6-dimethyl-2-oxo-3-pyridinyl)methyl]-5-[ethyl(tetrahydro-2*H*-pyran-4-yl)amino]-4-methyl-4'-(4-morpholinylmethyl)-, hydrobromide (1:1). The molecular formula of tazemetostat hydrobromide is C<sub>34</sub>H<sub>44</sub>N<sub>4</sub>O<sub>4</sub>•HBr. Tazemetostat hydrobromide has a molecular weight of 653.66 g/mol and the following structural formula:



Tazemetostat hydrobromide is a white to off-white solid that is slightly soluble in water and has pKa values of 5.26, 6.88, and 12.62. A saturated aqueous solution of tazemetostat hydrobromide has a pH of approximately 5 at ambient conditions.

TAZVERIK (tazemetostat) tablets for oral use contain 200 mg tazemetostat, equivalent to 228 mg tazemetostat hydrobromide.

Each tablet is film-coated and contains the following inactive ingredients in the tablet core: hydroxypropyl cellulose, lactose monohydrate, low-substituted hydroxypropyl cellulose, magnesium stearate, and sodium starch glycolate. The film-coat contains hypromellose, polyethylene glycol, red iron oxide, talc, and titanium dioxide.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

Tazemetostat is an inhibitor of the methyltransferase, EZH2, and some EZH2 gain-of-function mutations including Y646X, A682G, and A692V. Tazemetostat also inhibited EZH1 with a half-maximal inhibitory concentration (IC<sub>50</sub>) of 392 nM, approximately 36 times higher than the IC<sub>50</sub> for inhibition of EZH2.

The most well-characterized function of EZH2 is as the catalytic subunit of the polycomb repressive complex 2 (PRC2), catalyzing mono-, di-, and trimethylation of lysine 27 of histone H3. Trimethylation of histone H3 leads to transcriptional repression.

SWItch/Sucrose Non-Fermentable (SWI/SNF) complexes can antagonize PRC2 function in the regulation of the expression of certain genes of patients with epithelioid sarcoma. Preclinical in vitro and in vivo models with the loss or dysfunction of certain SWI/SNF

complex members (e.g., integrase interactor 1 [INI1/SNF5/SMARCB1/BAF47], SMARCA4, and SMARCA2) can lead to aberrant EZH2 activity or expression and a resulting oncogenic dependence on EZH2.

Tazemetostat suppressed proliferation of B-cell lymphoma cell lines in vitro and demonstrated antitumor activity in a mouse xenograft model of B-cell lymphoma with or without EZH2 gain-of-function mutations. Tazemetostat demonstrated greater effects on the inhibition of proliferation of lymphoma cell lines with mutant EZH2.

## **12.2 Pharmacodynamics**

Tazemetostat exposure-response relationships and the time course of pharmacodynamic responses are unknown.

### Cardiac Electrophysiology

The effect of orally administered TAZVERIK, at doses ranging from 100 mg to 1600 mg twice daily (0.125 to 2 times the approved recommended dosage) for 15 days, on the heart-rate corrected QT (QTc) interval was evaluated in a dose-finding study in 38 patients with advanced malignancies. Tazemetostat and its metabolite EPZ-6930 did not cause a large mean increase (i.e. >20 ms) on the QTc interval at the 800 mg twice daily dose. The largest mean increase (upper bound of 90% confidence interval) in QTc were 6.1 ms (8.5 ms) and 9.3 ms (12.5 ms) at a dose of 800 mg twice daily and 1600 mg twice daily, respectively.

## **12.3 Pharmacokinetics**

The systemic exposure of tazemetostat is approximately dose proportional over the dose range of 200 mg to 1600 mg twice daily of TAZVERIK (0.25 to 2 times the approved recommended dosage). Following TAZVERIK 800 mg orally twice daily, steady-state was reached by Day 15. The mean (coefficient of variation [CV]%) steady-state peak plasma concentration ( $C_{max}$ ) was 829 (56%) ng/mL and  $AUC_{0-12h}$  was 3340 (49%) ng•h/mL. Tazemetostat exhibited time-dependent pharmacokinetics (PK). The mean accumulation ratio (measured by AUC) was 0.58.

### Absorption

The mean absolute oral bioavailability of tazemetostat is approximately 33%. The median time to reach the peak plasma concentration of tazemetostat is 1 to 2 hours.

### *Effect of Food*

A high-fat, high-calorie (approximately 800 to 1000 calories) meal does not have a significant effect on tazemetostat exposure.

### Distribution

The mean (CV%) apparent volume of distribution at steady-state ( $V_{ss}/F$ ) is 1230 L (46%). Tazemetostat is 88% bound to human plasma proteins in vitro. The blood-to-plasma ratio is 0.73.

### Elimination

At steady-state, the estimated mean (CV%) terminal elimination half-life of tazemetostat is 3.1 hours (14%) and the apparent total clearance ( $CL_{ss}/F$ ) is 274 L/h (49%).

### *Metabolism*

In vitro, tazemetostat is metabolized by CYP3A to form the inactive major metabolites M5 (EPZ-6930) and M3 (EPZ006931). M5 undergoes further metabolism by CYP3A.

### *Excretion*

Following a single oral dose of radiolabeled tazemetostat, 94% of the total radioactivity was recovered over 12 days, with 15% excreted into urine and 79% into feces.

### Specific Populations

Age (16 to 91 years), sex, race (White, Black, Asian), body weight (37.3 to 173 kg), mild hepatic impairment (total bilirubin > 1 to 1.5 times ULN or AST > ULN) and renal impairment, including end stage renal disease, have no clinically meaningful effect on the pharmacokinetics of tazemetostat. The effect of moderate to severe hepatic impairment has not been studied.

### Drug Interaction Studies

#### *Clinical Studies*

##### *Effect of CYP3A Inhibitors on Tazemetostat:*

Coadministration of fluconazole (a moderate CYP3A inhibitor) with TAZVERIK 400 mg twice daily in patients increased tazemetostat steady-state  $AUC_{0-8h}$  by 3.1-fold and  $C_{max}$  by 2.3-fold.

##### *Effect of Gastric Acid Reducing Agents on Tazemetostat:*

Coadministration of omeprazole (a proton pump inhibitor) with TAZVERIK 800 mg twice daily in patients increased tazemetostat steady-state  $AUC_{0-8h}$  by 26% and  $C_{max}$  by 25%, which is not expected to have clinically relevant impact.

##### *Effect of Tazemetostat on CYP3A Substrate:*

Coadministration of TAZVERIK 800 mg twice daily with oral midazolam (a sensitive CYP3A substrate) in patients decreased midazolam  $AUC_{0-12h}$  by 40% and  $C_{max}$  by 21%.

##### *Effect of Tazemetostat on CYP2C8 and CYP2C19 Substrates:*

Coadministration of TAZVERIK 800 mg twice daily with repaglinide (a sensitive CYP2C8 substrate) and omeprazole (a sensitive CYP2C19 substrate) in patients increased repaglinide  $AUC_{0-8h}$  by 80% and  $C_{max}$  by 51%; and had no effect on the exposure of omeprazole.

#### *In Vitro Studies*

##### *Metabolic Enzymes:*

Tazemetostat does not inhibit CYP1A2, CYP2B6, CYP2C9, and CYP2D6 at clinically relevant concentrations.

##### *Drug Transporters:*

Tazemetostat is a substrate of p-glycoprotein (P-gp). Tazemetostat is not a substrate of breast cancer resistance protein (BCRP); renal transporters organic cation transporter 2 (OCT2), organic anion transporter 3 (OAT3), and multidrug and toxin extrusion transporter 1 (MATE1); or hepatic transporters organic anion transporting polypeptide 1B1 (OATP1B1) and organic anion transporting polypeptide 1B3 (OATP1B3).

Tazemetostat is an inhibitor of MATE1 and multidrug and toxin extrusion transporter 2-K (MATE2-K). Tazemetostat does not inhibit P-gp, BCRP, OATP1B1, OATP1B3, organic cation transporter 1 (OCT1), OCT2, organic anion transporter 1 (OAT1), OAT3, or bile salt export pump (BSEP) at clinically relevant concentrations.

## 13 NONCLINICAL TOXICOLOGY

### 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Dedicated carcinogenicity studies were not conducted with tazemetostat, but T-LBL, MDS, AML, and B-ALL have been reported clinically and T-LBL occurred in juvenile and adult rats after ~9 or more weeks of tazemetostat administration during 13-week toxicity studies. Based on nonclinical studies in rats, the risk of T-LBL appears to be greater with longer duration dosing.

Tazemetostat did not cause genetic damage in a standard battery of studies including a screening and pivotal bacterial reverse mutation (Ames) assay, an in vitro micronucleus assessment in human peripheral blood lymphocytes, and an in vivo micronucleus assessment in rats after oral administration.

Fertility and early embryonic development studies have not been conducted with tazemetostat; however, an assessment of male and female reproductive organs were included in 4- and 13-week repeat-dose toxicity studies in rats and Cynomolgus monkeys. Oral daily administration of tazemetostat did not result in any notable effects in the adult male and female reproductive organs [see *Use in Specific Populations (8.3)*].

## 14 CLINICAL STUDIES

### 14.1 Epithelioid Sarcoma

The efficacy of TAZVERIK was evaluated in an open-label, single-arm cohort (Cohort 5) of a multi-center study (Study EZH-202, NCT02601950) in patients with histologically confirmed, metastatic or locally advanced epithelioid sarcoma. Patients were required to have INI1 loss, detected using local tests, and an Eastern Cooperative Oncology Group (ECOG) performance status (PS) of 0-2. Patients received TAZVERIK 800 mg orally twice daily until disease progression or unacceptable toxicity. Tumor response assessments were performed every 8 weeks. The major efficacy outcome measures were confirmed overall response rate (ORR) according to Response Evaluation Criteria in Solid Tumors (RECIST) v1.1 as assessed by blinded independent central review (BICR) and duration of response (DOR). Median duration of follow-up was 14 months (range 0.4 to 31).

Among the 62 patients who received TAZVERIK, median age was 34 years (range 16 to 79); 63% were male, 76% were White, 11% were Asian, 44% had proximal disease, 92% had an ECOG PS of 0 or 1, and 8% had an ECOG PS of 2. Prior surgery occurred in 77% of patients; 61% received prior systemic chemotherapy.

Efficacy results are summarized in Table 8.

**Table 8. Efficacy Results for Patients with Epithelioid Sarcoma Enrolled in Cohort 5 of Study EZH-202**

---



<b>Efficacy Endpoints</b>	<b>TAZVERIK N=62</b>
<b>Overall Response Rate (95% CI)*</b>	15% (7%, 26%)
Complete Response	1.6%
Partial Response	13%
<b>Duration of Response</b>	
% with duration $\geq$ 6 months	67%
Range in months	3.7, 24.5+

CI = Confidence Interval

\*Time to response ranged from 1.4 to 18.4 months.

## 14.2 Relapsed or Refractory Follicular Lymphoma

The efficacy of TAZVERIK was evaluated in two open-label, single-arm cohorts (Cohorts 4 and 5) of a multi-center study (Study E7438-G000-101, NCT01897571) in patients with histologically confirmed follicular lymphoma after at least 2 prior systemic therapies. Patients were required to have ECOG PS of 0-2 and were enrolled based on EZH2 mutation status. EZH2 mutations were identified prospectively using formalin-fixed, paraffin-embedded tumor samples, which were centrally tested using the cobas<sup>®</sup> EZH2 Mutation Test; the cobas EZH2 Mutation test is designed to detect the following mutations: Y646X [S,H,C], Y646F, Y646N, A682G, and A692V. Patients received TAZVERIK 800 mg orally twice daily until confirmed disease progression or unacceptable toxicity. Tumor response assessments were performed every 8 weeks through Week 24 and then every 12 weeks. The major efficacy outcome measures were ORR and DOR according to the International Working Group Non-Hodgkin Lymphoma (IWG-NHL) criteria<sup>1</sup> as assessed by Independent Review Committee. Median duration of follow-up was 22 months (range 3 months to 44 months) for patients with EZH2 MT positive tumors and 36 months (range 32 months to 39 months) for patients whose tumors did not have an EZH2 mutation detected.

A total of 99 patients were enrolled, including 45 patients whose tumors had one of these EZH2 mutations (mutant) and 54 patients whose tumors did not have one of these mutations (wild-type).

Among the 45 patients with EZH2 mutant follicular lymphoma, median age was 62 years (range 38 to 80), 58% were female, 42% had early progression following front-line therapy (POD24), and all had an ECOG PS of 0 or 1. Race was reported in 84% of patients; of these patients, 82% were White. Based on the cobas EZH2 Mutation test, 36%, 29%, 27%, 11% and 2% of patients had the following mutations: Y646X [S,H,C], Y646F, Y646N, A682G, and A692V, respectively. The median number of lines of prior systemic therapy was 2 (range 1 to 11), with 49% refractory to rituximab, 49% refractory to their last therapy, and 9% had received prior stem cell transplant.

Among the 54 patients with EZH2 wild-type follicular lymphoma, median age was 61 years (range 36 to 87), 63% were male, 59% had POD24, and 91% had an ECOG PS of 0 or 1. Race was reported in 57% of patients; of these patients, 48% were White and 3% were Asian. The median number of lines of prior systemic therapy was 3 (range 1 to 8), with 59% refractory to rituximab, 41% refractory to their last therapy, and 39% had received prior stem cell transplant.

The approval of TAZVERIK was based upon the efficacy in 95 patients (42 EZH2 Mutant, 53 EZH2 Wild-Type) who had received at least 2 prior systemic therapies and is presented in Table 9.

**Table 9. Efficacy Results for Patients with Relapsed or Refractory Follicular Lymphoma Enrolled into Cohorts 4 and 5 of Study E7438-G000-101**

Efficacy Endpoints	TAZVERIK N=95	
	EZH2 Mutant Follicular Lymphoma N=42	EZH2 Wild-Type Follicular Lymphoma N=53
<b>Overall Response Rate</b> (95% CI)*	69% (53%, 82%)	34% (22%, 48%)
Complete Response	12%	4%
Partial Response	57%	30%
<b>Duration of Response</b>		
Median (95% CI) in months	10.9 (7.2, NE)	13.0 (5.6, NE)
Range in months	0.0+, 22.1+	1, 22.5+

CI = Confidence Interval; NE = Not Estimable.

\*Median time to response for patients with EZH2 MT follicular lymphoma was 3.7 months (range 1.6 to 10.9) and for patients with EZH2 WT follicular lymphoma was 3.9 months (range 1.6 to 16.3).

## 15 REFERENCES

1. Cheson BD, Pfistner B, Juweid ME, et al. Revised response criteria for malignant lymphoma. *J Clin Oncol.* 2007;25(5):579-586.

## 16 HOW SUPPLIED/STORAGE AND HANDLING

TAZVERIK 200 mg film-coated tablets are red, round, biconvex shape and debossed with "EZM 200" on one side and plain on the other. TAZVERIK is available in:

Bottles of 240 tablets with a desiccant; NDC 72607-100-00

Do not store above 30°C (86°F).

## 17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide).

### Secondary Malignancies

Advise patients of the increased risk of secondary hematologic malignancies. Advise patients to inform their healthcare provider if they experience fatigue, easy bruising, fever, bone pain, or paleness [see *Warnings and Precautions* (5.1)].

### Embryo-Fetal Toxicity

Advise pregnant women and females of reproductive potential of the potential risk to a fetus. Advise females to inform their healthcare provider of a known or suspected pregnancy [see *Use in Specific Populations* (8.1)].

Advise females of reproductive potential to use effective non-hormonal contraception during treatment with TAZVERIK and for 6 months after the final dose [see *Use in Specific Populations* (8.3)].

Advise males with female partners of reproductive potential to use effective contraception during treatment with TAZVERIK and for 3 months after the final dose [see *Use in Specific Populations* (8.3), *Nonclinical Toxicology* (13.1)].

### Lactation

Advise women not to breastfeed during treatment with TAZVERIK and for 1 week after the final dose [see *Use in Special Populations* (8.2)].

### Drug Interactions

Advise patients and caregivers to inform their healthcare provider of all concomitant medications, including prescription medicines, over-the-counter drugs, vitamins, and herbal products. Inform patients to avoid St. John's wort, grapefruit, and grapefruit juice while taking TAZVERIK [see *Drug Interactions* (7.1)].

Manufactured for:  
Epizyme, Inc.  
An Ipsen Company  
One Main St.  
Cambridge, MA 02142

©2023 Epizyme, Inc.

## **MEDICATION GUIDE** **TAZVERIK® (taz vayr' ik)** **(tazemetostat)** **tablets**

### **What is the most important information I should know about TAZVERIK?**

#### **TAZVERIK can cause serious side effects, including:**

- **Risk of new cancers.** An increase in new (second) cancers has happened in people who were treated with TAZVERIK. Talk with your healthcare provider about your risk of developing new cancers. Your healthcare provider will monitor you for new cancers after your treatment with TAZVERIK. Tell your healthcare provider if you are more tired than usual, or have easy bruising, fever, bone pain, or paleness.

See "**What are the possible side effects of TAZVERIK**" for more information about side effects.

### **What is TAZVERIK?**

TAZVERIK is a prescription medicine used to treat:

- adults and children aged 16 years and older with epithelioid sarcoma that has spread or grown and cannot be removed by surgery.
- adults with follicular lymphoma when the disease has come back or did not respond to treatment, whose tumors have an abnormal EZH2 gene, **and** who have been treated with at least two prior medicines. Your healthcare provider will perform a test to make sure TAZVERIK is right for you.
- adults with follicular lymphoma when the disease has come back or did not respond to treatment, who have no other satisfactory treatment options.

It is not known if TAZVERIK is safe and effective in children less than 16 years of age.

**Before taking TAZVERIK tell your healthcare provider about all of your medical conditions, including if you:**

- are pregnant or plan to become pregnant. TAZVERIK can harm your unborn baby. Your healthcare provider will give you a pregnancy test before you start treatment with TAZVERIK. Tell your healthcare provider right away if you become pregnant or think you may be pregnant.
  - **Females** who are able to become pregnant should use effective non-hormonal birth control (such as condoms) during treatment and for 6 months after the final dose of TAZVERIK. Birth control pills (oral contraceptives) and other hormonal forms of birth control may not be effective if used during treatment with TAZVERIK. Talk to your healthcare provider about birth control options that are right for you.
  - **Males** with female partners who are able to become pregnant should use effective birth control during treatment and for 3 months after the final dose of TAZVERIK.
- are breastfeeding or plan to breastfeed. It is not known if TAZVERIK passes into your breast milk. Do not breastfeed during treatment and for 1 week after the final dose of TAZVERIK.

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. TAZVERIK may affect the way other medicines work and other medicines may affect how TAZVERIK works.

**How should I take TAZVERIK?**

- Take TAZVERIK exactly as your healthcare provider tells you.
- Take TAZVERIK 2 times each day.
- Take TAZVERIK with or without food.
- Swallow TAZVERIK tablets whole. Do not cut, crush, or chew tablets.
- If you miss a dose or vomit after taking your dose, just skip that dose and take the next dose at your regular time.
- Your healthcare provider may change your dose, temporarily stop, or completely stop treatment with TAZVERIK if you get certain side effects.

**What should I avoid while taking TAZVERIK?**

- Avoid eating grapefruit or drinking grapefruit juice during treatment with TAZVERIK.
- Avoid taking St. John's wort during treatment with TAZVERIK.

**What are the possible side effects of TAZVERIK?**

**TAZVERIK can cause serious side effects. See "What is the most important information I should know about TAZVERIK?"**

**The most common side effects of TAZVERIK in people with epithelioid sarcoma include:**

- |             |                      |                |
|-------------|----------------------|----------------|
| • pain      | • nausea             | • vomiting     |
| • tiredness | • decreased appetite | • constipation |

**The most common side effects of TAZVERIK in people with follicular lymphoma include:**

- tiredness
- cold-like symptoms (upper respiratory infection)
- bone and muscle pain
- nausea
- stomach (abdominal) pain

These are not all the possible side effects of TAZVERIK.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

### **How should I store TAZVERIK?**

Do not store TAZVERIK tablets above 86°F (30°C).

**Keep TAZVERIK and all medicines out of the reach of children.**

### **General information about the safe and effective use of TAZVERIK.**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use TAZVERIK for a condition for which it was not prescribed. Do not give TAZVERIK to other people, even if they have the same symptoms that you have. It may harm them. You can ask your pharmacist or healthcare provider for information about TAZVERIK that is written for health professionals.

### **What are the ingredients in TAZVERIK?**

**Active Ingredient:** tazemetostat.

**Inactive Ingredients:** Tablet core: hydroxypropyl cellulose, lactose monohydrate, low-substituted hydroxypropyl cellulose, magnesium stearate, and sodium starch glycolate. Film-coating: hypromellose, polyethylene glycol, red iron oxide, talc, and titanium dioxide. Manufactured for: Epizyme, Inc.

An Ipsen Company

One Main St.

Cambridge, MA 02142 For more information, contact Epizyme at 855-463-5127.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Issued: November/2023

## **Principal Display Panel - 200 mg Bottle Label**

NDC 72607-100-00

240 Tablets

TAZVERIK®  
(tazemetostat) tablets

200 mg

Rx only

Dispense the Medication Guide, attached or provided separately, to each patient pursuant to Federal Law.

Epizyme



NDC 72607-100-00 240 Tablets

**TAZVERIK**  
(tazemetostat) tablets

**200 mg**

**Rx only**

Dispense the Medication Guide, attached or  
provided separately, to each patient pursuant  
to Federal Law.



**Storage:** Do not store above 30°C  
(86°F).

**Do not use if safety seal under  
cap is broken or missing.**

**Recommended Dosage:** See  
prescribing information.

Each tablet contains 200 mg  
tazemetostat equivalent to 228 mg  
tazemetostat hydrobromide.

**Manufactured for:**  
Epizyme, Inc.  
An Ipsen Company  
One Main St.  
Cambridge, MA 02142

© 2023 Epizyme, Inc.

1077297

70046516

## TAZVERIK

tazemetostat tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:72607-100
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>tazemetostat hydrobromide</b> (UNII: 6P89T5M073) (tazemetostat - UNII:Q40W93WPE1)	tazemetostat	200 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>Lactose monohydrate</b> (UNII: EWQ57Q8I5X)	
<b>Low-substituted hydroxypropyl cellulose, unspecified</b> (UNII: 2165RE0K14)	
<b>Hydroxypropyl cellulose, unspecified</b> (UNII: 9XZ8H6N6OH)	
<b>Sodium starch glycolate Type A potato</b> (UNII: 5856J3G2A2)	
<b>Magnesium stearate</b> (UNII: 70097M6I30)	
<b>Hypromellose, unspecified</b> (UNII: 3NXW29V3WO)	
<b>Talc</b> (UNII: 7SEV7J4R1U)	
<b>Polyethylene glycol 8000</b> (UNII: Q662QK8M3B)	
<b>Titanium dioxide</b> (UNII: 15FIX9V2JP)	

<b>Ferric oxide red</b> (UNII: 1K09F3G675)				
<b>Product Characteristics</b>				
<b>Color</b>	red (RED)	<b>Score</b>	no score	
<b>Shape</b>	ROUND (ROUND)	<b>Size</b>	10mm	
<b>Flavor</b>		<b>Imprint Code</b>	EZ M;200	
<b>Contains</b>				
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:72607-100-00	240 in 1 BOTTLE; Type 0: Not a Combination Product	01/23/2020	
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	
NDA	NDA211723	01/23/2020		

**Labeler** - Epizyme, Inc. (018795119)

Establishment			
Name	Address	ID/FEI	Business Operations
Patheon Pharmaceuticals Inc.		005286822	MANUFACTURE(72607-100)

Establishment			
Name	Address	ID/FEI	Business Operations
Sterling Pharma Solutions Limited		349157623	API MANUFACTURE(72607-100)