

RANITIDINE- ranitidine tablet, coated

Chain Drug Marketing Association

Drug Facts

Active ingredient (in each tablet)

Ranitidine 150 mg (as ranitidine hydrochloride USP, 168 mg)

Purpose

Acid reducer

Use(s)

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers
- if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

Stop use and ask doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water
 - to **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

Other information

- do not use if printed foil under bottle cap is open or torn
- store at 20°-25°C (68°-77°F)
- avoid excessive heat or humidity
- protect from light
- this product is sodium and sugar free

Inactive ingredients

FD&C red #40 aluminum lake, hypromellose, iron oxide black, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide.

Questions?

Call **1-888-375-3784**

Carton Label



6 0 8 1 7 0 0 5 1



**Maximum Strength
Acid Reducer**
Ranitidine Tablets USP,
150 mg

NDC 63868-480-24

Compare to the
Active Ingredient in
Zantac 150[®] Tablets*

**Maximum Strength
Acid Reducer**
Ranitidine Tablets USP,
150 mg

Acid Reducer

Prevents and Relieves
Heartburn Associated with
Acid Indigestion and Sour Stomach



24 Tablets (24 Doses)



6 35515 49643 4



Distributed by C.D.M.A., Inc.
43157 W 9 Mile Rd
Novi, MI 48375
www.qualitychoice.com
Questions: 248-449-9300

Drug Facts

Active ingredient (in each tablet)
Ranitidine 150 mg (as ranitidine hydrochloride USP, 168 mg) **Purpose**
Acid reducer

Uses ■ relieves heartburn associated with acid indigestion and sour stomach
■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings
Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers
Do not use ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. ■ with other acid reducers
■ if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have
■ had heartburn over 3 months. This may be a sign of a more serious condition.
■ heartburn with lightheadedness, sweating or dizziness
■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
■ frequent chest pain
■ frequent wheezing, particularly with heartburn
■ unexplained weight loss
■ nausea or vomiting
■ stomach pain

Stop use and ask a doctor if ■ your heartburn continues or worsens
■ you need to take this product for more than 14 days

Directions
■ adults and children 12 years and over:
■ to relieve symptoms, swallow 1 tablet with a glass of water
■ to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn
■ can be used up to twice daily (do not take more than 2 tablets in 24 hours)
■ children under 12 years: ask a doctor

Drug Facts (continued)

Other information
■ do not use if printed foil under bottle cap is open or torn
■ store at 20°-25°C (68°-77°F)
■ avoid excessive heat or humidity
■ protect from light
■ this product is sodium and sugar free

Inactive ingredients
FD&C red #40 aluminum lake, hydroxytoluene, iron oxide black, magnesium stearate, microcrystalline cellulose, polyethylene glycol, titanium dioxide.

Questions? Call 1-888-375-3784

TAMPER EVIDENT: DO NOT USE IF FOIL SEAL UNDER CAP PRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING.

IMPORTANT: Read the directions, consumer information leaflet and warnings before use. Keep the carton. It contains important information.

*This product is not manufactured or distributed by Chatham, Inc., owner of the registered trademark Zantac 150[®]

Bottle label



NDC 63868-480-24

**Maximum Strength
Acid Reducer**
Ranitidine Tablets USP,
150 mg

Acid Reducer

Prevents and Relieves
Heartburn Associated with
Acid Indigestion and Sour Stomach

24 Tablets (24 Doses)

TAMPER EVIDENT: DO NOT USE IF FOIL SEAL UNDER CAP PRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING.

IMPORTANT: This label does not contain full product information. See carton for complete information. Read the directions, consumer information leaflet and warnings before use. Retain carton and leaflet for reference.

Active ingredient (in each tablet) Ranitidine 150 mg (as ranitidine hydrochloride USP, 168 mg) **Purpose** Acid reducer

Uses ■ relieves heartburn associated with acid indigestion and sour stomach ■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages **Warnings. Allergy alert:** Do not use if you are allergic to ranitidine or other acid reducers **Do not use** ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. ■ with other acid reducers ■ if you have kidney disease, except under the advice and supervision of a doctor **Ask a doctor before use if you have** ■ had heartburn over 3 months. This may be a sign of a more serious condition ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain **Stop use and ask a doctor if** ■ your heartburn worsens or you need to take this product for more than 14 days **If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ adults and children 12 years and over:
■ to relieve symptoms, swallow 1 tablet with a glass of water
■ to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn ■ can be used up to twice daily (do not take more than 2 tablets in 24 hours)
■ children under 12 years: ask a doctor

Questions? Call 1-888-375-3784

Distributed by C.D.M.A., Inc.
43157 W 9 Mile Rd
Novi, MI 48375

LOT

EXP

150071808

LOT

EXP

RANITIDINE

ranitidine tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63868-480(NDC:55111-404)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Ranitidine Hydrochloride (UNII: BK76465IHM) (RANITIDINE - UNII:884KT10YB7)	RANITIDINE	150 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HYPROMELLOSE 2910 (6 MPA.S) (UNII: 0WZ8WG20P6)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
magnesium stearate (UNII: 70097M6I30)	
cellulose, microcrystalline (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
titanium dioxide (UNII: 15FIX9V2JP)	

Product Characteristics

Color	PINK	Score	no score
Shape	ROUND	Size	9mm
Flavor		Imprint Code	R150
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-480-50	1 in 1 CARTON	07/01/2018	09/30/2019
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:63868-480-24	1 in 1 CARTON	07/01/2018	09/30/2019
2		24 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA078192	07/01/2018	

