

**ASPIRIN LOW DOSE- aspirin tablet, chewable  
ARMY AND AIR FORCE EXCHANGE SERVICE**

-----  
**Exchange Select 44-218 Delisted**

***Active ingredient (in each chewable tablet)***

Aspirin 81 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever

***Uses***

for the temporary relief of minor aches and pains or as recommended by your doctor

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have had stomach ulcers or bleeding problems
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor
- if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have asthma
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - have bloody or black stools
  - vomit blood
  - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- new symptoms occur
- pain gets worse or lasts more than 10 days
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

***Directions***

- drink a full glass of water with each dose
- chew or crush tablets completely before swallowing; do not swallow tablets whole
- adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 chewable tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

***Other information***

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

## ***Inactive ingredients***

corn starch, dextrans hydrated, FD&C yellow #6 aluminum lake, flavor, maltodextrin, sucralose

## ***Questions or comments?***

**1-800-426-9391**

## ***Principal Display Panel***

exchange ✓ **select**<sup>™</sup>

Compare To The Active Ingredient of  
Bayer<sup>®</sup> Chewable Low Dose Aspirin<sup>†</sup>

Talk to your Doctor  
or other Healthcare  
provider before  
using this product  
for your heart.

## **LOW DOSE ASPIRIN Chewable Tablets**

Pain Reliever (**NSAID**)

•Aspirin Regimen •Orange flavored

**81** mg

Actual  
Size

Chew or crush tablets completely  
before swallowing.  
Do not swallow tablets whole.

## **3 BOTTLES INSIDE**

36 + 36 + 36 Tablets

✓ quality  
value

## **Value Pack**

**3 bottles x 36 tablets each =  
108 tablets**

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

<sup>†</sup>This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer<sup>®</sup> Chewable Low Dose Aspirin.

50844

REV1118A21807

**“SATISFACTION GUARANTEED OR YOUR MONEY BACK”**

Manufactured For Your Military Exchanges

Distributed by: LNK International, Inc.

Hauppauge, NY 11788

1-800-426-9391

exchange select™

# LOW DOSE ASPIRIN Chewable Tablets

Compare To The Active Ingredient of Bayer® Chewable Low Dose Aspirin†

Talk to your Doctor or other Healthcare provider before using this product for your heart.

## Pain Reliever (NSAID)

• Aspirin Regimen • Orange Flavored

**3 BOTTLES INSIDE**  
36 + 36 + 36 Tablets

Chew or crush tablets completely before swallowing. Do not swallow tablets whole.



exchange select™

# LOW DOSE ASPIRIN Chewable Tablets

Compare To The Active Ingredient of Bayer® Chewable Low Dose Aspirin†

Talk to your Doctor or other Healthcare provider before using this product for your heart.

## Pain Reliever (NSAID)

• Aspirin Regimen • Orange Flavored

**3 BOTTLES INSIDE**  
36 + 36 + 36 Tablets

Chew or crush tablets completely before swallowing. Do not swallow tablets whole.



# Value Pack

3 bottles x 36 tablets each = 108 tablets

**Value Pack**  
3 bottles x 36 tablets each = 108 tablets

exchange select™

# LOW DOSE ASPIRIN Chewable Tablets

Compare To The Active Ingredient of Bayer® Chewable Low Dose Aspirin†

Talk to your Doctor or other Healthcare provider before using this product for your heart.

## Pain Reliever (NSAID)

• Aspirin Regimen • Orange Flavored

**3 BOTTLES INSIDE**  
36 + 36 + 36 Tablets

Chew or crush tablets completely before swallowing. Do not swallow tablets whole.



exchange select™  
**LOW DOSE ASPIRIN**  
Chewable Tablets



TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

“SATISFACTION GUARANTEED OR YOUR MONEY BACK”  
Manufactured For Your Military Exchanges  
Distributed by: LNK International, Inc.  
Happauge, NY 11788  
1-800-426-9391

†This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Chewable Low Dose Aspirin.  
50844 REV1118A21807

B-0066E-218A-07BR  
REV1118A21807

Questions or comments?  
1-800-426-9391

as mentioned  
■ you have high blood pressure, heart disease,  
liver cirrhosis, or kidney disease  
■ you have asthma ■ you are taking a diuretic

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

**Drug Facts**

**Active ingredient**  
(in each chewable tablet)  
Aspirin 81 mg (NSAID) . . . . . Pain reliever  
nonsteroidal anti-inflammatory drug

**Purpose**  
Ask a doctor or pharmacist before use if you are taking a prescription drug for  
gout ■ diabetes ■ arthritis

**Stop use and ask a doctor if**  
■ an allergic reaction occurs. Seek medical help right away.  
■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ vomit blood

**Warnings**  
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. **Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:  
■ hives ■ facial swelling ■ shock ■ asthma (wheezing)  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ have had stomach ulcers or bleeding problems ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Do not use**  
■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor ■ if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**  
■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn

**Drug Facts (continued)**  
Ask a doctor or pharmacist before use if you are taking a prescription drug for  
gout ■ diabetes ■ arthritis

**Stop use and ask a doctor if**  
■ an allergic reaction occurs. Seek medical help right away.  
■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ vomit blood

**Warnings**  
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. **Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:  
■ hives ■ facial swelling ■ shock ■ asthma (wheezing)  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ have had stomach ulcers or bleeding problems ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Do not use**  
■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor ■ if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**  
■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn

**Inactive ingredients**  
dextrates hydroxyethylcellulose ■ aluminum lake, flavor, maltodextrin, sucralose

**Directions**  
■ drink a full glass of water with each dose ■ chew or crush tablets completely before swallowing; do not swallow tablets whole ■ adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 chewable tablets in 24 hours unless directed by a doctor ■ children under 12 years: do not use unless directed by a doctor

**Other information**  
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ see end flap for expiration date and lot number

**Drug Facts (continued)**  
Ask a doctor or pharmacist before use if you are taking a prescription drug for  
gout ■ diabetes ■ arthritis

**Stop use and ask a doctor if**  
■ an allergic reaction occurs. Seek medical help right away.  
■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ vomit blood

**Warnings**  
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. **Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:  
■ hives ■ facial swelling ■ shock ■ asthma (wheezing)  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you are age 60 or older ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others] ■ have had stomach ulcers or bleeding problems ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

**Do not use**  
■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor ■ if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if**  
■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn

**Inactive ingredients**  
dextrates hydroxyethylcellulose ■ aluminum lake, flavor, maltodextrin, sucralose

**Directions**  
■ drink a full glass of water with each dose ■ chew or crush tablets completely before swallowing; do not swallow tablets whole ■ adults and children 12 years and over: take 4 to 8 chewable tablets every 4 hours not to exceed 48 chewable tablets in 24 hours unless directed by a doctor ■ children under 12 years: do not use unless directed by a doctor

**Other information**  
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ see end flap for expiration date and lot number

**Inactive ingredients**  
dextrates hydroxyethylcellulose ■ aluminum lake, flavor, maltodextrin, sucralose

Exchange Select 44-218A

**ASPIRIN LOW DOSE**

aspirin tablet, chewable

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:55301-821
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

**Inactive Ingredients**

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
DEXTROSE MONOHYDRATE (UNII: LX22YL083G)	
FD&C YELLOW NO. 6 ALUMINUM LAKE (UNII: GYP6Z2JR6Q)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	

SUCRALOSE (UNII: 96K6UQ3ZD4)

### Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	8mm
<b>Flavor</b>	ORANGE	<b>Imprint Code</b>	44;218
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55301-821-07	3 in 1 CARTON	09/12/1990	03/02/2025
1		36 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	09/12/1990	03/02/2025

**Labeler** - ARMY AND AIR FORCE EXCHANGE SERVICE (001695568)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(55301-821)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(55301-821)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(55301-821)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(55301-821)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(55301-821)

Revised: 4/2024

ARMY AND AIR FORCE EXCHANGE SERVICE