ERGOCALCIFEROL- ergocalciferol liquid Virtus Pharmaceuticals

Ergocalciferol Oral Solution, USP

69543-234-60

Vitamin D Supplement Drops 8000 lU (0.2mg) Ergocalciferol per mL

Supplement Facts			
Serving Size: 400 IU			
Serving Per Container: 1200			
	Amount Per	% Daily Value	
	Serving	% Daily Value	
Vitamin D (ergocalciferol)	400 IU	100%	

Other Ingredients: Propylene Glycol

The US RDA for Vitamin D is 400 IU for children and adults.

Description

Each mL contains 8000 IU (0.2 mg) of ergocalciferol (Vitamin D₂) dissolved in propylene glycol.

Directions for Use

400 IU daily to be taken orally using the supplied pre-measured syringe. Submerge the tip of the syringe into the liquid, pull plunger back to bring the liquid to the 400 IU mark, and dispense into mouth.

DO NOT FILL SYRINGE ENTIRELY.

Contraindications

Contraindicated in people with hypercalcemia, malabsorption syndrome, abnormal sensitivity to the toxic effects of vitamin D, and hypervitaminosis D.

Warning

In case of accidental overdose contact a doctor or poison control center immediately.

Hypersensitivity to vitamin D may be a causative factor in infants with idiopathic hypercalcemia. In these children, vitamin D must be strictly restricted.

Caution

Adequate dietary calcium is necessary for response to vitamin D therapy.

Side Effects

Metabolic side effects include hypercalcemia.

You may report serious side effects by calling Virtus at 1-888-848-3593 or FDA at 1-800-FDA-1088

How Supplied

Each 60 mL polycarbonate screw top bottle contains 60 mL ergocalciferol in a propylene glycol base (0.2 mg ergocalciferol per mL).

Store at 20°-25° C (68°-77°F); excursion permitted between 15°-30° C (59°-86° F).

Adequate intake (AI) (minimum daily intake) of vitamin D is needed to support calcium metabolism and bone health, particularly in the young and old.¹

KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Tamper Resistant Feature:

Do **NOT** use if imprinted seal on cap is broken or missing.

Manufactured for:

Virtus Pharmaceuticals, LLC Tampa, FL 33619 1-888-848-3593 Made in USA

7274

Rev. 09/15

PRINCIPAL DISPLAY PANEL - 60 mL Bottle Carton

VIRTUS

PHARMACEUTICALS

69543-234-60

Ergocalciferol Oral Solution, USP

Vitamin D Supplement Drops 8000 IU (0.2mg) Ergocalciferol per mL

60 mL (2 fl. oz.)

¹ This statement has not been evaluated by the FDA. This product is not intended to diagnose, treat, cure, or prevent any disease.



69543-234-60

Ergocalciferol

Oral Solution, USP

Vitamin D Supplement Drops 8000 IU (0.2mg) Ergocalciferol per mL

60 mL (2 fl. oz.)



69543-234-60

Ergocalciferol

Oral Solution, USP

Vitamin D Supplement Drops 8000 IU (0.2mg) Ergocalciferol per mL

Supplement Facts

Serving Size: 400 IU Serving Per Container: 1200

> Amount Per % Daily Value Serving

Vitamin D (ergocalciferol) 400 IU

100%

Other Ingredients: Propylene Glycol

The US RDA for Vitamin D is 400 IU for children and adults.





Ergocalciferol

Oral Solution, USP

Vitamin D Supplement Drops 8000 IU (0.2mg) Ergocalciferol per mL

> Vitamin D Supplement Drops 8000 IU (0.2mg) Ergocalciferol per mL

Oral Solution, USP

Ergocalciferol

09-752-57369





69543-234-60

Ergocalciferol

Oral Solution, USP

Vitamin D Supplement Drops 8000 IU (0.2mg) Ergocalciferol per mL

Description: Each mL contains 8000 IU (0.2 mg) of ergocalciferol (Vitamin D₂) dissolved in propylene glycol.

Directions for Use: 400 IU daily to be taken orally using the supplied pre-measured syringe. Submerge the tip of the syringe into the liquid, pull plunger back to bring the liquid to the 400 IU mark, and dispense into mouth.



69543-234-60

Ergocalciferol

Oral Solution, USP

Vitamin D Supplement Drops 8000 IU (0.2mg) Ergocalciferol per mL

DO NOT FILL SYRINGE ENTIRELY.

Contraindications: Contraindicated in people with hypercalcemia, malabsorption syndrome, abnormal sensitivity to the toxic effects of vitamin D, and hypervitaminosis D.

Warning: In case of accidental overdose contact a doctor or poison control center immediately.

Hypersensitivity to vitamin D may be a causative factor in infants with idiopathic hypercalcemia. In these children, vitamin D must be strictly restricted,

Caution: Adequate dietary calcium is necessary for response to vitamin D therapy.

Side Effects: Metabolic side effects include hypercalcemia.

You may report serious side effects by calling Virtus at 1-888-848-3593 or FDA at 1-800-FDA-1088

How Supplied: Each 60 mL polycarbonate screw top bottle contains 60 mL ergocalciferol in a propylene glycol base (0.2 mg ergocalciferol per mL). Store at 20°-25° C (68°-77°F); excursion permitted between 15°-30° C

Manufactured for:

Virtus Pharmaceuticals, LLC Tampa, FL 33619 1-888-848-3593

7274 Rev. 09/15

Made in USA

(59°-86° F).

Adequate intake (AI) (minimum daily intake) of vitamin D is needed to support calcium metabolism and bone health, particularly in the young and old.*

*This statement has not been evaluated by the FDA. This product is not intended to diagnose, treat, cure, or prevent any disease.

KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

Tamper Resistant Feature:

Do **NOT** use if imprinted seal on cap is broken or missing.

ERGOCALCIFEROL

ergocalciferol liquid

Product Information

Product Type	DIETARY SUPPLEMENT	Item Code (Source)	NHRIC:69543-234
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Ergocalciferol (UNII: VS041H42XC) (Ergocalciferol - UNII:VS041H42XC)	Ergocalciferol	0.2 mg in 1 mL

Inactive Ingredients

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NHRIC:69543-234-60	1 in 1 CARTON		
1		60 mL in 1 BOTTLE		

Strength

Ingredient Name

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
DIETARY SUPPLEMENT		08/25/2015	

Labeler - Virtus Pharmaceuticals (079659493)

Propylene Glycol (UNII: 6DC9Q167V3)

Revised: 2/2016 Virtus Pharmaceuticals