

PAIN RELIEVER CHILDRENS- acetaminophen tablet, chewable
H E B

HEB 44-450-children

Active ingredient (in each chewable tablet)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - sore throat
 - flu
 - toothache
 - the common cold
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this

product

Ask a doctor before use if your child has

liver disease.

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin.

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.
- chew or crush tablets completely before swallowing; do not swallow tablets whole
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give for more than 5 days unless directed by a doctor

Weight (lb)	Age (yr)	Dose (chewable tablets)*
Under 24	Under 2	Ask a doctor
24-35	2-3	1 tablet
36-47	4-5	1 1/2 tablets
48-59	6-8	2 tablets
60-71	9-10	2 1/2 tablets
72-95	11	3 tablets

*or as directed by a doctor

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- avoid high humidity
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

citric acid, crospovidone, D&C red #27 aluminum lake, D&C red #30 aluminum lake, dextrates hydrated, ethylcellulose, flavor, magnesium stearate, mannitol, polyethylene, stearic acid, sucralose

Questions or comments?

1-800-426-9391

Principal display panel

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts
Active ingredient (in each chewable tablet)
 Acetaminophen 160 mg Pain reliever/fever reducer

Purpose
 Pain reliever/fever reducer

Uses
 temporarily relieves minor aches and pains due to:
 ■ headache ■ sore throat ■ flu
 ■ toothache ■ the common cold
 ■ temporarily reduces fever

Warnings
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes:
 ■ more than 5 doses in 24 hours, which is the maximum daily amount
 ■ with other drugs containing acetaminophen
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin reddening ■ blisters ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.
 Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Drug Facts (continued)
Do not use
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if your child is allergic to acetaminophen or any of the inactive ingredients in this product
Ask a doctor before use if your child has liver disease.
Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.
When using this product do not exceed recommended dosage.
Stop use and ask a doctor if
 ■ pain gets worse or lasts more than 5 days
 ■ fever gets worse or lasts more than 3 days
 ■ new symptoms occur
 ■ redness or swelling is present
 These could be signs of a serious condition.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ this product does not contain directions or complete warnings for adult use
 ■ do not give more than directed
 ■ find the right dose on chart below. If possible, use weight to dose; otherwise, use age.
 ■ chew or crush tablets completely before swallowing; do not swallow tablets whole
 ■ repeat dose every 4 hours while symptoms last
 ■ do not give more than 5 times in 24 hours
 ■ do not give for more than 5 days unless directed by a doctor

Weight (lb)	Age (yr)	Dose (chewable tablets)*
Under 24	Under 2	Ask a doctor
24-35	2-3	1 tablet
36-47	4-5	1 1/2 tablets
48-59	6-8	2 tablets
60-71	9-10	2 1/2 tablets
72-95	11	3 tablets

* or as directed by a doctor

Other information
 ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
 ■ avoid high humidity
 ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Drug Facts (continued)
 ■ use by expiration date on package

Inactive ingredients citric acid, crospovidone, D&C red #27 aluminum lake, D&C red #30 aluminum lake, dextrates hydrated, ethylcellulose, flavor, magnesium stearate, mannitol, polyethylene, stearic acid, sucralose

Questions or comments?
 1-800-426-9391

This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Children's Tylenol®.
 50844 REV0522C45008

100% GUARANTEE
Disentle
 If you aren't completely pleased with this product, we'll be happy to replace it or refund your money. You have our word on it.

MADE WITH PRIDE & CARE FOR H-E-B®
 SAN ANTONIO, TX 78204

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

20580-2110
 0 41220 36008 6

NO PRINT/NO VARNISH
 LOT & EXP DATE

B-0712-450-08-R
 REV0522C45008

Compare to Children's Tylenol® active ingredient†
 NDC 37808-945-08

H-E-B®
Children's Pain Reliever
Acetaminophen 160 mg
Chewable Tablets
Pain Reliever/Fever Reducer

For ages 2 to 11

- Relieves Pain + Fever
- Ibuprofen Free
- Aspirin Free

Bubblegum Flavor

24 CHEWABLE TABLETS

actual size
 Chew or crush tablets completely before swallowing. Do not swallow tablets whole.

HEB 44-450

Compare to **Children's Tylenol®**
 active ingredient†

NDC 37808-945-08

H-E-B®

Children's

Pain Reliever

Acetaminophen 160 mg

Chewable Tablets

Pain Reliever/Fever Reducer

For ages 2 to 11

- **Relieves Pain + Fever**
- **Ibuprofen Free**
- **Aspirin Free**

**24 CHEWABLE
TABLETS**

Bubblegum
Flavor

actual size

Chew or crush tablets
completely before swallowing.
Do not swallow tablets whole.

**TAMPER EVIDENT: DO NOT USE IF
PACKAGE IS OPENED OR IF BLISTER
UNIT IS TORN, BROKEN OR SHOWS
ANY SIGNS OF TAMPERING**

† This product is not manufactured or distributed
by Johnson & Johnson Corporation, owner of the
registered trademark Children's Tylenol®.
50844 REV0522C45008

H-E-B®

100%

Guarantee
promise

If you aren't completely pleased
with this product, we'll be happy to
replace it or refund your money.
You have our word on it.

**MADE WITH
PRIDE & CARE FOR H-E-B®
SAN ANTONIO, TX 78204**

PAIN RELIEVER CHILDRENS

acetaminophen tablet, chewable

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37808-945
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
CROSPROVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
D&C RED NO. 27 ALUMINUM LAKE (UNII: ZK64F7XSTX)	
D&C RED NO. 30 (UNII: 2S42T2808B)	
DEXTROSE MONOHYDRATE (UNII: LX22YL083G)	
ETHYLCELLULOSE, UNSPECIFIED (UNII: 7Z8S9VYZ4B)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MANNITOL (UNII: 3OWL53L36A)	
HIGH DENSITY POLYETHYLENE (UNII: UG00KM4WR7)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	pink	Score	2 pieces
Shape	ROUND	Size	16mm
Flavor	BUBBLE GUM	Imprint Code	44;450
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-945-08	4 in 1 CARTON	02/25/2005	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/25/2005	

Labeler - H E B (007924756)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(37808-945)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(37808-945) , pack(37808-945)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(37808-945)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(37808-945)

Revised: 11/2023

H E B