

**GLYCERIN- glycerin suppository**  
**Cardinal Health**

-----  
**Leader Children's Glycerin Suppository**

**Active ingredient (in each suppository)**

Glycerin 1g

**Purpose**

Laxative

**Uses**

- for relief of occasional constipation
- this product generally produces bowel movement in 1/4 to 1 hour

**Warnings**

**For rectal use only**

May cause rectal discomfort or a burning sensation.

**Ask a doctor before using any laxative if child has**

- abdominal pain, nausea or vomiting
- a sudden change in bowel habits lasting more than 2 weeks
- already used a laxative for more than 1 week

**Stop use and consult a doctor if child has**

- rectal bleeding
- no bowel movement within 1 hour using this product

These symptoms may indicate a serious condition.

**Keep out of reach of children**

**If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.**

**Directions - single daily dose**

children 2 to under 6 years - 1 suppository, or as directed by a doctor

children under 2 - ask a doctor

Insert suppository well up into rectum. Suppository need not melt completely to produce laxative action.

## Other information

- Store container tightly closed.
- Keep away from excessive heat.

## Inactive ingredients

purified water, sodium hydroxide, stearic acid

## Questions or comments?

1-800-492-5988

## Label

The product package shown below represents a sample of that currently in use. Additional packaging may also be available.

Children's Glycerin Suppositories, 25 count

Distributed by Cardinal Health, Dublin OH 43017

CH ChildSupp.jpg

**LEADER<sup>2</sup>**

NDC 70000-0429-1

**Children's Glycerin Suppositories**  
Laxative

Ages 2-5 Years  
For Relief of Occasional Constipation  
Rectal Laxative

**COMPARE TO FLEET<sup>®</sup> PEDIA-LAX<sup>®</sup> GLYCERIN SUPPOSITORIES active ingredient<sup>\*</sup>**

**25 SUPPOSITORIES**

**TAMPER EVIDENT: DO NOT USE IF PRINTED INNER SEAL IS DAMAGED OR MISSING.**

**Drug Facts**

**Active ingredient (in each suppository)**  
Glycerin 1g .....Laxative

**Uses**  
■ for relief of occasional constipation  
■ this product generally produces bowel movement in 1/4 to 1 hour

**Warnings**  
For rectal use only  
May cause rectal discomfort or a burning sensation.  
**Ask a doctor before using any laxative if child has**  
■ abdominal pain, nausea or vomiting  
■ a sudden change in bowel habits lasting more than 2 weeks  
■ already used a laxative for more than 1 week  
**Stop use and consult a doctor if child has**  
■ rectal bleeding  
■ no bowel movement within 1 hour of using this product  
These symptoms may indicate a serious condition.  
Keep out of reach of children.  
If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions** Single daily dosage  
children 2 to under 6 years 1 suppository, or as directed by a doctor  
children under 2 years ask a doctor  
Insert suppository well up into rectum. Suppository need not melt completely to produce laxative action.

**Other Information**  
■ Store container tightly closed.  
■ Keep away from excessive heat.

**Inactive ingredients**  
purified water, sodium hydroxide, stearic acid

**Questions or Comments?**  
1-800-492-5988

Dist. by CH, Dublin, OH 43017 ©2018 Cardinal Health  
\*This product is not manufactured or distributed by C.S. Peet Company, Inc., owner of the registered trademark, Fleet<sup>®</sup> Pedia-Lax<sup>®</sup> Glycerin Suppositories.  
**100% Money Back Guarantee**  
Return to place of purchase if not satisfied.  
CIN 5478557 REV. 12/18

0 96295 13613 5

**NO VARNISH**

## GLYCERIN

glycerin suppository

### Product Information

|                                |                |                           |                |
|--------------------------------|----------------|---------------------------|----------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:70000-0429 |
| <b>Route of Administration</b> | RECTAL         |                           |                |

### Active Ingredient/Active Moiety

| Ingredient Name  | Basis of Strength | Strength |
|--|-------------------|----------|
| GLYCERIN (UNII: PDC6A3C0OX) (GLYCERIN - UNII:PDC6A3C0OX) | GLYCERIN          | 1 g      |

### Inactive Ingredients

| Ingredient Name                            | Strength |
|--|----------|
| <b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I) |          |
| <b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)     |          |
| <b>WATER</b> (UNII: 059QF0KO0R)            |          |

### Packaging

| # | Item Code        | Package Description                            | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:70000-0429-1 | 25 in 1 JAR; Type 0: Not a Combination Product | 01/02/2019           |                    |

### Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M007                                     | 01/02/2019           |                    |

**Labeler** - Cardinal Health (063997360)

**Registrant** - DSC Laboratories, Inc. (097807374)

### Establishment

| Name                   | Address | ID/FEI    | Business Operations     |
|------------------------|---------|-----------|-------------------------|
| DSC Laboratories, Inc. |         | 097807374 | manufacture(70000-0429) |

Revised: 12/2024

Cardinal Health