

**CIRCLE K DAYTIME COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, and phenylephrine hydrochloride capsule, liquid filled  
Lil' Drug Store Products, Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Circle K Daytime Cold & Flu**

**Drug Facts**

<b>Active ingredients (in each softgel)</b>	<b>Purposes</b>
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

**Uses**

- temporarily relieves common cold and flu symptoms:
  - minor aches and pains
  - headache
  - sore throat
  - nasal congestion
  - fever
  - cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy Alert**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- diabetes
- heart disease
- thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin.

**When using this product, do not exceed recommended dosage.**

**Stop use and ask a doctor if**

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning**

Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in 24 hours

- adults and children 12 years and over: take 2 softgels with water every 4 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- **when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing**

#### **Other information**

- store between 15°-30°C (59°-86°F)
- avoid excessive heat

#### **Inactive ingredients**

butylated hydroxyanisole, butylated hydroxytoluene, FD&C yellow #6, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

#### **Questions or comments?**

call toll-free **1-877-507-6516**  
**(M-F 8AM-4:30PM CST)**

**Proudly distributed by Circle K Stores Inc.**

#### **PRINCIPAL DISPLAY PANEL - 16 Capsule Blister Pack Carton**

Compare to the Active Ingredients in  
Vicks® DayQuil® LiquiCaps®\*\*

CIRCLE K

Daytime Cold & Flu

Acetaminophen, Dextromethorphan HBr, and Phenylephrine HCl  
Pain Reliever/Fever Reducer, Cough Suppressant, Nasal Decongestant

Non-Drowsy Relief of

- Aches - Fever - Sore Throat
- Nasal Congestion - Cough

ACTUAL SIZE

16 Softgels  
(Liquid-filled capsules)

CIRCLE K

## Daytime Cold & Flu

**Acetaminophen**, Dextromethorphan HBr, and Phenylephrine HCl  
Pain Reliever/Fever Reducer, Cough Suppressant, Nasal Decongestant

Compare to the Active Ingredients in  
Vicks® DayQuil® LiquiCaps®\*\*

CIRCLE K

## Daytime Cold & Flu

**Acetaminophen**, Dextromethorphan HBr, and Phenylephrine HCl  
Pain Reliever/Fever Reducer, Cough Suppressant, Nasal Decongestant



ACTUAL SIZE

### Non-Drowsy Relief of

- Aches - Fever - Sore Throat
- Nasal Congestion - Cough

**16** Softgels  
(Liquid-filled capsules)

Product manufactured for:  
Lil' Drug Store Products, Inc.  
9300 Earhart Lane SW  
Cedar Rapids, IA 52404

Proudly distributed by Circle K Stores Inc.



**16** Softgels  
(Liquid-filled capsules)

CIRCLE K

## Daytime Cold & Flu

**Acetaminophen**, Dextromethorphan HBr, and Phenylephrine HCl  
Pain Reliever/Fever Reducer, Cough Suppressant, Nasal Decongestant



Lot No:  
Exp Date:

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

### Drug Facts

#### Active ingredients (in each softgel)

Acetaminophen 325 mg.....Pain reliever/fever reducer  
Dextromethorphan HBr 10 mg.....Cough suppressant  
Phenylephrine HCl 5 mg.....Nasal decongestant

#### Purposes

#### Uses

- temporarily relieves common cold and flu symptoms:
- minor aches and pains • headache • sore throat
- nasal congestion • fever
- cough due to minor throat and bronchial irritation

#### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy Alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's

### Drug Facts (continued)

disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

#### Ask a doctor before use if you have

- liver disease • diabetes
- heart disease • thyroid disease
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.**

**When using this product, do not exceed recommended dosage.**

#### Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- nervousness, dizziness, or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Drug Facts (continued)

#### Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 4 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

#### Other information

- store between 15°-30°C (59°-86°F)
- avoid excessive heat

#### Inactive ingredients

butylated hydroxyanisole, butylated hydroxytoluene, FD&C yellow #6, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

#### Questions or comments?

call toll-free 1-877-507-6516  
(M-F 8AM-4:30PM CST)

\*\*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, DayQuil®, and LiquiCaps® are registered trademarks of The Procter & Gamble Company.

### PARENTS:

Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

57718C-US-11-19-PA



## CIRCLE K DAYTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, and phenylephrine hydrochloride capsule, liquid filled

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66715-5671
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>BUTYLATED HYDROXYANISOLE</b> (UNII: REK4960K2U)	
<b>BUTYLATED HYDROXYTOLUENE</b> (UNII: 1P9D0Z171K)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GELATIN, UNSPECIFIED</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	

### Product Characteristics

Color	orange	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	P19
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start	Marketing End
---	-----------	---------------------	-----------------	---------------

#	Item Code	Package Description	Date	Date
1	NDC:66715-5671-8	8 in 1 CARTON	03/01/2020	07/13/2024
1		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
<b>Marketing Category</b>		<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC monograph final		part341	09/05/2014	07/13/2024

**Labeler -**
Lil' Drug Store Products, Inc. (093103646)