

ETODOLAC- etodolac tablet, film coated, extended release  
Zydus Pharmaceuticals (USA) Inc.

-----  
**Etodolac Extended-release Tablets, USP**

**Cardiovascular Thrombotic Events**

**WARNING: RISK OF SERIOUS CARDIOVASCULAR AND GASTROINTESTINAL EVENTS**

**Cardiovascular Thrombotic Events**

**Nonsteroidal anti-inflammatory drugs (NSAIDs) cause an increased risk of serious cardiovascular thrombotic events, including myocardial infarction and stroke, which can be fatal. This risk may occur early in treatment and may increase with duration of use [see *Warnings and Precautions*].**

- **Etodolac extended-release tablets are contraindicated in the setting of coronary artery bypass graft (CABG) surgery [see *Contraindications and Warnings and Precautions*].**


**Gastrointestinal Bleeding, Ulceration, and Perforation**

**NSAIDs cause an increased risk of serious gastrointestinal (GI) adverse events including bleeding, ulceration, and perforation of the stomach or intestines, which can be fatal. These events can occur at any time during use and without warning symptoms. Elderly patients and patients with a prior history of peptic ulcer disease and/or GI bleeding are at greater risk for serious GI events [see *Warnings and Precautions*].**

**DESCRIPTION**

Etodolac extended-release tablets, USP contain etodolac, which is a member of the pyranocarboxylic acid group of non-steroidal anti-inflammatory drugs (NSAIDs). Each tablet contains etodolac for oral administration. Etodolac is a racemic mixture of [+]*S* and [-]*R*-enantiomers. Etodolac, USP is a white to off-white crystalline powder, insoluble in water but soluble in alcohols, chloroform, dimethyl sulfoxide, and aqueous polyethylene glycol.

The chemical name is (±) 1,8-diethyl-1,3,4,9-tetrahydropyrano-[3,4-*b*]indole-1-acetic acid. It has the following structural formula:



C<sub>17</sub>H<sub>21</sub>NO<sub>3</sub> M.W. 287.36

Each etodolac extended-release tablet, USP intended for oral administration contains 400 mg or 500 mg or 600 mg of etodolac. In addition, each tablet contains the following inactive ingredients: disodium hydrogen phosphate, ethylcellulose, hypromellose, lactose monohydrate, magnesium stearate, polyethylene glycol, talc, titanium dioxide and triacetin. Additionally each 400 mg tablet contains: D&C yellow # 10 aluminum lake, FD&C Red # 40 aluminum lake and FD&C yellow # 6 aluminum lake and each 500 mg tablet contains: FD&C Blue # 2 aluminum lake, iron oxide black and iron oxide yellow and each 600 mg tablet contains: FD&C blue # 2 aluminum lake, iron oxide red and iron oxide yellow.

The USP Drug Release test complies with USP Dissolution Test 1.

## CLINICAL PHARMACOLOGY

### Pharmacodynamics

Etodolac extended-release tablets are a non-steroidal anti-inflammatory drug (NSAID) that exhibits anti-inflammatory, analgesic, and antipyretic activities in animal models. The mechanism of action of etodolac extended-release tablets, like that of other NSAIDs, is not completely understood, but may be related to prostaglandin synthetase inhibition.

### Pharmacokinetics

#### Absorption

Etodolac extended-release tablets and etodolac tablets both contain etodolac, but differ in their release characteristics. The systemic availability of etodolac from etodolac extended-release tablets is generally greater than 80%. Etodolac does not undergo significant first-pass metabolism following oral administration. After oral administration of etodolac extended-release tablets in doses up to 800 mg once daily, peak concentrations occur approximately 6 hours after dosing and are dose proportional for both total and free etodolac.

**Table 1** shows the comparison of etodolac pharmacokinetic parameters after the administration of etodolac tablets and etodolac extended-release tablets.

**Table 2** shows the etodolac pharmacokinetic parameters in various populations. The data from patients with renal and hepatic impairment were obtained following administration of (immediate-release) etodolac tablets.

**Table 1**

Pharmacokinetic Parameters	Mean (CV)%*	
	Etodolac Tablets	Etodolac Extended-release Tablets
Extent of Oral Absorption (Bioavailability) [F]	≥ 80%	≥ 80%
Time to Peak Concentration (T <sub>max</sub> ), h	1.4 (61%)	6.7 (47%)
Oral Clearance (CL/F), mL/h/kg	49.1 (33%)	46.8 (37%)
Apparent Volume of Distribution (V <sub>d</sub> /F), mL/kg	393 (29%)	566 (26%)
Terminal Half-Life (t <sub>1/2</sub> ), h	6.4 (22%)	8.4 (30%)

\* % Coefficient of variation

**Table 2 Mean (CV%)\* Pharmacokinetic Parameters of Etodolac in Normal Healthy Adults and Various Special Populations**

PK Parameters	Etodolac Extended-release Tablets				Etodolac Tablets			
	Normal Healthy Adults	Healthy Males	Healthy Females	Elderly (> 65 yr)	Hemodialysis <sup>†</sup> (24 to 65) (n = 9)		Renal Impairment <sup>†</sup>	Hepatic Impairment <sup>†</sup>
	(18 to 44) <sup>‡</sup> (n = 116)	(18 to 43) (n = 102)	(25 to 44) (n = 14)	(66 to 88) (n = 24)	Dialysis On	Dialysis Off	(46 to 73) (n = 10)	(34 to 60) (n = 9)
T <sub>max</sub> , h	6.7 (47%)*	6.8 (45%)	4.5 (56%)	6.2 (51%)	1.7 (88%)	0.9 (67%)	2.1 (46%)	1.1 (15%)
Oral Clearance, mL/h/kg (CL/F)	46.8 (37%)	46.8 (37%)	47.2 (38%)	51.6 (40%)	NA	NA	58.3 (19%)	42 (43%)
Apparent Volume of Distribution, mL/kg (V <sub>d</sub> /F)	566 (26%)	580 (26%)	459 (28%)	552 (34%)	NA	NA	NA	NA
Terminal Half-Life, h	8.4 (30%)	8.4 (29%)	7.6 (45%)	7.8 (26%)	5.1 (22%)	7.5 (34%)	NA	5.7 (24%)

\* % Coefficient of variation

† Pharmacokinetic parameters obtained following administration of etodolac tablets

‡ Age range (years)

NA = not available

## Food/Antacid Effects

Food has no significant effect on the extent of etodolac extended-release tablets absorption, however, food significantly increased  $C_{max}$  (54%) following a 600 mg dose.

The extent of absorption of etodolac is not affected when etodolac is administered with antacid. Coadministration, with an antacid, decreases the peak concentration reached by about 15 to 20% with no measurable effect on time-to-peak.

## Distribution

The mean apparent volume of distribution ( $V_d/F$ ) of etodolac following administration of etodolac extended-release tablets is 566 mL/kg. Etodolac is more than 99% bound to plasma proteins, primarily to albumin, and is independent of etodolac concentration over the dose range studied. It is not known whether etodolac is excreted in human milk. However, based on its physical-chemical properties, excretion into breast milk is expected.

## Metabolism

Etodolac metabolites do not contribute significantly to the pharmacological activity of etodolac extended-release tablets.

Following administration of immediate-release etodolac, several metabolites have been identified in human plasma and urine. Other metabolites remain to be identified. The metabolites include 6-, 7-, and 8-hydroxylated etodolac and etodolac glucuronide. After a single dose of  $^{14}C$ -etodolac, hydroxylated metabolites accounted for less than 10% of total drug in serum. On chronic dosing, hydroxylated-etodolac metabolites do not accumulate in the plasma of patients with normal renal function. The extent of accumulation of hydroxylated-etodolac metabolites in patients with renal dysfunction has not been studied. The role, if any, of a specific cytochrome P450 system in the metabolism of etodolac is unknown. The hydroxylated-etodolac metabolites undergo further glucuronidation followed by renal excretion and partial elimination in the feces.

## Excretion

The mean oral clearance of etodolac following oral etodolac extended-release tablets dosing is 47 ( $\pm 17$ ) mL/h/kg. The terminal half-life ( $t_{1/2}$ ) of etodolac after etodolac extended-release tablets administration is 8.4 hours compared to 6.4 hours for etodolac tablets. Approximately 1% of an etodolac tablet dose is excreted unchanged in the urine, with 72% of the dose excreted into the urine as parent drug plus metabolites:

- etodolac, unchanged	1%
- etodolac glucuronide	13%
- hydroxylated metabolites (6-, 7-, and 8-OH)	5%
- hydroxylated metabolite glucuronides	20%
- unidentified metabolites	33%

Fecal excretion accounted for 16% of the dose.

## Special Populations

### Geriatric

In clinical studies, age was not shown to have any effect on half-life or protein binding, and demonstrated no change in expected drug accumulation. No dosage adjustment is generally necessary in the elderly on the basis of pharmacokinetics. The elderly may need dosage adjustment, however, as they may be more sensitive to antiprostaglandin effects than younger patients (see **PRECAUTIONS, Geriatric Use**).

## ***Pediatric***

The pharmacokinetics of etodolac extended-release tablets were assessed in an open-label, 12 week clinical trial which included plasma sampling for population pharmacokinetics. Seventy-two (72) patients, 6 to 16 years of age, with juvenile rheumatoid arthritis, received etodolac extended-release tablets in doses of 13.3 to 21.3 mg/kg given as 400 to 1000 mg once daily. The results from a population pharmacokinetic analysis based on the 59 subjects who completed the trial are as follows:

**Table 3 Pharmacokinetic Parameter Estimates for Etodolac Extended-release Tablets in Patients with Juvenile Rheumatoid Arthritis**

<b>Parameter</b>	<b>JRA* (Age:6 to 16)† n = 59</b>
Oral Clearance (CL/F), mL/h/kg	47.8 (38%)
Apparent Volume of Distribution (Vd/F), mL/kg	78.9 (61%)
Half-life (t <sub>1/2</sub> ), h	12.1 (75%)

\* Mean (CV) of parameter estimates predicted from population pharmacokinetics

† Age range (years)

While similar, the pharmacokinetic parameters for children with juvenile rheumatoid arthritis did not directly correlate with adult pharmacokinetic data in rheumatoid arthritis. In the population pharmacokinetic analysis, body weights below 50 kg were found to correlate with CL/F (see **DOSAGE AND ADMINISTRATION**).

## **Race**

Pharmacokinetic differences due to race have not been identified. Clinical studies included patients of many races, all of whom responded in a similar fashion.

## ***Hepatic Insufficiency***

The pharmacokinetics of etodolac following administration of etodolac extended-release tablets have not been investigated in subjects with hepatic insufficiency. Following administration of etodolac tablets, the plasma protein binding and disposition of total and free etodolac were unchanged in the presence of compensated hepatic cirrhosis. Although no dosage adjustment is generally required in patients with chronic hepatic diseases, etodolac clearance is dependent on liver function and could be reduced in patients with severe hepatic failure.

## ***Renal Insufficiency***

The pharmacokinetics of etodolac following administration of etodolac extended-release tablets have not been investigated in subjects with renal insufficiency. Etodolac renal clearance following administration of etodolac tablets was unchanged in the presence of mild-to-moderate renal failure (creatinine clearance, 37 to 88 mL/min). Although renal elimination is a significant pathway of excretion for etodolac metabolites, no dosing adjustment in patients with mild to moderate renal dysfunction is generally necessary. Etodolac plasma protein binding decreases in patients with severe renal deficiency. Etodolac should be used with caution in such patients because, as with other NSAIDs, it may further decrease renal function in some patients. Etodolac is not significantly removed from the blood in patients undergoing hemodialysis.

## **CLINICAL STUDIES**

### **Arthritis**

The use of etodolac extended-release tablets in managing the signs and symptoms of osteoarthritis of the knee and rheumatoid arthritis was assessed in doubleblind, randomized, parallel, controlled clinical trials in 1552 patients. In these trials, etodolac extended-release tablets, given once daily, provided efficacy comparable to immediate-release etodolac.

The safety, efficacy, and pharmacokinetics of etodolac extended-release tablets were assessed in an open-label, 12 week clinical trial. Seventy-two (72) patients, 6 to 16 years of age, with juvenile

rheumatoid arthritis, received etodolac extended-release tablets in doses of 400 to 1000 mg (13.3 to 21.3 mg/kg body weight) once daily. At these doses, etodolac extended-release tablets controlled the signs and symptoms of juvenile rheumatoid arthritis. Based on the results of this study, the safety profile of etodolac extended-release tablets (at doses not exceeding 20 mg/kg) appeared to be similar to that observed in the adult arthritic patients in clinical trials (see **PRECAUTIONS, Pediatric Use**).

## **INDICATIONS AND USAGE**

Carefully consider the potential benefits and risks of etodolac extended-release tablets, USP and other treatment options before deciding to use etodolac extended-release tablets, USP. Use the lowest effective dose for the shortest duration consistent with individual patient treatment goals (see **WARNINGS**).

Etodolac extended-release tablets, USP are indicated:

- \* For relief of signs and symptoms of juvenile arthritis
- \* For relief of the signs and symptoms of rheumatoid arthritis
- \* For relief of the signs and symptoms of osteoarthritis

## **CONTRAINDICATIONS**

Etodolac extended-release tablets are contraindicated in patients with known hypersensitivity to etodolac.

Etodolac extended-release tablets should not be given to patients who have experienced asthma, urticaria, or allergic-type reactions after taking aspirin or other NSAIDs. Severe, rarely fatal, anaphylactoid-like reactions to NSAIDs have been reported in such patients (see **WARNINGS, Anaphylactoid Reactions** and **PRECAUTIONS, Preexisting Asthma**).

Etodolac extended-release tablets are contraindicated in the setting of coronary artery bypass graft (CABG) surgery (see **WARNINGS**).

## **WARNINGS**

### **Cardiovascular Effects**

#### ***Cardiovascular Thrombotic Events***

Clinical trials of several COX-2 selective and nonselective NSAIDs of up to three years duration have shown an increased risk of serious cardiovascular (CV) thrombotic events, including myocardial infarction (MI) and stroke, which can be fatal. Based on available data, it is unclear that the risk for CV thrombotic events is similar for all NSAIDs. The relative increase in serious CV thrombotic events over baseline conferred by NSAID use appears to be similar in those with and without known CV disease or risk factors for CV disease. However, patients with known CV disease or risk factors had a higher absolute incidence of excess serious CV thrombotic events, due to their increased baseline rate. Some observational studies found that this increased risk of serious CV thrombotic events began as early as the first weeks of treatment. The increase in CV thrombotic risk has been observed most consistently at higher doses.

To minimize the potential risk for an adverse CV event in NSAID-treated patients, use the lowest effective dose for the shortest duration possible. Physicians and patients should remain alert for the development of such events, throughout the entire treatment course, even in the absence of previous CV symptoms. Patients should be informed about the symptoms of serious CV events and the steps to take if they occur.

There is no consistent evidence that concurrent use of aspirin mitigates the increased risk of serious CV thrombotic events associated with NSAID use. The concurrent use of aspirin and an NSAID, such as etodolac, increases the risk of serious gastrointestinal (GI) events [*see Warnings*].

#### ***Status Post Coronary Artery Bypass Graft (CABG) Surgery***

Two large, controlled clinical trials of a COX-2 selective NSAID for the treatment of pain in the first

10–14 days following CABG surgery found an increased incidence of myocardial infarction and stroke. NSAIDs are contraindicated in the setting of CABG [see *Contraindications*].

### **Post-MI Patients**

Observational studies conducted in the Danish National Registry have demonstrated that patients treated with NSAIDs in the post-MI period were at increased risk of reinfarction, CV-related death, and all-cause mortality beginning in the first week of treatment. In this same cohort, the incidence of death in the first year post MI was 20 per 100 person years in NSAID-treated patients compared to 12 per 100 person years in non-NSAID exposed patients. Although the absolute rate of death declined somewhat after the first year post-MI, the increased relative risk of death in NSAID users persisted over at least the next four years of follow-up.

Avoid the use of etodolac extended-release tablets in patients with a recent MI unless the benefits are expected to outweigh the risk of recurrent CV thrombotic events. If etodolac extended-release tablets are used in patients with a recent MI, monitor patients for signs of cardiac ischemia.

### **Hypertension**

NSAIDs, including etodolac extended-release tablets, can lead to onset of new hypertension or worsening of preexisting hypertension, either of which may contribute to the increased incidence of CV events. Patients taking thiazides or loop diuretics may have impaired response to these therapies when taking NSAIDs. NSAIDs, including etodolac extended-release tablets, should be used with caution in patients with hypertension. Blood pressure (BP) should be monitored closely during the initiation of NSAID treatment and throughout the course of therapy.

### **Heart Failure and Edema**

The Coxib and traditional NSAID Trialists' Collaboration meta-analysis of randomized controlled trials demonstrated an approximately two-fold increase in hospitalizations for heart failure in COX-2 selective-treated patients and nonselective NSAID-treated patients compared to placebo-treated patients. In a Danish National Registry study of patients with heart failure, NSAID use increased the risk of MI, hospitalization for heart failure, and death.

Additionally, fluid retention and edema have been observed in some patients treated with NSAIDs. Use of etodolac may blunt the CV effects of several therapeutic agents used to treat these medical conditions [e.g., diuretics, ACE inhibitors, or angiotensin receptor blockers (ARBs)] [see *Drug Interactions*].

Avoid the use of etodolac extended-release tablets in patients with severe heart failure unless the benefits are expected to outweigh the risk of worsening heart failure. If etodolac extended-release tablets are used in patients with severe heart failure, monitor patients for signs of worsening heart failure.

### **Gastrointestinal Effects - Risk of Ulceration, Bleeding, and Perforation**

NSAIDs, including etodolac extended-release tablets, can cause serious gastrointestinal (GI) adverse events including inflammation, bleeding, ulceration, and perforation of the stomach, small intestine, or large intestine, which can be fatal. These serious adverse events can occur at any time, with or without warning symptoms, in patients treated with NSAIDs. Only one in five patients, who develop a serious upper GI adverse event on NSAID therapy, is symptomatic. Upper GI ulcers, gross bleeding, or perforation caused by NSAIDs occur in approximately 1% of patients treated for 3 to 6 months, and in about 2 to 4% of patients treated for one year. These trends continue with longer duration of use, increasing the likelihood of developing a serious GI event at some time during the course of therapy. However, even short-term therapy is not without risk.

NSAIDs should be prescribed with extreme caution in those with a prior history of ulcer disease or gastrointestinal bleeding. Patients with a *prior history of peptic ulcer disease and/or gastrointestinal bleeding* who use NSAIDs have a greater than 10 fold increased risk for developing a GI bleed compared to patients with neither of these risk factors. Other factors that increase the risk for GI bleeding in patients treated with NSAIDs include concomitant use of oral corticosteroids or anticoagulants, longer duration of NSAID therapy, smoking, use of alcohol, older age, and poor general health status. Most spontaneous reports of fatal GI events are in elderly or debilitated patients and

therefore, special care should be taken in treating this population.

To minimize the potential risk for an adverse GI event in patients treated with an NSAID, the lowest effective dose should be used for the shortest possible duration. Patients and physicians should remain alert for signs and symptoms of GI ulceration and bleeding during NSAID therapy and promptly initiate additional evaluation and treatment if a serious GI adverse event is suspected. This should include discontinuation of the NSAID until a serious GI adverse event is ruled out. For high risk patients, alternate therapies that do not involve NSAIDs should be considered.

### **Renal Effects**

Long-term administration of NSAIDs has resulted in renal papillary necrosis and other renal injury. Renal toxicity has also been seen in patients in whom renal prostaglandins have a compensatory role in the maintenance of renal perfusion. In these patients, administration of a nonsteroidal anti-inflammatory drug may cause a dose-dependent reduction in prostaglandin formation and, secondarily, in renal blood flow, which may precipitate overt renal decompensation. Patients at greatest risk of this reaction are those with impaired renal function, heart failure, liver dysfunction, those taking diuretics and ACE inhibitors, and the elderly. Discontinuation of NSAID therapy is usually followed by recovery to the pretreatment state.

### ***Advanced Renal Disease***

No information is available from controlled clinical studies regarding the use of etodolac extended-release tablets in patients with advanced renal disease. Therefore, treatment with etodolac extended-release tablets is not recommended in these patients with advanced renal disease. If etodolac extended-release tablet therapy must be initiated, close monitoring of the patient's renal function is advisable.

### **Anaphylactoid Reactions**

As with other NSAIDs, anaphylactoid reactions may occur in patients without known prior exposure to etodolac extended-release tablets. Etodolac extended-release tablets should not be given to patients with the aspirin triad. This symptom complex typically occurs in asthmatic patients who experience rhinitis with or without nasal polyps, or who exhibit severe, potentially fatal bronchospasm after taking aspirin or other NSAIDs (see **CONTRAINDICATIONS** and **PRECAUTIONS, Preexisting Asthma**). Emergency help should be sought in cases where an anaphylactoid reaction occurs.

### **Skin Reactions**

NSAIDs, including etodolac extended-release tablets, can cause serious skin adverse events such as exfoliative dermatitis, Stevens-Johnson Syndrome (SJS), and toxic epidermal necrolysis (TEN), which can be fatal. These serious events may occur without warning. Patients should be informed about the signs and symptoms of serious skin manifestations and use of the drug should be discontinued at the first appearance of skin rash or any other sign of hypersensitivity.

### **Pregnancy**

In late pregnancy, as with other NSAIDs, etodolac extended-release tablets should be avoided because it may cause premature closure of the ductus arteriosus.

## **PRECAUTIONS**

### **General**

Etodolac extended-release tablets cannot be expected to substitute for corticosteroids or to treat corticosteroid insufficiency. Abrupt discontinuation of corticosteroids may lead to disease exacerbation. Patients on prolonged corticosteroid therapy should have their therapy tapered slowly if a decision is made to discontinue corticosteroids.

The pharmacological activity of etodolac extended-release tablets in reducing fever and inflammation may diminish the utility of these diagnostic signs in detecting complications of presumed noninfectious, painful conditions.

### **Hepatic Effects**

Borderline elevations of one or more liver tests may occur in up to 15% of patients taking NSAIDs including etodolac extended-release tablets. These laboratory abnormalities may progress, may remain unchanged, or may be transient with continuing therapy. Notable elevations of ALT or AST (approximately three or more times the upper limit of normal) have been reported in approximately 1% of patients in clinical trials with NSAIDs. In addition, rare cases of severe hepatic reactions, including jaundice and fatal fulminant hepatitis, liver necrosis and hepatic failure, some of them with fatal outcomes have been reported.

A patient with symptoms and/or signs suggesting liver dysfunction, or in whom an abnormal liver test has occurred, should be evaluated for evidence of the development of a more severe hepatic reaction while on therapy with etodolac extended-release tablets. If clinical signs and symptoms consistent with liver disease develop, or if systemic manifestations occur (e.g., eosinophilia, rash, etc.), etodolac extended-release tablets should be discontinued.

### **Hematological Effects**

Anemia is sometimes seen in patients receiving NSAIDs, including etodolac extended-release tablets. This may be due to fluid retention, occult or gross GI blood loss, or an incompletely described effect upon erythropoiesis. Patients on long-term treatment with NSAIDs, including etodolac extended-release tablets, should have their hemoglobin or hematocrit checked if they exhibit any signs or symptoms of anemia.

NSAIDs inhibit platelet aggregation and have been shown to prolong bleeding time in some patients. Unlike aspirin, their effect on platelet function is quantitatively less, of shorter duration, and reversible. Patients receiving etodolac extended-release tablets who may be adversely affected by alterations in platelet function, such as those with coagulation disorders or patients receiving anticoagulants, should be carefully monitored.

### **Preexisting Asthma**

Patients with asthma may have aspirin-sensitive asthma. The use of aspirin in patients with aspirin-sensitive asthma has been associated with severe bronchospasm which can be fatal. Since cross reactivity, including bronchospasm, between aspirin and other nonsteroidal anti-inflammatory drugs has been reported in such aspirin-sensitive patients, etodolac extended-release tablets should not be administered to patients with this form of aspirin sensitivity and should be used with caution in patients with preexisting asthma.

### **Information for Patients**

**Patients should be informed of the following information before initiating therapy with an NSAID and periodically during the course of ongoing therapy. Patients should also be encouraged to read the NSAID Medication Guide that accompanies each prescription dispensed.**

1. **Cardiovascular Thrombotic Events:** Advise patients to be alert for the symptoms of cardiovascular thrombotic events, including chest pain, shortness of breath, weakness, or slurring of speech, and to report any of these symptoms to their health care provider immediately [see *Warnings*].
2. Etodolac extended-release tablets, like other NSAIDs, can cause GI discomfort and, rarely, serious GI side effects, such as ulcers and bleeding, which may result in hospitalization and even death. Although serious GI tract ulcerations and bleeding can occur without warning symptoms, patients should be alert for the signs and symptoms of ulcerations and bleeding, and should ask for medical advice when observing any indicative sign or symptoms including epigastric pain, dyspepsia, melena, and hematemesis. Patients should be apprised of the importance of this follow-up (see **WARNINGS, Gastrointestinal Effects - Risk of Ulceration, Bleeding, and Perforation**).
3. Etodolac extended-release tablets, like other NSAIDs, can cause serious skin side effects such as exfoliative dermatitis, SJS, and TEN, which may result in hospitalizations and even death. Although serious skin reactions may occur without warning, patients should be alert for the signs and symptoms of skin rash and blisters, fever, or other signs of hypersensitivity such as itching, and should ask for medical advice when observing any indicative signs or symptoms. Patients should be advised to stop the drug immediately if they develop any type of rash and contact their physicians as soon as possible.



4. **Heart Failure And Edema:** Advise patients to be alert for the symptoms of congestive heart failure including shortness of breath, unexplained weight gain, or edema and to contact their healthcare provider if such symptoms occur [*see Warnings*].
5. Patients should be informed of the warning signs and symptoms of hepatotoxicity (e.g., nausea, fatigue, lethargy, pruritus, jaundice, right upper quadrant tenderness, and "flu-like" symptoms). If these occur, patients should be instructed to stop therapy and seek immediate medical therapy.
6. Patients should be informed of the signs of an anaphylactoid reaction (e.g., difficulty breathing, swelling of the face or throat). If these occur, patients should be instructed to seek immediate emergency help (see **WARNINGS**).
7. In late pregnancy, as with other NSAIDs, etodolac extended-release tablets should be avoided because it will cause premature closure of the ductus arteriosus.

### **Laboratory Tests**

Because serious GI tract ulcerations and bleeding can occur without warning symptoms, physicians should monitor for signs or symptoms of GI bleeding. Patients on long-term treatment with NSAIDs, should have their CBC and a chemistry profile checked periodically. If clinical signs and symptoms consistent with liver or renal disease develop, systemic manifestations occur (e.g., eosinophilia, rash, etc.) or if abnormal liver tests persist or worsen, etodolac extended-release tablets should be discontinued.

### **Drug Interactions**

#### ***ACE-inhibitors***

Reports suggest that NSAIDs may diminish the antihypertensive effect of ACE-inhibitors. This interaction should be given consideration in patients taking NSAIDs concomitantly with ACE-inhibitors.

#### ***Aspirin***

When etodolac extended-release tablets are administered with aspirin, its protein binding is reduced, although the clearance of free etodolac is not altered. The clinical significance of this interaction is not known; however, as with other NSAIDs, concomitant administration of etodolac extended-release tablets and aspirin is not generally recommended because of the potential of increased adverse effects.

#### ***Cyclosporine and Digoxin***

Etodolac extended-release tablets, like other NSAIDs, through effects on renal prostaglandins, may cause changes in the elimination of these drugs, leading to elevated serum levels of cyclosporine and digoxin and increased toxicity. Nephrotoxicity associated with cyclosporine may also be enhanced. Patients receiving these drugs who are given etodolac extended-release tablets, or any other NSAID, and particularly those patients with altered renal function, should be observed for the development of the specific toxicities of these drugs.

#### ***Furosemide***

Clinical studies, as well as post marketing observations, have shown that etodolac extended-release tablets can reduce the natriuretic effect of furosemide and thiazides in some patients. This response has been attributed to inhibition of renal prostaglandin synthesis. During concomitant therapy with NSAIDs, the patient should be observed closely for signs of renal failure (see **WARNINGS, Renal Effects**), as well as to assure diuretic efficacy.

#### ***Lithium***

NSAIDs have produced an elevation of plasma lithium levels and a reduction in renal lithium clearance. The mean minimum lithium concentration increased 15% and the renal clearance was decreased by approximately 20%. These effects have been attributed to inhibition of renal prostaglandin synthesis by the NSAID. Thus, when NSAIDs and lithium are administered concurrently, subjects should be observed carefully for signs of lithium toxicity.

#### ***Methotrexate***

NSAIDs have been reported to competitively inhibit methotrexate accumulation in rabbit kidney slices.

This may indicate that they could enhance the toxicity of methotrexate. Caution should be used when NSAIDs are administered concomitantly with methotrexate.

### **Phenylbutazone**

Phenylbutazone causes an increase (by about 80%) in the free fraction of etodolac. Although *in vivo* studies have not been done to see if etodolac clearance is changed by coadministration of phenylbutazone, it is not recommended that they be coadministered.

### **Warfarin**

The effects of warfarin and NSAIDs on GI bleeding are synergistic, such that users of both drugs together have a risk of serious GI bleeding higher than users of either drug alone.

### **Drug/Laboratory Test Interactions**

The urine of patients who take etodolac can give a false-positive reaction for urinary bilirubin (urobilin) due to the presence of phenolic metabolites of etodolac. Diagnostic dip-stick methodology, used to detect ketone bodies in urine, has resulted in false-positive findings in some patients treated with etodolac. Generally, this phenomenon has not been associated with other clinically significant events. No dose relationship has been observed.

Etodolac treatment is associated with a small decrease in serum uric acid levels. In clinical trials, mean decreases of 1 to 2 mg/dL were observed in arthritic patients receiving etodolac (600 mg to 1000 mg/day) after 4 weeks of therapy. These levels then remained stable for up to 1 year of therapy.

### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

No carcinogenic effect of etodolac was observed in mice or rats receiving oral doses of 15 mg/kg/day (45 to 89 mg/m<sup>2</sup>, respectively) or less for periods of 18 months or 2 years, respectively. Etodolac was not mutagenic in *in vitro* tests performed with *S. typhimurium* and mouse lymphoma cells as well as in an *in vivo* mouse micronucleus test. However, data from the *in vitro* human peripheral lymphocyte test showed an increase in the number of gaps (3% to 5% unstained regions in the chromatid without dislocation) among the etodolac-treated cultures (50 to 200 g/mL) compared to negative controls (2%); no other difference was noted between the controls and drug-treated groups. Etodolac showed no impairment of fertility in male and female rats up to oral doses of 16 mg/kg (94 mg/m<sup>2</sup>). However, reduced implantation of fertilized eggs occurred in the 8 mg/kg group.

### **Pregnancy**

#### **Teratogenic Effects**

##### *Pregnancy category C*

Reproductive studies conducted in rats and rabbits have not demonstrated evidence of developmental abnormalities. However, animal reproduction studies are not always predictive of human response. There are no adequate and well-controlled studies in pregnant women.

#### **Nonteratogenic Effects**

Because of the known effects of nonsteroidal anti-inflammatory drugs on the fetal cardiovascular system (closure of ductus arteriosus), use during pregnancy (particularly late pregnancy) should be avoided.

#### **Labor and Delivery**

In rat studies with NSAIDs, as with other drugs known to inhibit prostaglandin synthesis, an increased incidence of dystocia, delayed parturition, and decreased pup survival occurred. The effects of etodolac extended-release tablets on labor and delivery in pregnant women are unknown.

#### **Nursing Mothers**

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk and because of the potential for serious adverse reactions in nursing infants from etodolac extended-release tablets, a decision should be made whether to discontinue nursing or to discontinue

the drug, taking into account the importance of the drug to the mother.

### **Pediatric Use**

Safety and effectiveness in pediatric patients below the age of 6 years have not been established.

### **Geriatric Use**

As with any NSAIDs, caution should be exercised in treating the elderly (65 years and older).

### **ADVERSE REACTIONS**

A total of 1552 patients were exposed to etodolac extended-release tablets in controlled clinical studies of at least 4 weeks in length and using daily doses in the range of 400 to 1200 mg. In the tabulations below, adverse event rates are generally categorized based on the incidence of events in the first 30 days of treatment with etodolac extended-release tablets. As with other NSAIDs, the cumulative adverse event rates may increase significantly over time with extended therapy.

In patients taking NSAIDs, including etodolac extended-release tablets, the most frequently reported adverse experiences occurring in approximately 1 to 10% of patients are:

Gastrointestinal experiences including

abdominal pain dyspepsia gross bleeding/perforation*	constipation flatulence nausea	Diarrhea GI ulcers (gastric/duodenal)* Vomiting
--	--------------------------------------	---

Other events including:

abnormal renal function* dizziness headaches infection rashes	Anemia* edema* hypertension pharyngitis rhinitis	asthenia elevated liver enzymes* increased bleeding time* pruritus tinnitus*
---	--	--

\* Adverse events that were observed in < 1% of patients in the first 30 days of treatment with etodolac extended-release tablets in clinical trials.

### **Additional NSAID Adverse Experiences Reported Occasionally with NSAIDs or Etodolac Extended-release Tablets Include**

#### ***Body as a whole***

Allergic reaction, anaphylactic/anaphylactoid reactions (including shock), chills, fever, sepsis

#### ***Cardiovascular system***

Congestive heart failure, flushing, palpitations, tachycardia, syncope, vasculitis (including necrotizing and allergic)

#### ***Digestive system***

Anorexia, cholestatic hepatitis, cholestatic jaundice, dry mouth, duodenitis, eructation, esophagitis, gastritis, gastric/peptic ulcers, glossitis, hepatic failure, hepatitis, hematemesis, intestinal ulceration, jaundice, liver necrosis, melena, pancreatitis, rectal bleeding, stomatitis

#### ***Hemic and lymphatic system***

Agranulocytosis, ecchymosis, eosinophilia, hemolytic anemia, leukopenia, neutropenia, pancytopenia, purpura, thrombocytopenia

#### ***Metabolic and nutritional***

Hyperglycemia in previously controlled diabetic patients

***Nervous system***

Anxiety, confusion, depression, dream abnormalities, insomnia, nervousness, paresthesia, somnolence, tremors, vertigo

***Respiratory system***

Asthma, dyspnea, pulmonary infiltration with eosinophilia

***Skin and appendages***

Angioedema, cutaneous vasculitis with purpura, erythema multiforme, hyperpigmentation, sweating, urticaria, vesiculobullous rash

***Special senses***

Blurred vision, photophobia, transient visual disturbances

***Urogenital system***

Dysuria, elevated BUN, oliguria/polyuria, proteinuria, renal failure, renal insufficiency, renal papillary necrosis, serum creatinine increase, urinary frequency

**Other NSAID Adverse Reactions, Which Occur Rarely Are**

***Body as a whole***

Anaphylactic reactions, appetite changes, death

***Cardiovascular system***

Arrhythmia, cerebrovascular accident, hypotension, myocardial infarction

***Digestive system***

Colitis, esophagitis with or without stricture or cardiospasm, thirst, ulcerative stomatitis

***Hemic and lymphatic system***

Aplastic anemia, lymphadenopathy

***Metabolic and nutritional***

Change in weight

***Nervous system***

Coma, convulsions, hallucinations, meningitis

***Respiratory***

Bronchitis, pneumonia, respiratory depression, sinusitis

***Skin and appendages***

Alopecia, exfoliative dermatitis, maculopapular rash, photosensitivity, skin peeling, Stevens-Johnson syndrome, toxic epidermal necrosis

***Special senses***

Conjunctivitis, deafness, hearing impairment, taste perversion

***Urogenital system***

Cystitis, hematuria, interstitial nephritis, leukorrhea, renal calculus, uterine bleeding irregularities

**OVERDOSAGE**

Symptoms following acute NSAID overdose are usually limited to lethargy, drowsiness, nausea, vomiting, and epigastric pain, which are generally reversible with supportive care. Gastrointestinal bleeding can occur. Hypertension, acute renal failure, respiratory depression and coma may occur, but are rare. Anaphylactoid reactions have been reported with therapeutic ingestion of NSAIDs, and may occur following an overdose.

Patients should be managed by symptomatic and supportive care following an NSAID overdose. There are no specific antidotes.

Emesis and/or activated charcoal (60 to 100 g in adults, 1 to 2 g/kg in children) and/or osmotic cathartic may be indicated in patients seen within 4 hours of ingestion with symptoms or following a large overdose (5 to 10 times the usual dose). Forced diuresis, alkalinization of the urine, hemodialysis, or hemoperfusion may not be useful due to high protein binding.

## **DOSAGE AND ADMINISTRATION**

Carefully consider the potential benefits and risks of etodolac extended-release tablets and other treatment options before deciding to use etodolac extended-release tablets. Use the lowest effective dose for the shortest duration consistent with individual patient treatment goals (see **WARNINGS**).

After observing the response to initial therapy with etodolac extended-release tablets, the dose and frequency should be adjusted to suit an individual patient's needs.

### **Juvenile Rheumatoid Arthritis**

For the relief of the signs and symptoms of juvenile rheumatoid arthritis in patients 6 to 16 years of age, the recommended dose given orally once per day should be based on body weight, according to the following table:

**Table 4**

<b>Body Weight Range (kg)</b>	<b>Dose</b>
20 to 30	400 mg Tablet x 1
31 to 45	600 mg Tablet x 1
46 to 60	400 mg Tablet x 2
> 60	500 mg Tablet x 2

### **Rheumatoid Arthritis and Osteoarthritis**

For the relief of the signs and symptoms of osteoarthritis or rheumatoid arthritis, the recommended starting dose of etodolac extended-release tablets is 400 to 1000 mg given orally once per day.

As with other NSAIDs, the lowest effective dose should be sought for each patient. In chronic conditions, a therapeutic response to therapy with etodolac extended-release tablets is sometimes seen within one week of therapy, but most often is observed by two weeks.

## **HOW SUPPLIED**

Etodolac Extended-release Tablets USP, 400 mg are orange-colored, oval-shaped, beveled edged, film-coated tablets, debossed with "271" on one side and plain on other side and are supplied as follows:

NDC 68382-271-14 in bottles of 60 tablets

NDC 68382-271-01 in bottles of 100 tablets

NDC 68382-271-05 in bottles of 500 tablets

NDC 68382-271-10 in bottles of 1000 tablets

NDC 68382-271-77 in unit-dose blister cartons of 100 (10 x 10) unit dose tablets

Etodolac Extended-release Tablets USP, 500 mg are grey-colored, oval-shaped, beveled edged, film-coated tablets, debossed with "272" on one side and plain on other side and are supplied as follows:

NDC 68382-272-14 in bottles of 60 tablets

NDC 68382-272-01 in bottles of 100 tablets

NDC 68382-272-05 in bottles of 500 tablets

NDC 68382-272-10 in bottles of 1000 tablets

NDC 68382-272-77 in unit-dose blister cartons of 100 (10 x 10) unit dose tablets

Etodolac Extended-release Tablets USP, 600 mg are blue-colored, oval-shaped, beveled edged, film-coated tablets debossed with "273" on one side and plain on other side and are supplied as follows:

NDC 68382-273-14 in bottles of 60 tablets

NDC 68382-273-01 in bottles of 100 tablets

NDC 68382-273-05 in bottles of 500 tablets

NDC 68382-273-10 in bottles of 1000 tablets

NDC 68382-273-77 in unit-dose blister cartons of 100 (10 x 10) unit dose tablets

**Storage:**

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Protect from excessive heat and humidity.

Dispense in a tight, light-resistant container with a child-resistant closure.

**Manufactured by:**

Cadila Healthcare Ltd.

Ahmedabad, India

**Distributed by:**

**Zydus Pharmaceuticals USA Inc.**

Pennington, NJ 08534

Rev.: 06/16

**Medication Guide**

**For Nonsteroidal Anti-inflammatory Drugs (NSAIDs)**

**Etodolac**

**(e-TOE-doe-lak)**

**Extended-release Tablets, USP**

**What is the most important information I should know about medicines called Nonsteroidal Anti-inflammatory Drugs (NSAIDs)?**

**NSAIDs can cause serious side effects, including:**

- **Increased risk of a heart attack or stroke that can lead to death** . This risk may happen early in treatment and may increase:
  - with increasing doses of NSAIDs
  - with longer use of NSAIDs

**Do not take NSAIDs right before or after a heart surgery called a "coronary artery bypass graft (CABG)." Avoid taking NSAIDs after a recent heart attack, unless your healthcare provider tells you to. You may have an increased risk of another heart attack if you take NSAIDs after a recent heart attack.**

- **Increased risk of bleeding, ulcers, and tears (perforation) of the esophagus (tube leading from the mouth to the stomach), stomach and intestines:**

- anytime during use
- without warning symptoms
- that may cause death

**The risk of getting an ulcer or bleeding increases with:**

- past history of stomach ulcers, or stomach or intestinal bleeding with use of NSAIDs
- taking medicines called "corticosteroids", "anticoagulants", "SSRIs", or "SNRIs"
- increasing doses of NSAIDs
- longer use of NSAIDs
- smoking
- drinking alcohol
- older age
- poor health
- advanced liver disease
- bleeding problems

**NSAIDs should only be used:**

- exactly as prescribed
- at the lowest dose possible for your treatment
- for the shortest time needed

**What are NSAIDs?**

NSAIDs are used to treat pain and redness, swelling, and heat (inflammation) from medical conditions such as different types of arthritis, menstrual cramps, and other types of short-term pain.

**Who should not take NSAIDs? Do not take NSAIDs:**

- if you have had an asthma attack, hives, or other allergic reaction with aspirin or any other NSAIDs.
- right before or after heart bypass surgery.

**Before taking NSAIDs, tell your healthcare provider about all of your medical conditions, including if you:**

- have liver or kidney problems
- have high blood pressure
- have asthma
- are pregnant or plan to become pregnant. Talk to your healthcare provider if you are considering taking NSAIDs during pregnancy. **You should not take NSAIDs after 29 weeks of pregnancy.**
- are breastfeeding or plan to breast feed.

**Tell your healthcare provider about all of the medicines you take, including prescription or over-the-counter medicines, vitamins or herbal supplements.** NSAIDs and some other medicines can interact with each other and cause serious side effects. **Do not start taking any new medicine without talking to your healthcare provider first.**

**What are the possible side effects of NSAIDs?**

**NSAIDs can cause serious side effects, including:**

**See "What is the most important information I should know about medicines called Nonsteroidal Anti-inflammatory Drugs (NSAIDs)?"**

- new or worse high blood pressure
- heart failure
- liver problems including liver failure
- kidney problems including kidney failure
- low red blood cells (anemia)
- life-threatening skin reactions
- life threatening allergic reactions
- **Other side effects of NSAIDs include:** stomach pain, constipation, diarrhea, gas, heartburn, nausea, vomiting, and dizziness.

**Get emergency help right away if you get any of the following symptoms :**

- shortness of breath or trouble breathing
- chest pain
- weakness in one part or side of your body
- slurred speech
- swelling of the face or throat

**Stop taking your NSAID and call your healthcare provider right away if you get any of the following symptoms :**

- nausea
- more tired or weaker than usual
- diarrhea
- itching
- your skin or eyes look yellow
- indigestion or stomach pain
- flu-like symptoms
- vomit blood
- there is blood in your bowel movement or it is black and sticky like tar
- unusual weight gain
- skin rash or blisters with fever
- swelling of the arms, legs, hands and feet

**If you take too much of your NSAID, call your healthcare provider or get medical help right away.**

These are not all the possible side effects of NSAIDs. For more information, ask your healthcare provider or pharmacist about NSAIDs.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

**Other information about NSAIDs**

- Aspirin is an NSAID but it does not increase the chance of a heart attack. Aspirin can cause bleeding in the brain, stomach, and intestines. Aspirin can also cause ulcers in the stomach and intestines.
- Some NSAIDs are sold in lower doses without a prescription (over-the counter). Talk to your healthcare provider before using over-the-counter NSAIDs for more than 10 days.

**General information about the safe and effective use of NSAIDs**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use NSAIDs for a condition for which it was not prescribed. Do not give NSAIDs to other people, even if they have the same symptoms that you have. It may harm them.

If you would like more information about NSAIDs, talk with your healthcare provider. You can ask your pharmacist or healthcare provider for information about NSAIDs that is written for health professionals.

Please address medical inquiries to, (MedicalAffairs@zydususa.com) Tel.: 1-877-993-8779.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

**This product's package insert may have been updated. For current package insert, please visit [www.zydususa.com](http://www.zydususa.com)**

**Manufactured by:**

Cadila Healthcare Ltd.  
Ahmedabad, India

**Distributed by:**

**Zydus Pharmaceuticals USA Inc.**  
Pennington, NJ 08534



Rev.: 01/16

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**

NDC 68382-271-14 in bottle of 60 tablets

Etodolac Extended-release Tablets USP, 400 mg

R<sub>x</sub> only

60 tablets

ZYDUS

No Varnished Area (Do Not Print)  
(18 x 41 mm)

NDC 68382-271-14

**Etodolac  
Extended-release  
Tablets, USP**

**400 mg**

PHARMACIST: Dispense the Medication Guide  
Provided Separately to each Patient.

**zydus**  
pharmaceuticals

**60 TABLETS  
Rx only**

Each extended-release tablet contains:  
Etodolac, USP .....400 mg

**Usual Dosage:** See package insert for  
complete prescribing information.

Store at 20° to 25°C (68° to 77°F)  
[See USP Controlled Room Temperature].  
Protect from excessive heat and humidity.

Dispense in a tight, light-resistant  
container with a child-resistant closure.

**KEEP THIS AND ALL DRUGS OUT  
OF THE REACH OF CHILDREN.**

Code No.: GUJ/DRUG/1486

**Manufactured by:**  
Cadila Healthcare Ltd.  
Ahmedabad, India

**Distributed by:**  
Zydus Pharmaceuticals (USA) Inc.  
Pennington, NJ 08534

2055925  
Rev: 04/17

NDC 68382-272-14 in bottle of 60 tablets

Etodolac Extended-release Tablets USP, 500 mg

R<sub>x</sub> only

60 tablets

ZYDUS

No Varnished Area (Do Not Print)  
(18 x 50.5 mm)

NDC 68382-272-14

# Etodolac Extended-release Tablets, USP

**500 mg**

PHARMACIST: Dispense the Medication Guide  
Provided Separately to each Patient.

60 TABLETS  
Rx only

Each extended-release tablet contains:  
Etodolac, USP .....500 mg

**Usual Dosage:** See package insert for  
complete prescribing information.

Store at 20° to 25°C (68° to 77°F)  
[See USP Controlled Room Temperature].  
Protect from excessive heat and humidity.

Dispense in a tight, light-resistant  
container with a child-resistant closure.

**KEEP THIS AND ALL DRUGS OUT  
OF THE REACH OF CHILDREN.**

Code No.: GUJ/DRUG/1486

**Manufactured by:**  
Cadila Healthcare Ltd.  
Ahmedabad, India

**Distributed by:**  
Zydus Pharmaceuticals (USA) Inc.  
Pennington, NJ 08534

2055927  
Rev.: 04/17

NDC 68382-273-14 in bottle of 60 tablets  
Etodolac Extended-release Tablets USP, 600 mg  
Rx only  
60 tablets  
ZYDUS

No Varnished Area (Do Not Print)  
(18 x 50.5 mm)

NDC 68382-273-14

# Etodolac Extended-release Tablets, USP

**600 mg**

PHARMACIST: Dispense the Medication Guide  
Provided Separately to each Patient.

60 TABLETS  
Rx only

Each extended-release tablet contains:  
Etodolac, USP .....600 mg

**Usual Dosage:** See package insert for  
complete prescribing information.

Store at 20° to 25°C (68° to 77°F)  
[See USP Controlled Room Temperature].  
Protect from excessive heat and humidity.

Dispense in a tight, light-resistant  
container with a child-resistant closure.

**KEEP THIS AND ALL DRUGS OUT  
OF THE REACH OF CHILDREN.**

Code No.: GUJ/DRUG/1486

**Manufactured by:**  
Cadila Healthcare Ltd.  
Ahmedabad, India

**Distributed by:**  
Zydus Pharmaceuticals (USA) Inc.  
Pennington, NJ 08534

2055929  
Rev.: 04/17

# ETODOLAC

etodolac tablet, film coated, extended release

## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:68382-271
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ETODOLAC (UNII: 2M36281008) (ETODOLAC - UNII:2M36281008)	ETODOLAC	400 mg

## Inactive Ingredients

Ingredient Name	Strength
SODIUM PHOSPHATE, DIBASIC, ANHYDROUS (UNII: 22ADO53M6F)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
ETHYLCELLULOSE (100 MPA.S) (UNII: 47MLB0F1MV)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

## Product Characteristics

<b>Color</b>	ORANGE (ORANGE)	<b>Score</b>	no score
<b>Shape</b>	OVAL (OVAL)	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	271
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68382-271-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/15/2014	
2	NDC:68382-271-05	500 in 1 BOTTLE; Type 0: Not a Combination Product	02/15/2014	
3	NDC:68382-271-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	02/15/2014	
4	NDC:68382-271-77	100 in 1 CARTON	02/15/2014	
4	NDC:68382-271-30	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
5	NDC:68382-271-14	60 in 1 BOTTLE; Type 0: Not a Combination Product	02/15/2014	

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA091134	02/15/2014	

# ETODOLAC

etodolac tablet, film coated, extended release

## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:68382-272
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ETODOLAC (UNII: 2M36281008) (ETODOLAC - UNII:2M36281008)	ETODOLAC	500 mg

## Inactive Ingredients

Ingredient Name	Strength
SODIUM PHOSPHATE, DIBASIC, ANHYDROUS (UNII: 22ADO53M6F)	
ETHYLCELLULOSE (100 MPAS) (UNII: 47MLB0F1MV)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
TALC (UNII: 7SEV7J4RIU)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

## Product Characteristics

<b>Color</b>	GRAY (GRAY)	<b>Score</b>	no score
<b>Shape</b>	OVAL (OVAL)	<b>Size</b>	18mm
<b>Flavor</b>		<b>Imprint Code</b>	272
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68382-272-05	500 in 1 BOTTLE; Type 0: Not a Combination Product	02/15/2014	
2	NDC:68382-272-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	02/15/2014	
3	NDC:68382-272-77	100 in 1 CARTON	02/15/2014	
3	NDC:68382-272-30	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
4	NDC:68382-272-14	60 in 1 BOTTLE; Type 0: Not a Combination Product	02/15/2014	
5	NDC:68382-272-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/15/2014	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091134	02/15/2014	

# ETODOLAC

etodolac tablet, film coated, extended release

## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:68382-273
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ETODOLAC (UNII: 2M36281008) (ETODOLAC - UNII:2M36281008)	ETODOLAC	600 mg

## Inactive Ingredients

Ingredient Name	Strength
SODIUM PHOSPHATE, DIBASIC, ANHYDROUS (UNII: 22ADO53M6F)	
ETHYLCELLULOSE (100 MPAS) (UNII: 47MLB0F1MV)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

## Product Characteristics

<b>Color</b>	BLUE (BLUE)	<b>Score</b>	no score
<b>Shape</b>	OVAL (OVAL)	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	273
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68382-273-14	60 in 1 BOTTLE; Type 0: Not a Combination Product	02/15/2014	
2	NDC:68382-273-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/15/2014	
3	NDC:68382-273-05	500 in 1 BOTTLE; Type 0: Not a Combination Product	02/15/2014	
4	NDC:68382-273-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	02/15/2014	
5	NDC:68382-273-77	100 in 1 CARTON	02/15/2014	
5	NDC:68382-273-30	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091134	02/15/2014	

**Labeler** - Zydus Pharmaceuticals (USA) Inc. (156861945)

**Registrant** - Zydus Pharmaceuticals (USA) Inc. (156861945)

**Establishment**

Name	Address	ID/FEI	Business Operations
Cadila Healthcare Limited		918596198	ANALYSIS(68382-271, 68382-272, 68382-273) , MANUFACTURE(68382-271, 68382-272, 68382-273)

Revised: 1/2020

Zydus Pharmaceuticals (USA) Inc.