

TOPCARE ALL DAY PAIN RELIEF- naproxen sodium tablet, film coated
Topco Associates LLC

Topco Associates LLC. All Day Pain Relief Drug Facts

Active ingredient (in each tablet)

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts more than 10 days

- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose
- **Adults and children 12 years and older:**
 - take 1 tablet every 8 to 12 hours while symptoms last
 - for the first dose you may take 2 tablets within the first hour
 - do not exceed 2 tablets in any 8- to 12-hour period
 - do not exceed 3 tablets in a 24-hour period
- **Children under 12 years:**
 - ask a doctor

Other information

- **each tablet contains:** sodium 22 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients

FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments?

1-888-423-0139

Principal Display Panel

COMPARE TO ALEVE® ACTIVE INGREDIENT
 All Day Pain Relief
 NAPROXEN SODIUM TABLETS, 220 mg EACH
 PAIN RELIEVER • FEVER REDUCER (NSAID)
 12 HOUR
 Strength to last 12 hours
 50 TABLETS
 actual size



DISTRIBUTED BY TOPCO ASSOCIATES LLC
 ELK GROVE VILLAGE, IL 60007
 ©TOPCO PERA0321
 QUESTIONS? 1-888-423-0139
 topcare@topco.com
 www.topcarebrand.com



Scan here for more information or call 1-888-423-0139

GLUTEN FREE

✓ QUALITY GUARANTEED

TopCare
 health

COMPARE TO ALEVE®
 ACTIVE INGREDIENT**

NDC 36800-490-71

All Day Pain Relief

NAPROXEN SODIUM TABLETS,
 220 mg EACH

PAIN RELIEVER • FEVER REDUCER (NSAID)

12 HOUR Strength to last 12 hours

50 TABLETS

actual size

KEEP CARTON FOR REFERENCE

**This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Aleve®.



49071 88 C14

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are under a doctor's care for any serious condition taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin taking any other drug

When using this product

take with food or milk if stomach upset occurs

Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: feel faint vomit blood have bloody or black stools have stomach pain that does not get better you have symptoms of heart problems or stroke: chest pain trouble breathing weakness in one part or side of body slurred speech leg swelling pain gets worse or lasts more than 10 days fever gets worse or lasts more than 3 days you have difficulty swallowing it feels like the pill is stuck in your throat redness or swelling is present in the painful area any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions do not take more than directed

the smallest effective dose should be used
 drink a full glass of water with each dose
Adults and children 12 years and older:
 take 1 tablet every 8 to 12 hours while symptoms last
 for the first dose you may take 2 tablets within the first hour
 do not exceed 2 tablets in any 8- to 12-hour period
 do not exceed 3 tablets in a 24-hour period
Children under 12 years: ask a doctor

Other information each tablet contains: sodium 22 mg
 store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Drug Facts (continued)

Inactive ingredients FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments? 1-888-423-0139

actual size



DO NOT USE IF PRINTED FOIL UNDER CAP IS BROKEN OR MISSING

Drug Facts

Active ingredient (in each tablet)	Purpose
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*	Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug	

Uses temporarily relieves minor aches and pains due to: minor pain of arthritis muscular aches backache menstrual cramps headache toothache the common cold temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives facial swelling asthma (wheezing) shock skin reddening rash blisters If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older have had stomach ulcers or bleeding problems take a blood thinning (anticoagulant) or steroid drug take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) have 3 or more alcoholic drinks every day while using this product take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer right before or after heart surgery

Ask a doctor before use if the stomach bleeding warning applies to you you have a history of stomach problems, such as heartburn you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke you are taking a diuretic you have problems or serious side effects from taking pain relievers or fever reducers

TOPCARE ALL DAY PAIN RELIEF

naproxen sodium tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-490
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	BLUE (Light Blue)	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	L490
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-490-62	1 in 1 CARTON	07/22/1997	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:36800-490-71	1 in 1 CARTON	01/14/1997	
2		50 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:36800-490-78	1 in 1 CARTON	06/11/2002	12/31/2023
3		100 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:36800-490-76	1 in 1 CARTON	04/07/2009	

4		120 in 1 BOTTLE; Type 0: Not a Combination Product		
5	NDC:36800-490-75	1 in 1 CARTON	12/09/2021	
5		90 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074661	01/14/1997	

Labeler - Topco Associates LLC (006935977)

Revised: 12/2023

Topco Associates LLC