BIO-COX TYPE A MEDICATED ARTICLE- salinomycin sodium powder Huvepharma, Inc.

Bio-Cox®

Type A Medicated Article

Brand of Salinomycin Sodium
Type A Medicated Article

Net wt 50 LB (22.68 kg)

HUVEPHARMA®

Bio-Cox® Type A Medicated Article

Brand of Salinomycin Sodium
Type A Medicated Article

For use in broiler chickens, replacement chickens, and quail only

Active drug ingredient

Salinomycin equivalent to 60 grams salinomycin sodium activity per pound

For use in manufacturing medicated feeds only

Broiler chickens and replacement chickens

Use Bio-Cox Type A Medicated Article to provide salinomycin sodium concentrations ranging from 40-60 g/ton (0.0044% to 0.0066%) for the prevention of coccidiosis in broiler chickens and replacement chickens caused by *Eimeria tenella*, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti* and *E. mivati*. The dosage should be adjusted to meet the severity of the coccidial challenge, which varies with environmental and managment conditions.

Quail

Use Bio-Cox Type A Medicated Article to provide salinomycin sodium concentrations of 50 g/ton (0.0055%) for the prevention of coccidiosis in quail caused by *Eimeria dispersa* and *E. lettyae*.

Mixing directions

Thoroughly mix Bio-Cox Type A Medicated Article with non-medicated feed to provide the level of salinomycin sodium per ton as indicated in the table below.

Salinomycin Sodium Activity	Amount of E Type A Medi Per Ton of F	cated Article (60 g/lb)
(grams per ton)	lb	grams
40	0.67	304
45	0.75	341

50	0.83	377
55	0.92	418
60	1.00	454

Feeding directions

Feed continuously as the sole ration

Important

Must be thoroughly mixed in feeds before use. Not approved for use with pellet binders.

Caution

May be fatal if accidentally fed to adult turkeys or to horses.

Warning

Do not feed to birds producing eggs for human consumption. When mixing and handling Bio-Cox Type A Medicated Article premix, use protective clothing, impervious gloves, eye protection, and an approved dust respirator. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water.

Store below 25°C (77°F), excursions permitted to 37°C (99°F) Restricted drug (California) - Use only as directed. Not for human use. **Approved by FDA under NADA # 128-686**

HUVEPHARMA®

Distributed by: Huvepharma, Inc. Peachtree City, GA 30269

HUVEPHARMA and Bio-Cox are registered trademarks of Huvepharma EOOD.

Made in USA

Net wt 50 LB (22.68 kg)



Bio-Cox® Type A Medicated Article

Bio-Cox® Type A Medicated Articl

Brand of Salinomycin Sodium
Type A Medicated Article

Net wt 50 LB (22.68 kg)



Bio-Cox[®]
Type A Medicated Article

Bio-Cox® Type A Medicated Article Brand of Salinomycin Sodium Type A Medicated Article For use in broiler chickens, replacement chickens, and quail only Active drug Salinomycin equivalent to 60 grams salinomycin sodium activity per pound Ingredient For use in manufacturing medicated feeds only Use Bio-Cox Type A Medicated Article to provide salinomycin sodium concentrations ranging from 40-60 g/ton (0.0044% to 0.0066%) for the prevention of ococidicals in broiler chickens and replacement chickens caused by Emeria tensella E. necatric, E. acenvulina, E. maxima, E. brunetti and E. mixati. The dosage-should be adjusted to meet the severity of the cocidial challenge, which varies with environmental and managment conditions. Broiler chickens and replacement chickens Use Bio-Cox Type A Medicated Article to provide salinomycin sodium concentrations of 50 g/ton (0.0055%) for the prevention of coccidiosis in quail caused by *Eimaria dispersa and E. lettyae*. Quall Mixing directions Amount of Bio-Cox Type A Medicated Article (60 g/lb) Per Ton of Feed Salinomycin Sodium Activity (grams per ton) 0.67 0.75 0.83 0.92 304 341 377 418 Feeding Feed continuously as the sole ration directions Important Must be thoroughly mixed inteeds before use. Not approved for use with pellet binders. Caution May be fatal if accidentally fed to adult turkeys or to horses Do not feed to birds producing eggs for human consumption. When mixing and handling Bio-Cox Type A Medicated Article premix, use protective clathing, impervious gloves, eye protection, and an approved dust respirator. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rings thoroughly with water. Warning Store below 25°C (77°F), excursions permitted to 37°C (98°F) Restricted drug (Californis) *Use only as directed. Not for human use. Approved by FDA under NADA # 128-886 HUVEPHARMA Take Time Made in USA Distributed by: Huvepharms, Inc. Peachtree City, GA 30289 Observe Label HUVEPHAPMA and Bis-Cox are registered frederierts of Herepterra. 5000. Net wt 50 LB (22.68 kg) GUARANTEED BY 8814 13-000462 Polyamide

salinomycin sodium powder

Product Information			
Product Type	OTC TYPE A MEDICATED ARTICLE ANIMAL DRUG	Item Code (Source)	NDC:23243- 6807
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
SALINOMYCIN SODIUM (UNII: 92UOD3BMEK) (SALINOMYCIN - UNII:62UXS86T64)	SALINOMYCIN SODIUM	60 g in 0.45 kg

P	ackaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:23243-6807-5	22.68 kg in 1 BAG		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA128686	01/01/2020	

Labeler - Huvepharma, Inc. (619153559)

Registrant - Huvepharma EOOD (552671651)

Establishment			
Name	Address	ID/FEI	Business Operations
Huvepharma, Inc - Van Buren		080346352	medicated animal feed manufacture, analysis, pack, label, manufacture

Revised: 9/2023 Huvepharma, Inc.