

NICOTINE- nicotine polacrilex lozenge
CVS Pharmacy

Drug Facts

Active ingredient (in each lozenge)

Nicotine polacrilex 4 mg

Purpose

Stop smoking aid

Use

- reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding,

only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth problems occur
- persistent indigestion or severe sore throat occurs
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness,

diarrhea, weakness and rapid heartbeat

- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets.

Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222)

Directions (4 mg lozenge)

- **if you are under 18 years of age, ask a doctor before use. No studies have been done to show if this product will work for you.**
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the lozenge on your quit day
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2 mg nicotine lozenge
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine lozenge according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours

- **nicotine lozenge is a medicine and must be used a certain way to get the best results**
- place the lozenge in your mouth and allow the lozenge to slowly dissolve. Minimize swallowing. **Do not chew or swallow lozenge.**
- you may feel a warm or tingling sensation
- occasionally move the lozenge from one side of your mouth to the other until completely dissolved
- do not eat or drink 15 minutes before using or while the lozenge is in your mouth
- to improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks
- do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you hiccups, heartburn, nausea or other side effects
- **do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day.**
- It is important to complete treatment. If you feel you need to use the lozenge for a longer period to keep from smoking, talk to your health care provider.

Other information

- Phenylketonurics: Contains Phenylalanine 2.8 mg per lozenge
- store at 20-25°C (68-77°F)
- keep vial tightly closed and protect from light

Inactive ingredients

aspartame, calcium polycarbophil, flavor, maltodextrin, mannitol, sodium alginate, sodium bicarbonate, sodium stearyl fumarate, talc, xanthan gum

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel (4 mg lozenge)

Compare to the active ingredient in Nicorette® Lozenge*

Sugar Free

Nicotine Lozenges

Nicotine Polacrilex 4 mg

STOP SMOKING AID

Mint Flavor

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP.

If you smoke your first cigarette **MORE THAN 30 MINUTES** after waking up, use Nicotine Polacrilex 2 mg Lozenge.

LOZENGES, 4 mg

TO INCREASE YOUR SUCCESS IN QUITTING:

- 1.** You must be motivated to quit.
- 2. Use Enough** - Use **at least 9** Nicotine mini lozenges per day during the first six weeks.
- 3. Use Long Enough** - Use Nicotine mini lozenges for the full 12 weeks.
- 4. Use With a Support Program** as directed in the enclosed User's Guide.
 - **not for sale to those under 18 years of age**
 - **proof of age required**
 - **not for sale in vending machines or from any source where proof of age cannot be verified**

*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., distributor of Nicorette® Lozenge.

See inside for more details or visit www.nitthehabit.com

TAMPER EVIDENT FEATURE: DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING.

RETAIN OUTER CARTON FOR FULL PRODUCT USES, DIRECTIONS AND WARNINGS.

Distributed By: CVS Pharmacy, Inc.

One CVS Drive, Woonsocket, RI 02895

Package Label (4 mg lozenge)

Drug Facts

Active ingredient (in each lozenge)
Nicotine polacrilex 4 mg

Purpose
Stop smoking aid

Use
reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have
heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
stomach ulcer or diabetes history of seizures

Ask a doctor or pharmacist before use if you are
taking a non-nicotine stop smoking drug
taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if
mouth problems occur
persistent indigestive or severe sore throat occurs
irregular heartbeat or palpitations occur
you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions
If you are under 18 years of age, ask a doctor before use. No studies have been done to show if this product will work for you.
before using this product, read the enclosed User's Guide for complete directions and other important information
begin using the lozenge on your quit day
if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine lozenges
if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine lozenge according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours

nicotine lozenge is a medicine and must be used a certain way to get the best results
place the lozenge in your mouth and allow the lozenge to slowly dissolve. Minimize swallowing. Do not chew or swallow lozenges.
you may feel a warm or tingling sensation

Drug Facts (continued)

occasionally move the lozenge from one side of your mouth to the other until completely dissolved
do not eat or drink 15 minutes before using or while the lozenge is in your mouth
to improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks
do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you hiccups, heartburn, nausea or other side effects
do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day.
it is important to complete treatment. If you feel you need to use the lozenge for a longer period to keep from smoking, talk to your health care provider.

Other information
Phenylethanolamine: Contains Phenylethanolamine 2.8 mg per lozenge
store at 20 -25°C (68 -77°F)
keep bottle tightly closed and protect from light

Inactive ingredients aspartame, calcium polyacrylate, flavor, methylcellulose, mannitol, sodium alginate, sodium bicarbonate, sodium acrylate fumarate, talc, sorbitol gum

Questions or comments?
Call 1-877-783-8888 Monday-Friday 9AM-5PM EST

*This product is not manufactured or distributed by Ciba-Geigy/Novartis Consumer Healthcare, L.P., distributor of Nicorette® Lozenge.

CVS Health. Compare to the active ingredient in Nicorette® Lozenge* NDC 69842-564-35

Sugar Free Nicotine Lozenge
Nicotine Polacrilex 4 mg
STOP SMOKING AID

4
mg

not for sale to those under 18 years of age

proof of age required

not for sale in vending machines or from any source where proof of age cannot be verified

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP.

If you smoke your first cigarette **MORE THAN 30 MINUTES** after waking up, use Nicotine Polacrilex 2 mg Lozenge.

Actual Size

Mint Mini Lozenge

135 LOZENGES, 4 mg EACH
(5 BOTTLES OF 27)

Includes User's Guide

CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2020 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-36037

CVS Quality
Money Back Guarantee

#491745

0 50426 33078 4

PLD-A215E FC007066

Lot No:
Exp. Date:

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivated to quit.
2. Use Enough – Use at least 9 Nicotine Mini Lozenges per day during the first six weeks.
3. Use Long Enough – Use Nicotine Mini Lozenges for the full 12 weeks.
4. Use With a Support Program as directed in the enclosed User's Guide.

Opening Directions:
Push down on cap and turn to open.

TAMPER EVIDENT FEATURE: DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING.

RETAIN OUTER CARTON FOR FULL PRODUCT USES, DIRECTIONS AND WARNINGS.

CVS HEALTH Sugar Free Nicotine Lozenge 4 mg

NICOTINE

nicotine polacrilex lozenge

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69842-564
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength

NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	4 mg
---	----------	------

Inactive Ingredients

Ingredient Name	Strength
ASPARTAME (UNII: Z0H242BBR1)	
CALCIUM POLYCARBOPHIL (UNII: 8F049NKY49)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MANNITOL (UNII: 3OWL53L36A)	
SODIUM ALGINATE (UNII: C269C4G2ZQ)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM STEARYL FUMARATE (UNII: 7CV7WJK4UI)	
TALC (UNII: 7SEV7J4R1U)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	white	Score	no score
Shape	OVAL	Size	10mm
Flavor	MINT	Imprint Code	131
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69842-564-81	3 in 1 BOX	05/29/2020	
1		27 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:69842-564-27	1 in 1 BOX	05/29/2020	
2		27 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:69842-564-20	1 in 1 BOX	05/29/2020	
3		20 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
4	NDC:69842-564-10	1 in 1 BOX	05/29/2020	
4		10 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
5	NDC:69842-564-35	5 in 1 BOX	05/29/2020	
5		27 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA207868	05/29/2020	

Labeler - CVS Pharmacy (062312574)

Revised: 11/2022

CVS Pharmacy