

**LEADER TRIPLE ANTIBIOTIC PLUS- bacitracin zinc, neomycin sulfate, polymyxin b sulfate, pramoxine hydrochloride ointment**  
**CARDINAL HEALTH**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Leader Triple Antibiotic Ointment Plus 1 oz. NBE Neosporin + Pain Relief**

Active Ingredients (in each gram)	Purpose
Bacitracin Zinc 500 units .....	First aid antibiotic
Neomycin Sulfate 3.5 mg.....	First aid antibiotic
Polymyxin-B Sulfate 10,000 units.....	First aid antibiotic
Pramoxine hydrochloride 10 mg.....	Pain reliever

**Uses**

help prevent infection and temporarily relieves pain due to

- minor cuts
- scrapes
- burns

**Warnings**

For external use only

Allergy alert: do not use if allergic to any of the ingredients

Do not use

- in or near the eyes
- over large areas of the body
- longer than 1 week

Ask a doctor before use in case of deep or puncture wounds, animal bites, or serious burns

When using this product do not use longer than 1 week

Stop use and ask a doctor if

- condition persists or gets worse
- symptoms last for more than 7 days or clear up and comes back within a few days
- a rash or other allergic reaction occurs

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

Adults and children 2 years and older:

- clean affected area
- apply a small amount (equal to the surface area of the tip of a finger) on area 1 to 3 times daily
- may be covered with a sterile bandage

**Other information**

- store at 15 ° to 25 ° C (59 ° to 77 ° F)
- Lot No. and Exp. Date: see box or crimp of tube

Inactive ingredients

white petrolatum, water

DISTRIBUTED BY CARDINAL HEALTH

DUBLIN, OHIO 43017

CIN 2372167

[www.myleader.com](http://www.myleader.com)

1-800-200-6313

Made in Korea



Compare to Neosporin<sup>®</sup> Plus active ingredients

Polymyxin B Sulfate • Bacitracin Zinc • Neomycin Sulfate • Pramoxine HCl  
**Triple Antibiotic Ointment Plus**

Helps prevent infection and provide temporary relief of pain or discomfort in minor cuts, scrapes and burns

NDC 37205-266-10



Polymyxin B Sulfate • Bacitracin Zinc • Neomycin Sulfate • Pramoxine HCl  
**Triple Antibiotic Ointment Plus**  
FIRST AID ANTIBIOTIC / PAIN RELIEF



NET WT 1 OZ (28 g)

DISTRIBUTED BY CARDINAL HEALTH  
DUBLIN, OHIO 43017  
CIN 2372167  
www.myleader.com  
1-800-200-6313



0 96295 10487 5  
Made in Korea

All Leader<sup>®</sup> Brand products are 100% satisfaction guaranteed or return to place of purchase for a full refund.

This product is not manufactured or distributed by Johnson & Johnson Consumer Companies, Inc., owner of the registered trademark Neosporin<sup>®</sup> Plus.

**Drug Facts**

**Active ingredients (in each gram)**  
Bacitracin zinc 500 units  
Neomycin sulfate 3.5 mg  
Polymyxin B sulfate 10,000 units  
Pramoxine hydrochloride 10 mg  
External analgesic

**Purpose**  
First aid antibiotic  
First aid antibiotic  
First aid antibiotic

**Uses** helps prevent infection and temporarily relieves pain due to minor cuts, scrapes and burns.

**Warnings**  
For external use only

**Allergy alert:** do not use if you are allergic to any of the ingredients

**Do not use** ■ in or near the eyes ■ over large areas of the body

**Ask a doctor before use if you have** ■ deep or puncture wounds ■ animal bites ■ serious burns

**When using this product do not use longer than 1 week**  
■ condition lasts or gets worse ■ symptoms last for more than 1 week, or clear up and occur again within a few days ■ rash or other allergic reaction develops

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**  
■ adults and children 2 years of age and older:  
■ clean affected area ■ apply a small amount (equal to the surface area of tip of a finger) on area 1 to 3 times daily  
■ may be covered with a sterile bandage  
■ children under 2 years of age: ask a doctor

**Other information** ■ store at 20° to 25°C (68° to 77°F) ■ Lot No. & Exp. Date: see box or see crimp of the tube

**Inactive ingredients** liquid paraffin, white petrolatum

**Questions or Comments?** 1-800-814-8028

LOT & EXP.

# LEADER TRIPLE ANTIBIOTIC PLUS

bacitracin zinc, neomycin sulfate, polymyxin b sulfate, pramoxine hydrochloride ointment

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:37205-266
<b>Route of Administration</b>	TOPICAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>BACITRACIN ZINC</b> (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RWO52I)	BACITRACIN	500 [USP <sup>U</sup> ] in 1 g
<b>NEOMYCIN SULFATE</b> (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN	3.5 mg in 1 g
<b>POLYMYXIN B SULFATE</b> (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ07J96K)	POLYMYXIN B	10000 [USP <sup>U</sup> ] in 1 g
<b>PRAMOXINE HYDROCHLORIDE</b> (UNII: 88AYB867L5) (PRAMOXINE - UNII:068X84E056)	PRAMOXINE HYDROCHLORIDE	10 mg in 1 g

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>PETROLATUM</b> (UNII: 4T6H12BN9U)	
<b>WATER</b> (UNII: 059QF0KO0R)	

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:37205-266-10	1 in 1 CARTON	03/01/2012	
1		28 g in 1 TUBE; Type 0: Not a Combination Product		

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC monograph final	part333B	03/01/2012	

**Labeler** - CARDINAL HEALTH (097537435)

Revised: 12/2017

CARDINAL HEALTH