

**CIRCLE K IBUPROFEN- ibuprofen tablet, film coated**  
**Lil Drug Store Products, Inc**

-----  
**Circle K™ Ibuprofen**

***Drug Facts***

**Active ingredient (in each tablet)**

Ibuprofen 200 mg (NSAID <sup>1</sup>)

---

<sup>1</sup> nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever/Fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - toothache
  - backache
  - menstrual cramps
  - the common cold
  - muscular aches
  - minor pain of arthritis
- temporarily reduces fever

**Warnings**

**Allergy alert**

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Heart attack and stroke warning**

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic

### **Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

### **When using this product**

- take with food or milk if stomach upset occurs

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of the body
  - slurred speech
  - leg swelling

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over:
  - take 1 tablet every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 tablet, 2 tablets may be used
  - do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

### **Other information**

- store between 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F).

### **Inactive ingredients**

carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

### **Questions or comments?**

call toll-free **1-877-507-6516 (M-F 8 AM-4:30PM CST)**

**Proudly distributed by Circle K Stores Inc.**

### **PRINCIPAL DISPLAY PANEL - 200 mg Tablet Pouch Carton**

Compare to the Active  
Ingredient in Advil<sup>®</sup> Tablets\*\*

CIRCLE K<sup>®</sup>

Ibuprofen

Pain Reliever/Fever Reducer (NSAID), 200 mg

ACTUAL SIZE

6 Coated Tablets

3 POUCHES, 2 TABLETS PER POUCH

ibuprofen



### Drug Facts (continued)

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over:
  - take 1 tablet every 4 to 6 hours while symptoms persist
  - if pain or fever does not respond to 1 tablet, 2 tablets may be used
  - do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

**Other information** • store between 20-25°C (68-77°F). Avoid excessive heat above 40°C (104°F).

**Inactive ingredients** carnauba wax, colloidal silicon dioxide, corn starch, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

### Questions or comments?

call toll-free 1-877-507-6516 (M-F 8 AM-4:30PM CST)

### Product manufactured for:

Lif Drug Store Products, Inc., 9300 Earhart Lane SW, Cedar Rapids, IA 52404  
www.lifdrugstore.com 57393C-US-06-17-A

Proudly distributed by Circle K Stores Inc.

\*This product is not manufactured or distributed by Pfizer Consumer Healthcare. Lif Drug Store Products does not own the Advil® trademark.



Ibuprofen  
6 Tablets

Compare to the Active  
Ingredient in Advil® Tablets\*\*



# Ibuprofen

Pain Reliever/Fever Reducer (NSAID), 200 mg



ACTUAL SIZE

## 6 Coated Tablets

3 POUCHES, 2 TABLETS PER POUCH

Ibuprofen 6 Tablets



**DO NOT USE IF INNER POUCH OR CARTON IS OPEN OR DAMAGED!**

### Drug Facts

#### Active ingredient (in each tablet)

Ibuprofen 200 mg (NSAID)\*.....Pain reliever/Fever reducer  
\*nonsteroidal anti-inflammatory drug

#### Purpose

### Uses

- temporarily relieves minor aches and pains due to:
  - headache
  - toothache
  - backache
  - menstrual cramps
  - the common cold
  - muscular aches
  - minor pain of arthritis
- temporarily reduces fever

### Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

LIFT PANEL FOR CONTINUED DRUG FACTS

### Drug Facts (continued)

#### Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

#### Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic

#### Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product** • take with food or milk if stomach upset occurs

#### Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of the body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

PDP/Package - NDC 66715-5539

TEMPORARILY RELIEVES MINOR ACHES & PAINS

Compare to the active ingredient in  
Advil Tablets\*

CIRCLE K™

200 mg  
ibuprofen  
Ibuprofen 200 mg (NSAID),  
Pain Reliever/Fever Reducer

[tablet image]

actual size

6 coated tablets (2 per pouch)



CIRCLE K IBUPROFEN

ibuprofen tablet, film coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66715-5839
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ )	
STARCH, CORN (UNII: O8232NY3SJ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>ANHYDROUS LACTOSE</b> (UNII: 3SY5LH9PMK)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	brown	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	44;291
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-5839-2	2 in 1 CARTON	06/21/2016	12/10/2022
1		2 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:66715-5839-3	2 in 1 CARTON	10/15/2019	05/12/2024
2		2 in 1 POUCH; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075010	06/21/2016	05/12/2024

## CIRCLE K IBUPROFEN

ibuprofen tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:66715-5539
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>IBUPROFEN</b> (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>ANHYDROUS LACTOSE</b> (UNII: 3SY5LH9PMK)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I3O)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	brown	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	44;291
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-5539-3	2 in 1 CARTON	03/01/2021	
1		2 in 1 POUCH; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075010	03/01/2021	

**Labeler -** Lil Drug Store Products, Inc (093103646)