

TEPMETKO- tepotinib hydrochloride tablet

EMD Serono, Inc.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use TEPMETKO safely and effectively. See full prescribing information for TEPMETKO.

TEPMETKO® (tepotinib) tablets, for oral use
Initial U.S. Approval: 2021

RECENT MAJOR CHANGES

Indication and Usage (1)	02/2024
Dosage and Administration (2.3, 2.4)	02/2024
Warnings and Precautions (5.3)	02/2024

INDICATIONS AND USAGE

TEPMETKO is a kinase inhibitor indicated for the treatment of adult patients with metastatic non-small cell lung cancer (NSCLC) harboring mesenchymal-epithelial transition (*MET*) exon 14 skipping alterations. (1)

DOSAGE AND ADMINISTRATION

- Select patients for treatment with TEPMETKO on the presence of *MET*ex14 skipping. (2.1, 14)
- Recommended dosage: 450 mg orally once daily with food until disease progression or unacceptable toxicity. (2.2)

DOSAGE FORMS AND STRENGTHS

Tablets: 225 mg. (3)

CONTRAINDICATIONS

None. (4)

WARNINGS AND PRECAUTIONS

- Interstitial Lung Disease (ILD)/Pneumonitis: Immediately withhold TEPMETKO in patients with suspected ILD/pneumonitis. Permanently discontinue TEPMETKO in patients diagnosed with ILD/pneumonitis of any severity. (2.4, 5.1)
- Hepatotoxicity: Monitor liver function tests. Withhold, dose reduce, or permanently discontinue TEPMETKO based on severity. (5.2)
- Pancreatic Toxicity: Monitor amylase and lipase. Withhold, dose reduce, or permanently discontinue TEPMETKO based on severity. (5.3)
- Embryo-fetal toxicity: TEPMETKO can cause fetal harm. Advise of potential risk to a fetus and use of effective contraception. (5.4, 8.1, 8.3)

ADVERSE REACTIONS

Most common adverse reactions ($\geq 20\%$) were edema, nausea, fatigue, musculoskeletal pain, diarrhea, dyspnea, decreased appetite, and rash. The most common Grade 3 to 4 laboratory abnormalities ($\geq 2\%$) were decreased lymphocytes, decreased albumin, decreased sodium, increased gamma-glutamyltransferase, increased amylase, increased lipase, increased ALT, increased AST, and decreased hemoglobin. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact EMD Serono at 1-800-283-8088 ext. 5563 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- Certain P-gp substrates: Avoid coadministration of TEPMETKO with P-gp substrates where minimal concentration changes may lead to serious or life-threatening toxicities. (7.1)

USE IN SPECIFIC POPULATIONS

Lactation: Advise not to breastfeed. (8.2)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 2/2024

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Patient Selection for METex14 Skipping Alterations
- 2.2 Recommended Dosage
- 2.3 Administration to Patients Who Have Difficulty Swallowing Solids
- 2.4 Dose Modifications for Adverse Reactions

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Interstitial Lung Disease (ILD)/Pneumonitis
- 5.2 Hepatotoxicity
- 5.3 Pancreatic Toxicity
- 5.4 Embryo-Fetal Toxicity

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience

7 DRUG INTERACTIONS

- 7.1 Effects of TEPMETKO on Other Drugs

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.3 Females and Males of Reproductive Potential
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment
- 8.7 Hepatic Impairment

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

TEPMETKO is indicated for the treatment of adult patients with metastatic non-small cell lung cancer (NSCLC) harboring mesenchymal-epithelial transition (*MET*) exon 14 skipping alterations.

2 DOSAGE AND ADMINISTRATION

2.1 Patient Selection for METex14 Skipping Alterations

Select patients for treatment with TEPMETKO based on the presence of *MET* exon 14 skipping alterations in plasma or tumor specimens. Testing for the presence of *MET* exon 14 skipping alterations in plasma specimens is recommended only in patients for whom a tumor biopsy cannot be obtained. If an alteration is not detected in a plasma specimen, re-evaluate the feasibility of biopsy for tumor tissue testing. An FDA-approved test for detection of *MET* exon 14 skipping alterations in NSCLC for selecting patients for treatment with TEPMETKO is not available.

2.2 Recommended Dosage

The recommended dosage of TEPMETKO is 450 mg orally once daily with food [*see Clinical Pharmacology (12.3)*] until disease progression or unacceptable toxicity.

Instruct patients to take their dose of TEPMETKO at approximately the same time every day and to swallow tablets whole. Do not chew, crush or split tablets. Patients who have difficulty swallowing solids can disperse tablets in water [*see Dosage and Administration (2.3)*].

Advise patients not to make up a missed dose within 8 hours of the next scheduled dose.

If vomiting occurs after taking a dose of TEPMETKO, advise patients to take the next dose at the scheduled time.

2.3 Administration to Patients Who Have Difficulty Swallowing Solids

Place TEPMETKO tablet(s) in a glass containing 30 mL (1 ounce) of non-carbonated water. No other liquids should be used or added. Stir, without crushing, until the tablet(s) is dispersed into small pieces (tablets will not completely dissolve) and drink immediately or within 1 hour. Swallow the tablet dispersion. Do not chew pieces of the tablet. Rinse the glass with an additional 30 mL and drink immediately ensuring no residue remains in the glass and the full dose is administered.

If an administration via a naso-gastric tube (with at least 8 French gauge) is required, disperse the tablet(s) in 30 mL of non-carbonated water as described above. Administer the 30 mL of liquid immediately or within 1 hour as per naso-gastric tube manufacturer's instructions. Immediately rinse twice with 30 mL each time to ensure that no residue remains in the glass or syringe and the full dose is administered.

2.4 Dose Modifications for Adverse Reactions

The recommended dose reduction of TEPMETKO for the management of adverse reactions is 225 mg orally once daily.

Permanently discontinue TEPMETKO in patients who are unable to tolerate 225 mg orally

once daily.

The recommended dosage modifications of TEPMETKO for adverse reactions are provided in Table 1.

Table 1: Recommended TEPMETKO Dosage Modifications for Adverse Reactions

Adverse Reaction	Severity*	Dose Modification
Interstitial Lung Disease (ILD)/Pneumonitis [see <i>Warnings and Precautions (5.1)</i>]	Any Grade	Withhold TEPMETKO if ILD is suspected. Permanently discontinue TEPMETKO if ILD is confirmed.
Increased ALT and/or AST without increased total bilirubin [see <i>Warnings and Precautions (5.2)</i>]	Grade 3	Withhold TEPMETKO until recovery to baseline ALT/AST. If recovered to baseline within 7 days, then resume TEPMETKO at the same dose; otherwise resume TEPMETKO at a reduced dose.
	Grade 4	Permanently discontinue TEPMETKO.
Increased ALT and/or AST with increased total bilirubin in the absence of cholestasis or hemolysis [see <i>Warnings and Precautions (5.2)</i>]	ALT and/or AST greater than 3 times ULN with total bilirubin greater than 2 times ULN	Permanently discontinue TEPMETKO.
Increased total bilirubin without concurrent increased ALT and/or AST [see <i>Warnings and Precautions (5.2)</i>]	Grade 3	Withhold TEPMETKO until recovery to baseline bilirubin. If recovered to baseline within 7 days, then resume TEPMETKO at a reduced dose; otherwise permanently discontinue.
	Grade 4	Permanently discontinue TEPMETKO.
Increased lipase or amylase [see <i>Warnings and Precautions (5.3)</i>]	Grade 3	Withhold TEPMETKO until \leq Grade 2 or baseline. If recovered to baseline or \leq Grade 2 within 14 days, resume TEPMETKO at a reduced dose; otherwise permanently discontinue TEPMETKO.
	Grade 4	Permanently discontinue TEPMETKO.
Pancreatitis [see <i>Warnings and</i>	Grade 3 or 4	Permanently discontinue

<i>warnings and Precautions (5.3)]</i>	Grade 3 or 4	TEPMETKO.
Other adverse reactions [see Adverse Reactions (6.1)]	Grade 2	Maintain dose level. If intolerable, consider withholding TEPMETKO until resolved, then resume TEPMETKO at a reduced dose.
	Grade 3	Withhold TEPMETKO until resolved, then resume TEPMETKO at a reduced dose.
	Grade 4	Permanently discontinue TEPMETKO.

* Severity as defined by National Cancer Institute Common Terminology Criteria for Adverse Events (NCI CTCAE) Version 5.

3 DOSAGE FORMS AND STRENGTHS

Tablets: 225 mg, white-pink, oval, biconvex film-coated tablets with embossment "M" on one side and plain on the other side.

4 CONTRAINDICATIONS

None.

5 WARNINGS AND PRECAUTIONS

5.1 Interstitial Lung Disease (ILD)/Pneumonitis

ILD/pneumonitis, which can be fatal, occurred in patients treated with TEPMETKO [see Adverse Reactions (6.1)]. ILD/pneumonitis occurred in 2% patients treated with TEPMETKO, with one patient experiencing a Grade 3 or higher event; this event resulted in death. Five patients (1%) discontinued TEPMETKO due to ILD/pneumonitis.

Monitor patients for new or worsening pulmonary symptoms indicative of ILD/pneumonitis (e.g., dyspnea, cough, fever). Immediately withhold TEPMETKO in patients with suspected ILD/pneumonitis and permanently discontinue if no other potential causes of ILD/pneumonitis are identified [see Dosage and Administration (2.4)].

5.2 Hepatotoxicity

Hepatotoxicity occurred in patients treated with TEPMETKO [see Adverse Reactions (6.1)]. Increased alanine aminotransferase (ALT)/increased aspartate aminotransferase (AST) occurred in 18% of patients treated with TEPMETKO. Grade 3 or 4 increased ALT/AST occurred in 4.7% of patients. A fatal adverse reaction of hepatic failure occurred in one patient (0.2%). Four patients (0.8%) discontinued TEPMETKO due to increased ALT/AST. The median time-to-onset of Grade 3 or higher increased ALT/AST was 47 days (range 1 to 262).

Monitor liver function tests (including ALT, AST, and total bilirubin) prior to the start of TEPMETKO, every 2 weeks during the first 3 months of treatment, then once a month or as clinically indicated, with more frequent testing in patients who develop increased

transaminases or bilirubin. Based on the severity of the adverse reaction, withhold, dose reduce, or permanently discontinue TEPMETKO [see *Dosage and Administration (2.4)*].

5.3 Pancreatic Toxicity

Elevations in amylase and lipase levels occurred in patients treated with TEPMETKO [see *Adverse Reactions (6.1)*]. Increased amylase and/or lipase occurred in 13% of patients treated with TEPMETKO. Grade 3 and 4 increased amylase and/or lipase occurred in 5% and 1.2% of patients, respectively. Monitor amylase and lipase at baseline and regularly during treatment with TEPMETKO. Based on the severity of the adverse drug reaction, temporarily withhold, dose reduce, or permanently discontinue TEPMETKO [see *Dosage and Administration (2.4)*].

5.4 Embryo-Fetal Toxicity

Based on findings in animal studies and its mechanism of action TEPMETKO can cause fetal harm when administered to a pregnant woman. Oral administration of tepotinib to pregnant rabbits during the period of organogenesis resulted in malformations (teratogenicity) and anomalies at exposures less than the human exposure based on area under the curve (AUC) at the 450 mg daily clinical dose. Advise pregnant women of the potential risk to a fetus. Advise females of reproductive potential or males with female partners of reproductive potential to use effective contraception during treatment with TEPMETKO and for one week after the last dose. [See *Use in Specific Populations (8.1, 8.3)*]

6 ADVERSE REACTIONS

The following adverse reactions are described in greater detail elsewhere in the labeling:

- Interstitial Lung Disease/Pneumonitis [see *Warnings and Precautions (5.1)*]
- Hepatotoxicity [see *Warnings and Precautions (5.2)*]
- Pancreatic Toxicity [see *Warnings and Precautions (5.3)*]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The pooled safety population described in the WARNINGS AND PRECAUTIONS reflect exposure to TEPMETKO in 506 patients with solid tumors enrolled in five open-label, single-arm studies receiving TEPMETKO as single agent at a dose of 450 mg once daily. This included 313 patients with NSCLC positive for *MET*ex14 skipping alterations, who received TEPMETKO in VISION. Among 506 patients who received TEPMETKO, 44% were exposed for 6 months or longer, and 22% were exposed for more than one year.

The data described below reflect exposure to TEPMETKO 450 mg once daily in 313 patients with metastatic non-small cell lung cancer (NSCLC) with *MET*ex14 skipping alterations in VISION [see *Clinical Studies (14)*].

Serious adverse reactions occurred in 51% of patients who received TEPMETKO. Serious adverse reactions in > 2% of patients included pleural effusion (6%), pneumonia (6%), edema (5%), general health deterioration (3.8%), dyspnea (3.5%), musculoskeletal pain

(2.9%), and pulmonary embolism (2.2%). Fatal adverse reactions occurred in 1.9% of patients who received TEPMETKO, including pneumonitis (0.3%), hepatic failure (0.3%), dyspnea from fluid overload (0.3%), pneumonia (0.3%), sepsis (0.3%), and death of unknown cause (0.3%).

Permanent discontinuation due to an adverse reaction occurred in 25% of patients who received TEPMETKO. The most frequent adverse reactions (> 1%) leading to permanent discontinuations of TEPMETKO were edema (8%), pleural effusion (1.6%), and general health deterioration (1.6%).

Dosage interruptions due to an adverse reaction occurred in 53% of patients who received TEPMETKO. Adverse reactions which required dosage interruption in > 2% of patients who received TEPMETKO included edema (28%), increased blood creatinine (6%), pleural effusion (3.5%), nausea (3.2%), increased ALT (2.9%), pneumonia (2.6%), decreased appetite (2.2%), and dyspnea (2.2%).

Dose reductions due to an adverse reaction occurred in 36% of patients who received TEPMETKO. Adverse reactions which required dose reductions in > 2% of patients who received TEPMETKO included edema (22%), increased blood creatinine (2.9%), fatigue (2.2%), and pleural effusion (2.2%).

The most common adverse reactions ($\geq 20\%$) in patients who received TEPMETKO were edema, nausea, fatigue, musculoskeletal pain, diarrhea, dyspnea, decreased appetite, and rash. The most common Grade 3 to 4 laboratory abnormalities ($\geq 2\%$) were decreased lymphocytes, decreased albumin, decreased sodium, increased gamma-glutamyltransferase, increased amylase, increased lipase, increased ALT, increased AST, and decreased hemoglobin.

Table 2 summarizes the adverse reactions in VISION.

Table 2: Adverse Reactions in $\geq 10\%$ of Patients with NSCLC with METex14 Skipping Alterations Who Received TEPMETKO in VISION

Adverse Reactions	TEPMETKO (N=313)	
	All Grades* (%)	Grades 3 to 4* (%)
General disorders and administration-site conditions		
Edema [†]	81	16
Fatigue [‡]	30	1.9
Gastrointestinal disorders		
Nausea	31	1.3
Diarrhea	29	0.6
Abdominal pain [§]	19	0.6
Constipation	19	0.3
Vomiting [¶]	15	1
Musculoskeletal and Connective Tissue Disorders		
Musculoskeletal pain [#]	30	3.2
Respiratory, thoracic, and mediastinal disorders		
Dyspnea ^p	24	2.6
Cough ^β	18	0.3

Pleural effusion	14	4.2
Metabolism and nutrition disorders		
Decreased appetite	21	1.9
Skin and Subcutaneous Tissue Disorders		
Rash ^à	21	1.3
Infections and Infestations		
Pneumonia ^è	12	3.8

* Severity as defined by National Cancer Institute Common Terminology Criteria for Adverse Events (NCI CTCAE) version 4.03.

† Edema includes eye edema, face edema, generalized edema, localized edema, edema, genital edema, peripheral edema, peripheral swelling, periorbital edema, and scrotal edema.

‡ Fatigue includes asthenia and fatigue.

§ Abdominal pain includes abdominal discomfort, abdominal pain, abdominal pain lower, abdominal pain upper, gastrointestinal pain, and hepatic pain.

¶ Vomiting includes retching and vomiting.

Musculoskeletal pain includes arthralgia, arthritis, back pain, bone pain, musculoskeletal chest pain, musculoskeletal pain, myalgia, non-cardiac chest pain, pain in extremity, and spinal pain.

Ⓟ Dyspnea includes dyspnea, dyspnea at rest, and dyspnea exertional.

Ⓠ Cough includes cough, and productive cough.

à Rash includes rash, palmar-plantar erythrodysesthesia syndrome, rash maculopapular, eczema, exfoliative rash, rash erythematous, rash pustular, skin exfoliation, dermatitis acneiform, drug eruption, dermatitis, rash pruritic, dermatitis bullous, toxic skin eruption.

è Pneumonia includes pneumonia, pneumonia aspiration, and pneumonia bacterial.

Clinically relevant adverse reactions in < 10% of patients who received TEPMETKO included ILD/pneumonitis, fever, dizziness, pruritus, and headache.

Table 3 summarizes the laboratory abnormalities observed in VISION.

Table 3: Select Laboratory Abnormalities (≥ 20%) That Worsened from Baseline in Patients Who Received TEPMETKO in VISION

Laboratory Abnormalities	TEPMETKO*	
	Grades 1 to 4 [†] (%)	Grades 3 to 4 [†] (%)
Chemistry		
Decreased albumin	81	9
Increased creatinine	60	1
Increased alkaline phosphatase aminotransferase	52	1.6
Increased alanine aminotransferase	50	4.9
Increased aspartate aminotransferase	40	3.6
Decreased sodium	36	9
Increased gamma-glutamyltransferase	29	6
Increased potassium	26	1.9
Increased amylase	25	5
Increased lipase	21	5

Hematology		
Decreased lymphocytes	57	15
Decreased hemoglobin	31	3.6
Decreased leukocytes	25	1.9
Decreased platelets	24	0.6

* The denominator used to calculate the rate varied from 268 to 309 based on the number of patients with a baseline value and at least one post-treatment value.

† Severity as defined by National Cancer Institute Common Terminology Criteria for Adverse Events (NCI CTCAE) version 4.03.

Increased Creatinine

A median increase in serum creatinine of 30% was observed 21 days after initiation of treatment with TEPMETKO. The serum creatinine increases persisted throughout treatment and were reversible upon treatment completion.

7 DRUG INTERACTIONS

7.1 Effects of TEPMETKO on Other Drugs

Certain P-gp Substrates

Tepotinib is a P-gp inhibitor. Concomitant use of TEPMETKO increases the concentration of P-gp substrates [see *Clinical Pharmacology (12.3)*], which may increase the incidence and severity of adverse reactions of these substrates. Avoid concomitant use of TEPMETKO with certain P-gp substrates where minimal concentration changes may lead to serious or life-threatening toxicities. If concomitant use is unavoidable, reduce the P-gp substrate dosage if recommended in its approved product labeling.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Based on findings in animal studies and the mechanism of action [see *Clinical Pharmacology (12.1)*], TEPMETKO can cause fetal harm when administered to a pregnant woman. There are no available data on the use of TEPMETKO in pregnant women. Oral administration of tepotinib to pregnant rabbits during the period of organogenesis resulted in malformations (teratogenicity) and anomalies at maternal exposures less than the human exposure based on area under the curve (AUC) at the 450 mg daily clinical dose (see *Data*). Advise pregnant women of the potential risk to a fetus.

In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively.

Data

Animal Data

In embryo-fetal development studies, pregnant rabbits received oral doses of 0.5, 5, 25, 50, 150, or 450 mg/kg tepotinib hydrochloride hydrate daily during organogenesis. Severe maternal toxicity occurred at the 450 mg/kg dose (approximately 0.75 times the human exposure at the 450 mg clinical dose). At 150 mg/kg (approximately 0.5 times the human exposure by AUC at the 450 mg clinical dose), two animals aborted and one animal died prematurely; mean fetal body weight was also decreased. A dose-dependent increase of skeletal malformations, including malrotations of fore and/or hind paws with concomitant misshapen scapula and/or malpositioned clavicle and/or calcaneus and/or talus, occurred at doses \geq 5 mg/kg (approximately 0.003 times the human exposure by AUC at the 450 mg clinical dose); there was also an incidence of spina bifida at the 5 mg/kg dose level.

8.2 Lactation

Risk Summary

There are no data regarding the secretion of tepotinib or its metabolites in human milk or its effects on the breastfed infant or milk production. Advise women not to breastfeed during treatment with TEPMETKO and for one week after the last dose.

8.3 Females and Males of Reproductive Potential

Based on animal data, TEPMETKO can cause malformations at doses less than the human exposure based on AUC at the 450 mg clinical dose [*see Use in Specific Populations (8.1)*].

Pregnancy Testing

Verify pregnancy status in females of reproductive potential prior to initiating TEPMETKO [*see Use in Specific Populations (8.1)*].

Contraception

Females

Advise females of reproductive potential to use effective contraception during TEPMETKO treatment and for one week after the last dose.

Males

Advise male patients with female partners of reproductive potential to use effective contraception during TEPMETKO treatment and for one week after the last dose.

8.4 Pediatric Use

The safety and efficacy of TEPMETKO in pediatric patients have not been established.

8.5 Geriatric Use

Of 313 patients with NSCLC positive for *MET*ex14 skipping alterations in VISION who received 450 mg TEPMETKO once daily, 79% were 65 years or older, and 41% were 75 years or older. No clinically important differences in safety or efficacy were observed between patients aged 65 years or older and younger patients.

8.6 Renal Impairment


No dosage modification is recommended in patients with mild or moderate renal impairment (creatinine clearance [CL_{cr}] 30 to 89 mL/min, estimated by Cockcroft-Gault). The recommended dosage has not been established for patients with severe renal impairment (CL_{cr} < 30 mL/min) [see *Clinical Pharmacology* (12.3)].

8.7 Hepatic Impairment

No dosage modification is recommended in patients with mild (Child Pugh Class A) or moderate (Child Pugh Class B) hepatic impairment. The pharmacokinetics and safety of tepotinib in patients with severe hepatic impairment (Child Pugh Class C) have not been studied [see *Clinical Pharmacology* (12.3)].

11 DESCRIPTION

Tepotinib is a kinase inhibitor. TEPMETKO (tepotinib) tablets for oral use are formulated with tepotinib hydrochloride hydrate. The chemical name for tepotinib hydrochloride hydrate is 3-{1-[(3-{5-[(1-methylpiperidin-4-yl)methoxy]pyrimidin-2-yl}phenyl)methyl]-6-oxo-1,6-dihydropyridazin-3-yl}benzotrile hydrochloride hydrate. The molecular formula is C₂₉H₂₈N₆O₂•HCl•H₂O and the molecular weight is 547.05 g/mol for tepotinib hydrochloride hydrate and 492.58 g/mol for tepotinib (free base). The chemical structure is shown below:



Tepotinib hydrochloride hydrate is a white to off-white powder with a pK_a of 9.5.

TEPMETKO is supplied as film-coated tablets containing 225 mg of tepotinib (equivalent to 250 mg tepotinib hydrochloride hydrate). Inactive ingredients in the tablet core are mannitol, microcrystalline cellulose, crospovidone, magnesium stearate, and colloidal silicon dioxide. The tablet coating consists of hypromellose, titanium dioxide, lactose monohydrate, polyethylene glycol, triacetin, and red iron oxides.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Tepotinib is a kinase inhibitor that targets MET, including variants with exon 14 skipping alterations. Tepotinib inhibits hepatocyte growth factor (HGF)-dependent and -independent MET phosphorylation and MET-dependent downstream signaling pathways. Tepotinib also inhibited melatonin 2 and imidazoline 1 receptors at clinically achievable concentrations.

In vitro, tepotinib inhibited tumor cell proliferation, anchorage-independent growth, and

migration of MET-dependent tumor cells. In mice implanted with tumor cell lines with oncogenic activation of MET, including *MET*_{ex14} skipping alterations, tepotinib inhibited tumor growth, led to sustained inhibition of MET phosphorylation, and, in one model, decreased the formation of metastases.

12.2 Pharmacodynamics

Exposure-Response

Tepotinib exposure-response relationships and the time course of pharmacodynamic response have not been fully characterized.

Cardiac Electrophysiology

At the recommended dosage, no large mean increases in QTc (i.e. > 20 ms) were detected in patients with various solid tumors. A concentration-dependent increase in QTc interval was observed. The QTc effect of tepotinib at high clinical exposures has not been evaluated.

12.3 Pharmacokinetics

The pharmacokinetics of tepotinib were evaluated in patients with cancer administered 450 mg once daily unless otherwise specified. Tepotinib exposure (AUC_{0-12h} and C_{max}) increases dose-proportionally over the dose range of 27 mg (0.06 times the recommended daily dosage) to 450 mg. At the recommended dosage, the geometric mean (coefficient of variation [CV] %) steady state C_{max} was 1,291 ng/mL (48.1%) and the AUC_{0-24h} was 27,438 ng•h/mL (51.7%). The oral clearance of tepotinib did not change with respect to time. The median accumulation was 2.5-fold for C_{max} and 3.3-fold for AUC_{0-24h} after multiple daily doses of tepotinib.

Absorption

The median T_{max} of tepotinib is 8 hours (range from 6 to 12 hours). The geometric mean (CV%) absolute bioavailability of TEPMETKO in the fed state was 71.6% (10.8%) in healthy subjects.

Effect of Food

The mean AUC_{0-INF} of tepotinib increased by 1.6-fold and C_{max} increased by 2-fold, following administration of a high-fat, high-calorie meal (approximately 800 to 1,000 calories, 150 calories from protein, 250 calories from carbohydrate, and 500 to 600 calories from fat). The median T_{max} shifted from 12 hours to 8 hours.

Distribution

The geometric mean (CV%) apparent volume of distribution (V_Z/F) of tepotinib is 1,038 L (24.3%). Protein binding of tepotinib is 98% and is independent of drug concentration at clinically relevant exposures.

Elimination

The apparent clearance (CL/F) of tepotinib is 23.8 L/h (87.5%) and the half-life is 32 hours following oral administration of TEPMETKO in patients with cancer.

Metabolism

Tepotinib is primarily metabolized by CYP3A4 and CYP2C8. One major circulating plasma

metabolite (M506) has been identified.

Excretion

Following a single oral administration of a radiolabeled dose of 450 mg tepotinib, approximately 85% of the dose was recovered in feces (45% unchanged) and 13.6% in urine (7% unchanged). The major circulating metabolite M506 accounted for about 40.4% of the total radioactivity in plasma.

Specific Populations

No clinically significant effects on tepotinib pharmacokinetics were observed based on age (18 to 89 years), race/ethnicity (White, Black, Asian, Japanese, and Hispanic), sex, body weight (35.5 to 136 kg), mild to moderate renal impairment (CL_{cr} 30 to 89 mL/min), or mild to moderate hepatic impairment (Child-Pugh A and B). The effect of severe renal impairment (CL_{cr} < 30 mL/min) and severe hepatic impairment (Child-Pugh C) on the pharmacokinetics of tepotinib has not been studied.

Drug Interaction Studies

Clinical Studies and Model-Informed Approaches

No clinically significant differences in the pharmacokinetics of tepotinib were observed when coadministered with the following drugs: itraconazole (strong CYP3A and P-gp inhibitor), carbamazepine (strong CYP3A inducer) or omeprazole (proton pump inhibitor/acid reducing agent) under fed conditions.

No clinically significant differences in the pharmacokinetics of the following drugs were observed or predicted when coadministered with tepotinib: midazolam (sensitive CYP3A substrate) or CYP2C9 substrates.

P-gp Substrates: Coadministration of TEPMETKO with dabigatran etexilate (P-gp substrate) increased dabigatran C_{max} by 40% and AUC_{0-∞} by 50%.

MATE2 and OCT2 Substrates: No clinically relevant differences in glucose levels were observed when metformin (MATE2 and OCT2 substrate) was coadministered with tepotinib.

In Vitro Studies

Cytochrome P450 Enzymes: Tepotinib is a substrate of CYP3A4 and CYP2C8. Tepotinib and M506 do not inhibit CYP1A2, CYP2A6, CYP2B6, CYP2C8, CYP2C19, CYP2D6 or CYP2E1, and do not induce CYP1A2 or 2B6 at clinically relevant concentrations.

UDP-Glucuronosyltransferase (UGT): Tepotinib and M506 do not inhibit UGT 1A1, 1A9, 2B17, 1A3/4/6 and 2B7/15 at clinically relevant concentrations.

Transporter Systems: Tepotinib is a P-gp substrate. Tepotinib may inhibit intestinal BCRP at clinically relevant concentrations. Tepotinib does not inhibit bile salt export pump (BSEP), organic anion transporter polypeptide (OATP) 1B1, B3, or organic anion transporter (OAT) 1 and 3.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenicity studies have not been performed with tepotinib. Tepotinib and its major circulating metabolite were not mutagenic in vitro in the bacterial reverse mutation (Ames) assay, or a mouse lymphoma assay. In vivo, tepotinib was not genotoxic in a rat micronucleus test.

Fertility studies of tepotinib have not been performed. There were no morphological changes in male or female reproductive organs in repeat-dose toxicity studies in dogs.

14 CLINICAL STUDIES

The efficacy of TEPMETKO was evaluated in a single-arm, open-label, multicenter, non-randomized, multicohort study (VISION, NCT02864992). Eligible patients were required to have advanced or metastatic NSCLC harboring *MET*ex14 skipping alterations, epidermal growth factor receptor (EGFR) wild-type and anaplastic lymphoma kinase (ALK) negative status, at least one measurable lesion as defined by Response Evaluation Criteria in Solid Tumors (RECIST) version 1.1, and Eastern Cooperative Oncology Group (ECOG) Performance Status of 0 to 1. Patients with symptomatic CNS metastases, clinically significant uncontrolled cardiac disease, or who received treatment with any MET or hepatocyte growth factor (HGF) inhibitor were not eligible for the study.

Identification of *MET*ex14 skipping alterations was prospectively determined using central laboratories employing either a PCR-based or next-generation sequencing-based clinical trial assay using tissue (66%) and/or plasma (57%) samples.

Patients received TEPMETKO 450 mg once daily until disease progression or unacceptable toxicity. The major efficacy outcome measure was confirmed overall response rate (ORR) according to Response Evaluation Criteria in Solid Tumors (RECIST v1.1) as evaluated by a Blinded Independent Review Committee (BIRC). An additional efficacy outcome measure was duration of response (DOR) by BIRC.

The efficacy population included 164 treatment naïve patients and 149 previously treated patients. The median age was 72 years (range 41 to 94 years); 51% female; 62% White, 34% Asian; 26% had Eastern Cooperative Oncology Group (ECOG) Performance Status (PS) 0 and 74% had ECOG PS 1; 49% never smoked; 81% had adenocarcinoma; 94% had metastatic disease; and 13% had CNS metastases. Amongst previously treated patients, 84% received prior platinum-based chemotherapy.

Efficacy results are presented in Table 4.

Table 4: Efficacy Results in the VISION study

Efficacy parameter	Treatment-Naïve (N=164)	Previously Treated (N=149)
Overall response rate (95% CI) ^{*,†}	57% (49, 65)	45% (37, 53)
Duration of Response		
Range in months [‡]	1.3+, 56.6+	1.4+, 67.6+
Patients with DOR ≥ 6 months [‡]	66%	66%
Patients with DOR ≥ 12 months [‡]	40%	36%

CI: Confidence interval, + denotes ongoing response.

* Blinded Independent Review Committee (BIRC) review.

† Confirmed responses.

‡ Based on observed duration of response.

16 HOW SUPPLIED/STORAGE AND HANDLING

TEPMETKO (tepotinib) tablets: 225 mg tepotinib, white-pink, oval, biconvex film-coated tablet with embossment "M" on one side and plain on the other side.

NDC number	Size
44087-5000-3	Box of 30 tablets: 3 blister cards each containing 10 tablets
44087-5000-6	Box of 60 tablets: 6 blister cards each containing 10 tablets

The blister cards consist of a child-resistant blister foil.

Store TEPMETKO at 20°C-25°C (68°F-77°F); excursions permitted to 15°C-30°C (59°F-86°F) [see USP-NF Controlled Room Temperature]. Store in original package.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information).

Interstitial Lung Disease (ILD)/Pneumonitis

Inform patients of the risk of severe or fatal ILD/pneumonitis. Advise patients to contact their healthcare provider immediately to report new or worsening respiratory symptoms [see *Warnings and Precautions (5.1)*].

Hepatotoxicity

Inform patients that they will need to undergo lab tests to monitor liver function. Advise patients to immediately contact their healthcare provider for signs and symptoms of liver dysfunction [see *Warnings and Precautions (5.2)*].

Pancreatic Toxicity

Inform patients that they will need to undergo lab tests to monitor pancreatic function. Advise patients to immediately contact their healthcare provider for signs and symptoms of pancreatitis [see *Warnings and Precautions (5.3)*].

Embryo-Fetal Toxicity

Advise males and females of reproductive potential that TEPMETKO can cause fetal harm.

Advise females of reproductive potential to use effective contraception during and for one week after the last dose of TEPMETKO [see *Warnings and Precautions (5.4)* and *Use in Specific Populations (8.3)*].

Advise male patients with female partners of reproductive potential to use effective contraception during treatment with TEPMETKO and for one week after the last dose of TEPMETKO [see *Warnings and Precautions (5.4)* and *Use in Specific Populations (8.3)*].

Lactation

Advise women not to breastfeed during treatment with TEPMETKO and for one week after the last dose [see *Use in Specific Populations (8.2)*].

Drug Interactions

Advise patients to inform their healthcare provider of all concomitant medications, including prescription medicines, over-the-counter drugs and herbal products [see *Drug Interactions (7)*].

Dosing and Administration

Instruct patients to take 450 mg TEPMETKO once daily with food [see *Dosage and Administration (2.2)*].

Missed Dose

Advise patients that a missed dose of TEPMETKO can be taken as soon as remembered on the same day, unless the next dose is due within 8 hours. If vomiting occurs after taking a dose of TEPMETKO, advise patients to take the next dose at the scheduled time [see *Dosage and Administration (2.2)*].

Manufactured for:

EMD Serono, Inc.
Rockland, MA 02370
U.S.A.

TEPMETKO is a trademark of Merck KGaA,
Darmstadt, Germany

PATIENT INFORMATION

TEPMETKO® (tep-MET-co)
(tepotinib)
tablets, for oral use

What is the most important information I should know about TEPMETKO? TEPMETKO may cause serious side effects, including:

- **Lung problems.** TEPMETKO may cause severe or life-threatening swelling (inflammation) of the lungs during treatment that can lead to death. Tell your healthcare provider right away if you develop any new or worsening symptoms of lung problems, including:
 - trouble breathing
 - shortness of breath
 - cough
 - fever

See "What are possible side effects of TEPMETKO?" for more information about side effects.

What is TEPMETKO?

TEPMETKO is a prescription medicine used to treat adults with non-small cell lung cancer (NSCLC) that:

- has spread to other parts of the body (metastatic), and
- whose tumors have an abnormal mesenchymal epithelial transition (MET) gene. Your healthcare provider will perform a test to make sure that TEPMETKO is right for you.

It is not known if TEPMETKO is safe and effective in children.

Before you take TEPMETKO, tell your healthcare provider about all of your medical conditions, including if you:

- have or have had lung or breathing problems other than your lung cancer
- have or have had liver problems
- have or have had pancreatic problems
- are pregnant or plan to become pregnant. TEPMETKO can harm your unborn baby.

Females who are able to become pregnant:

- Your healthcare provider may do a pregnancy test before you start treatment with TEPMETKO.
- You should use effective birth control (contraception) during treatment and for 1 week after the last dose of TEPMETKO. Talk to your healthcare provider about birth control methods that may be right for you.

Males with female partners who are able to become pregnant should use effective birth control (contraception) during treatment with TEPMETKO and for 1 week after the last dose of TEPMETKO.

- are breastfeeding or plan to breastfeed. It is not known if TEPMETKO passes into your breast milk. Do not breastfeed during treatment and for 1 week after the last dose of TEPMETKO.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

How should I take TEPMETKO?

- Take TEPMETKO exactly as your healthcare provider tells you to. Do not change your dose or stop taking TEPMETKO unless your healthcare provider tells you to.
- Take TEPMETKO 1 time a day with food.
- Take your dose of TEPMETKO at about the same time each day.
- Swallow TEPMETKO tablets whole. **Do not chew, crush or split the tablets.**

If you cannot swallow TEPMETKO tablets whole:

- Place your prescribed dose of TEPMETKO tablets in a glass that contains 30 mL (1 ounce) of non-carbonated water. **Do not** use or add any other liquids.
- Stir the TEPMETKO tablets and water until the TEPMETKO tablets are in small pieces (the tablets will not completely dissolve). **Do not** crush TEPMETKO tablets.
- Drink the TEPMETKO and water mixture right away or within 1 hour. Make sure to swallow the mixture. **Do not** chew pieces of the tablet.
- Add another 30 mL of non-carbonated water to the glass and drink it right away to get your full dose of TEPMETKO.

If you have a nasogastric (NG) tube that is 8 French gauge or larger:

- Follow the same instructions for mixing TEPMETKO tablets in a glass that contains 30 mL of non-carbonated water.
- **Do not** use or add any other liquids.
- Stir the TEPMETKO tablets and water until the TEPMETKO tablets are in small pieces (the tablets will not completely dissolve). **Do not** crush TEPMETKO tablets.
- Give the TEPMETKO tablet and water mixture using the NG tube manufacturer instructions right away or within 1 hour.
- Pieces of the tablet may remain in the glass. Right away, add 30 mL of non-carbonated water into the glass. Give the water and any remaining TEPMETKO

through the NG tube right away to make sure that all of the medicine is given. Repeat this step. This will help to ensure that no medicine is left in the glass or syringe and the full prescribed dose of TEPMETKO is given.

- If you miss a dose of TEPMETKO, take it as soon as you remember. If your next dose is due within 8 hours, skip the missed dose and take your next dose at your regular scheduled time.
- If you vomit after taking a dose of TEPMETKO, take your next dose at your regular scheduled time.

What are the possible side effects of TEPMETKO?

TEPMETKO may cause serious side effects, including:

- See "**What is the most important information I should know about TEPMETKO?**"
- **Liver problems.** TEPMETKO may cause abnormal liver blood test results. Your healthcare provider will do blood tests to check your liver function before you start treatment and during treatment with TEPMETKO. Tell your healthcare provider right away if you develop any signs and symptoms of liver problems, including:
 - your skin or the white part of your eyes turns yellow
 - dark or "tea colored" urine
 - light-colored stools (bowel movements)
 - confusion
 - tiredness
 - loss of appetite for several days or longer
 - nausea and vomiting
 - pain, aching, or tenderness on the right side of your stomach-area (abdomen)
 - weakness
 - swelling in your stomach-area
- **Pancreas problems.** TEPMETKO may cause increases in your blood amylase and lipase levels that may indicate a problem with your pancreas. Your healthcare provider will do blood tests to check your pancreatic function before you start treatment and during treatment with TEPMETKO. Tell your healthcare provider right away if you develop any signs and symptoms of pancreas problems, including:
 - upper stomach (abdominal) pain that may spread to your back and get worse with eating
 - weight loss
 - nausea
 - vomiting

The most common side effects of TEPMETKO include:

- swelling in your face or other parts of your body
- nausea
- tiredness
- muscle and joint pain
- diarrhea
- shortness of breath
- rash
- changes in certain blood tests

Your healthcare provider may change your dose, temporarily stop, or permanently stop treatment with TEPMETKO if you develop serious side effects during treatment.

These are not all of the possible side effects of TEPMETKO.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store TEPMETKO?

- Store TEPMETKO at room temperature between 68°F to 77°F (20°C to 25°C).
- TEPMETKO tablets come in blister cards with a child-resistant blister foil.
- Store TEPMETKO in original package.

Keep TEPMETKO and all medicines out of the reach of children.

General information about the safe and effective use of TEPMETKO.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use TEPMETKO for a condition for which it was not prescribed. Do not give TEPMETKO to other people, even if they have the same symptoms you have. It may harm them. You can ask your pharmacist or healthcare provider for information about TEPMETKO that is written for health professionals.

What are the ingredients in TEPMETKO?

Active ingredient: tepotinib

Inactive ingredients: mannitol, microcrystalline cellulose, crospovidone, magnesium stearate, and colloidal silicon dioxide.

Tablet coating: hypromellose, titanium dioxide, lactose monohydrate, polyethylene glycol, triacetin, and red iron oxides.

Manufactured for: EMD Serono, Inc., Rockland, MA 02370, U.S.A.

TEPMETKO is a trademark of Merck KGaA, Darmstadt, Germany.

Product of Germany.

For more information, call toll-free 1-844-662-3631 or go to www.TEPMETKO.com.

This Patient Information has been approved
by the U.S. Food and Drug Administration.

Revised 02/2024

PRINCIPAL DISPLAY PANEL - 30 Tablet Blister Pack Carton

NDC 44087-5000-3

TEPMETKO®

(tepotinib) tablets

225 mg per tablet

Rx Only

Each tablet contains 225 mg of tepotinib
(equivalent to 250 mg tepotinib hydrochloride hydrate)

Each carton contains 3 child resistant blister
cards of 10 tablets each

30 tablets

EMD

SERONO

TEPMETKO®
(tepotinib) tablets
225 mg per tablet

NDC 44087-5000-3

Rx Only

Each tablet contains 225 mg of tepotinib
(equivalent to 250 mg tepotinib hydrochloride hydrate)

Each carton contains 3 child resistant blister
cards of 10 tablets each

30 tablets

**EMD
SERONO**

TEPMETKO

tepotinib hydrochloride tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:44087-5000
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Tepotinib Hydrochloride (UNII: VY5YX2TQ1F) (Tepotinib - UNII:1IJV77E107)	Tepotinib	225 mg

Inactive Ingredients

Ingredient Name	Strength
Mannitol (UNII: 3OWL53L36A)	
Microcrystalline Cellulose (UNII: OP1R32D61U)	
Crospovidone, Unspecified (UNII: 2S7830E561)	
Magnesium stearate (UNII: 70097M6I30)	
Silicon Dioxide (UNII: ETJ7Z6XBU4)	
Hypromellose, Unspecified (UNII: 3NXW29V3WO)	
Lactose monohydrate (UNII: EWQ57Q8I5X)	
Polyethylene Glycol, Unspecified (UNII: 3WJQ0SDW1A)	
Triacetin (UNII: XHX3C3X673)	
Ferric oxide red (UNII: 1K09F3G675)	
Titanium dioxide (UNII: 15FIX9V2JP)	

Product Characteristics

Color	PINK (White-pink)	Score	no score
Shape	OVAL (oval, biconvex)	Size	7mm
Flavor		Imprint Code	M
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:44087-5000-3	3 in 1 CARTON	02/03/2021	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:44087-5000-6	6 in 1 CARTON	02/03/2021	
2		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA214096	02/03/2021	

Labeler - EMD Serono, Inc. (088514898)

Revised: 2/2024

EMD Serono, Inc.