

DAYTIME NITETIME COLD AND FLU- acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl
Meijer Distribution Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Meijer Distribution, Inc. DayTime NiteTime Cold & Flu Drug Facts

Nighttime Cold & Flu

Active ingredients (in each 30 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- a sodium-restricted diet

Ask doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 30 mL contains:** sodium 39 mg
- store at 20-25°C (68-77°F)

Inactive ingredients

alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments?

1-800-719-9260

Daytime Cold & Flu

Active ingredients (in each 15 mL)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Uses

- temporarily relieves common cold/flu symptoms:
- nasal congestion
- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

When using this product

do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 15 mL contains:** sodium 7 mg
- store at 20-25°C (68-77°F)

Inactive ingredients

butylated hydroxyanisole, edetate disodium, FD&C yellow no. 6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

Questions or comments?

1-800-719-9260

Principal Display Panel

DAYTIME & NITETIME COMBO PACK

meijer®

Compare to Vicks® NyQuil® and DayQuil® active ingredients

Compare to Vicks® DayQuil® active ingredients

meijer®

MAXIMUM STRENGTH

NON-DROWSY

daytime cold & flu

Acetaminophen

Pain Reliever | Fever Reducer

Dextromethorphan HBr

Cough Suppressant

Phenylephrine HCl

Nasal Decongestant

Compare to Vicks® NyQuil® active ingredients

meijer®

MAXIMUM STRENGTH

nitetime cold & flu

Acetaminophen

Pain Reliever | Fever Reducer

Dextromethorphan HBr

Cough Suppressant

Doxylamine Succinate

Antihistamine

Powerful Non-Drowsy Relief

Cherry Flavor

ALCOHOL 10%

Multi-Symptom Relief

2-12 FL OZ (355 mL) BOTTLES

TOTAL 24 FL OZ (1.5 pt) (710 mL)

MAXIMUM STRENGTH
NON-DROWSY
**daytime
cold & flu**

Powerful Non-Drowsy Relief

MAXIMUM STRENGTH
**nitetime
cold & flu**

Powerful Nighttime Relief

Cherry Flavor

**DAYTIME & NITETIME
COMBO PACK**

NDC 41250-733-02

meijer

Compare to Vicks®
NyQuil® and DayQuil®
active ingredients*

Compare to
Vicks® NyQuil®
active ingredients*

MAXIMUM
STRENGTH
**nitetime
cold & flu**

Powerful
Nighttime Relief

Cherry Flavor

ALCOHOL 10%



**DO NOT USE IF PRINTED NECKBAND
IS BROKEN OR MISSING**

meijer

MAXIMUM
STRENGTH
NON-DROWSY
**daytime
cold & flu**

Acetaminophen
Pain Reliever | Fever Reducer
Dextromethorphan HBr
Cough Suppressant
Phenylephrine HCl
Nasal Decongestant

Powerful
Non-Drowsy Relief

meijer

MAXIMUM
STRENGTH
**nitetime
cold & flu**

Acetaminophen
Pain Reliever | Fever Reducer
Dextromethorphan HBr
Cough Suppressant
Doxylamine Succinate
Antihistamine

Cherry Flavor
ALCOHOL 10%

Multi-Symptom Relief

2-12 FL OZ (355 mL) BOTTLES
TOTAL 24 FL OZ (1.5 pt) (710 mL)



DIST. BY MEIJER DISTRIBUTION, INC.
GRAND RAPIDS, MI 49544
www.meijer.com



WWW.MEIJER.COM/
SATISFACTION

PD 316913



7 13733 64847 9

73302 6E C6

Nighttime Cold & Flu

Active ingredients (in each 30 mL)	Purpose
Acetaminophen 650 mg	Pain relief/fever reducer
Dextromethorphan HBr 30 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine

Uses: temporarily relieves common cold/flu symptoms:
 ■ cough due to minor throat and bronchial irritation
 ■ sore throat ■ headache ■ minor aches and pains
 ■ fever ■ runny nose and sneezing

Warnings: Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
 ■ more than 4,000 mg of acetaminophen in 24 hours
 ■ with other drugs containing acetaminophen
 ■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin redness ■ blisters ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use:
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking the product.
 ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have:
 ■ liver disease ■ glaucoma
 ■ cough that occurs with too much phlegm (mucus)
 ■ a breathing problem such as emphysema or chronic bronchitis
 ■ trouble urinating due to an enlarged prostate gland
 ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema
 ■ a sodium-restricted diet

Ask a doctor or pharmacist before use if you are:
 ■ taking sedatives or tranquilizers ■ taking the blood-thinning drug warfarin

When using this product:
 ■ excitability may occur, especially in children
 ■ marked drowsiness may occur ■ avoid alcoholic drinks
 ■ be careful when driving a motor vehicle or operating machinery
 ■ if alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if:
 ■ pain or cough gets worse or lasts more than 7 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present
 ■ new symptoms occur
 ■ cough comes back or occurs with rash or headache that lasts.
 These could be signs of a serious condition.

Directions:
 ■ take only as directed - see Overdose warning
 ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information:
 ■ each 30 mL contains sodium 39 mg
 ■ store at 20-25°C (68-77°F)

Inactive ingredients: alcohol, a hydroxy citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments? 1-800-719-9290

Daytime Cold & Flu

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg	Pain relief/fever reducer
Dextromethorphan HBr 30 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses: temporarily relieves common cold/flu symptoms:
 ■ nasal congestion ■ cough due to minor throat and bronchial irritation
 ■ sore throat ■ headache ■ minor aches and pains ■ fever

Warnings: Liver warning: This product contains acetaminophen. Severe liver damage may occur if:
 ■ adult takes more than 4,000 mg of acetaminophen in 24 hours
 ■ child takes more than 5 doses in 24 hours
 ■ takes with other drugs containing acetaminophen
 ■ adult has 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 ■ skin redness ■ blisters ■ rash
 If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use:
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking the product.
 ■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have:
 ■ liver disease ■ heart disease ■ high blood pressure
 ■ thyroid disease ■ diabetes
 ■ trouble urinating due to an enlarged prostate gland
 ■ cough that occurs with too much phlegm (mucus)
 ■ persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood-thinning drug warfarin

When using this product do not use more than directed:
Stop use and ask a doctor if:
 ■ you get nervous, dizzy or sleepless
 ■ pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present
 ■ new symptoms occur
 ■ cough comes back or occurs with rash or headache that lasts.
 These could be signs of a serious condition.

Directions:
 ■ take only as directed - see Overdose warning
 ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
child ren 6 to under 12 yrs	1.5 mL every 4 hrs
child ren 4 to under 6 yrs	ask a doctor
child ren under 4 yrs	do not use

Other information:
 ■ each 15 mL contains sodium 7 mg ■ store at 20-25°C (68-77°F)

Inactive ingredients: butylated hydroxyanisole, edetate disodium, FD & C yellow no. 6, flavor, glycerin, methyl, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

Questions or comments? 1-800-719-9290

meijer Compare to Vicks® DayQuil® active ingredients*

MAXIMUM STRENGTH NON-DROWSY

daytime cold & flu

Powerful Non-Drowsy Relief

Alcohol Free Antihistamine Free

PARENTS: Learn about teen medicine abuse
www.StopMedicineAbuse.org



*These products are not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademarks Vicks® DayQuil® and DayQuil®.



DAYTIME NITETIME COLD AND FLU

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-733
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-733-02	1 in 1 CARTON; Type 0: Not a Combination Product	10/24/2012	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE	355 mL
Part 2	1 BOTTLE	355 mL

Part 1 of 2

NITETIME COLD AND FLU

acetaminophen, dextromethorphan hbr, doxylamine succinate solution

Product Information

Item Code (Source)	NDC:41250-459
---------------------------	---------------

Route of Administration	ORAL
--------------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

ALCOHOL (UNII: 3K9958V90M)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	

Product Characteristics

Color	RED (Clear/Dark Red)	Score	
Shape		Size	
Flavor	CHERRY (Menthol Aroma)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-459-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	09/09/2011	

Part 2 of 2

DAYTIME COLD AND FLU NON DROWSY

acetaminophen, dextromethorphan hbr, phenylephrine hcl solution

Product Information

Item Code (Source)	NDC:41250-656
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH)	DEXTROMETHORPHAN	10 mg

(DEXTROMETHORPHAN - UNII:7355X3ROTS)	HYDROBROMIDE	in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
BUTYLATED HYDROXYANISOLE (UNII: REK4960K2U)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GLYCERIN (UNII: PDC6A3C0OX)	
MENTHOL, UNSPECIFIED FORM (UNII: L7T10EIP3A)	
SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM (UNII: 3980JIH2SW)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	ORANGE (clear)	Score	
Shape		Size	
Flavor	MENTHOL (with fruit)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-656-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	09/15/1989	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	10/24/2012	

Labeler - Meijer Distribution Inc (006959555)

