

**SEVERE CONGESTION AND COUGH RELIEF MAXIMUM STRENGTH-
dextromethorphan hbr, guaifenesin, phenylephrine hcl solution
L.N.K. International, Inc.**

Quality Plus 44-004

Active ingredients (in each 20 mL)

Dextromethorphan HBr 20 mg
Guaifenesin 400 mg
Phenylephrine HCl 10 mg

Purpose

Cough suppressant
Expectorant
Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help you get to sleep
 - nasal congestion due to a cold

Warnings

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- diabetes
- high blood pressure
- thyroid disease
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough persists more than 7 days, tends to recur, or is accompanied by a fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- do not take more than 6 doses in any 24-hour period
- mL = milliliter
- only use the dose cup provided
- adults and children 12 years and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years: do not use

Other information

- **each 20 mL contains:** sodium 9 mg
- use by expiration date on package
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)

Inactive ingredients

anhydrous citric acid, FD&C blue #1, FD&C red #40, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sorbitol, sucralose, xanthan gum

Questions or comments?

1-888-423-0139

Principal Display Panel

**QUALITY
+PLUS**

NDC 50844-004-45

*Compare to active ingredients
in Mucinex® FAST-MAX®
Severe Congestion & Cough

MAXIMUM
STRENGTH

**SEVERE
CONGESTION &
COUGH RELIEF**

Dextromethorphan HBr
Guaifenesin
Phenylephrine HCl

Cough Suppressant
Expectorant
Nasal Decongestant

Ages 12 Years and Over

Mixed
Berry
Flavored

6 FL OZ (177 mL)

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

PARENTS:

Learn about teen medicine abuse
www.StopMedicineAbuse.org

*This product is not manufactured or
distributed by RB Health (US) LLC, owner of
the registered trademark Mucinex®
Fast-Max® Severe Congestion & Cough.
50844 REV0423A00445

Distributed by
LNK INTERNATIONAL, INC.
60 Arkay Drive
Hauppauge, NY 11788
USA

NDC 50844-004-45

QUALITY PLUS

*Compare to active ingredients
in Mucinex® FAST-MAX®
Severe Congestion & Cough

MAXIMUM STRENGTH SEVERE CONGESTION & COUGH RELIEF

Dextromethorphan HBr
Guaifenesin
Phenylephrine HCl



**Cough Suppressant
Expectorant
Nasal Decongestant**

Mixed Berry Flavored

Ages 12 Years and Over

6 FL OZ (177 mL)

F-004-45
REV A

PEEL BACK TAB TO READ COMPLETE
DRUG FACTS AND INFORMATION

Drug Facts TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Active ingredients (in each 20 mL) Purpose
Dextromethorphan HBr 20 mg.....Cough suppressant
Guaifenesin 400 mg.....Expectorant
Phenylephrine HCl 10 mg.....Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help you get to sleep
 - nasal congestion due to a cold

PARENTS: Learn about teen medicine abuse
www.StopMedicineAbuse.org

Distributed by
LNK INTERNATIONAL, INC.
60 Arkay Drive
Hauppauge, NY 11788
USA



B-004-45
REV A

**No Print / No Varnish Area
Lot # and Exp. Info**

Drug Facts (continued)

Warnings
Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease ■ thyroid disease
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)
- high blood pressure ■ diabetes

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough persists more than 7 days, tends to recur, or is accompanied by a fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions

- do not take more than directed
- do not take more than 6 doses in any 24-hour period
- mL = milliliter
- only use the dose cup provided
- adults and children 12 years and over: 20 mL in dosing cup provided every 4 hours
- children under 12 years: do not use

Other information

- each 20 mL contains: sodium 9 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients anhydrous citric acid, FD&C blue #1, FD&C red #40, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sorbitol, sucralose, xanthan gum

Questions or comments?
1-800-426-9391

*This product is not manufactured or distributed by RB Health (US) LLC, owner of the registered trademark Mucinex® Fast-Max® Severe Congestion & Cough.
50844 REV0423A00445

Quality Plus 44-004

SEVERE CONGESTION AND COUGH RELIEF MAXIMUM STRENGTH

dextromethorphan hbr, guaifenesin, phenylephrine hcl solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50844-004
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	blue	Score	
Shape		Size	
Flavor	BERRY (MIXED)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50844-004-45	177 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/01/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/01/2017	

Labeler - L.N.K. International, Inc. (038154464)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	manufacture(50844-004) , pack(50844-004)

Revised: 8/2023

L.N.K. International, Inc.