

NAPROXEN- naproxen sodium tablet, film coated
L.N.K. International, Inc.

Convenience Valet 44-417

Active ingredient (in each tablet)

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*
*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - the common cold
 - toothache
 - menstrual cramps
 - backache
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- asthma (wheezing)
- skin reddening
- shock
- blisters
- rash
- facial swelling
- hives

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

- take more or for a longer time than directed
- have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - leg swelling
 - chest pain
 - slurred speech
 - trouble breathing
 - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose
- adults and children 12 years and older
 - take 1 tablet every 8 to 12 hours while symptoms last
 - for the first dose you may take 2 tablets within the first hour
 - do not exceed 2 tablets in any 8- to 12-hour period
 - do not exceed 3 tablets in a 24-hour period
- children under 12 years: ask a doctor

Other information

- **each tablet contains:** sodium 20 mg
- store at 20°-25°C (68°-77°F). Avoid high humidity and excessive heat above 40°C (104°F)
- see end flap for expiration date and lot number

Inactive ingredients

croscarmellose sodium, FD&C blue #2 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, talc, titanium dioxide

Questions or comments?

1-844-428-2538

Principal display panel

24/7 Life

BY 7-ELEVEN™

Naproxen

Naproxen sodium tablets 220 mg
Pain Reliever/Fever Reducer **(NSAID)**

STRENGTH TO LAST 12 HOURS

compare to

Aleve® Tablets

active ingredient**

24
TABLETS

Actual
Size

**QUALITY
GUARANTEED**

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**This product is not manufactured or distributed by
Bayer HealthCare LLC, distributor of Aleve® Tablets.
50844 REV1221B41708

DISTRIBUTED BY 7-ELEVEN, INC.
IRVING, TX 75063 WWW.7-ELEVEN.COM

Satisfaction Guaranteed 1-800-255-0711

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy, unless definitely directed to do so by a doctor, because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose
- adults and children 12 years and older
- take 1 tablet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 tablets within the first hour
- do not exceed 2 tablets in any 8- to 12-hour period
- do not exceed 3 tablets in a 24-hour period
- children under 12 years: ask a doctor

Other information

- each tablet contains sodium 20 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F)
- see end flap for expiration date and lot number

Inactive ingredients crosscarmellose sodium, FD&C blue #7 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, polyvinylpyrrolidone, silicon dioxide, talc, titanium dioxide

Questions or comments? 1-844-429-2538

This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Aleve® Tablets. 50844 REV1221B41708 5628C-SE-01-21

DISTRIBUTED BY 7-ELEVEN, INC. WWW.7-ELEVEN.COM IRVING, TX 75063

Satisfaction Guaranteed 1-800-255-0711

24 TABLETS Actual Size

Naproxen
Naproxen sodium tablets 220 mg
Pain Reliever/Fever Reducer (NSAID)
STRENGTH TO LAST 12 HOURS

compare to Aleve® Tablets active ingredient**

QUALITY GUARANTEED

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SHEET SEAL UNDER CAP IS BROKEN OR MISSING

0 52548 68982 7

No Print / No Varnish
Lot no. & Exp. date

Drug Facts **KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION**

Active ingredient (in each tablet) Purpose
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)* Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - the common cold
 - toothache
 - menstrual cramps
 - backache
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- asthma (wheezing)
- skin redness
- shock
- hives
- blisters
- facial swelling
- throat swelling

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- have a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- take more or for a longer time than directed
- have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease the benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- stop use and ask a doctor if you experience any of the following signs of stomach bleeding:
 - red, black, or vomit blood
 - dark stools
 - stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - leg swelling
 - chest pain
 - sturred speech
 - trouble breathing
- weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

B-1205-417-08
REV1221B41708

Convenience Valet 44-417

| | | | |
|--|----------------|---------------------------|---------------|
| NAPROXEN | | | |
| naproxen sodium tablet, film coated | | | |
| Product Information | | | |
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:50844-261 |
| Route of Administration | ORAL | | |
| Active Ingredient/Active Moiety | | | |

| Ingredient Name | | Basis of Strength | Strength | |
|---|--|--|----------------------|--------------------|
| NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ) | | NAPROXEN SODIUM | 220 mg | |
| Inactive Ingredients | | | | |
| Ingredient Name | | | Strength | |
| CROSCARMELOSE SODIUM (UNII: M28OL1HH48) | | | | |
| FD&C BLUE NO. 2--ALUMINUM LAKE (UNII: 4AQJ3LG584) | | | | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | | | | |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U) | | | | |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A) | | | | |
| POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990) | | | | |
| POVIDONE, UNSPECIFIED (UNII: FZ989GH94E) | | | | |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | | | | |
| TALC (UNII: 7SEV7J4R1U) | | | | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | | | | |
| Product Characteristics | | | | |
| Color | blue | Score | no score | |
| Shape | ROUND | Size | 10mm | |
| Flavor | | Imprint Code | 44;417 | |
| Contains | | | | |
| Packaging | | | | |
| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
| 1 | NDC:50844-261-08 | 1 in 1 CARTON | 08/06/2019 | |
| 1 | | 24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | | |
| Marketing Information | | | | |
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date | |
| ANDA | ANDA204872 | 08/06/2019 | | |

Labeler - L.N.K. International, Inc. (038154464)

Establishment

| Name | Address | ID/FEI | Business Operations |
|-------------------------|---------|-----------|---------------------|
| LNK International, Inc. | | 038154464 | pack(50844-261) |

Establishment

| Name | Address | ID/FEI | Business Operations |
|------|---------|--------|---------------------|
|------|---------|--------|---------------------|

| | | | |
|-------------------------|--|-----------|------------------------|
| LNK International, Inc. | | 832867837 | manufacture(50844-261) |
|-------------------------|--|-----------|------------------------|

Establishment

| Name | Address | ID/FEI | Business Operations |
|-------------------------|---------|-----------|------------------------|
| LNK International, Inc. | | 832867894 | manufacture(50844-261) |

Establishment

| Name | Address | ID/FEI | Business Operations |
|-------------------------|---------|-----------|---------------------|
| LNK International, Inc. | | 967626305 | pack(50844-261) |

Establishment

| Name | Address | ID/FEI | Business Operations |
|-------------------------|---------|-----------|------------------------|
| LNK International, Inc. | | 117025878 | manufacture(50844-261) |

Revised: 8/2023

L.N.K. International, Inc.