

ASPIRIN- aspirin tablet, delayed release
Walgreen Company

Walgreens 44-587A

Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

for the temporary relief of minor aches and pains or as recommended by your doctor.
Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have had stomach ulcers or bleeding problems
- take more or for a longer time than directed
- have 3 or more alcoholic drinks every day while using this product

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- you have asthma
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - vomit blood
 - have bloody or black stools
 - feel faint
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

colloidal anhydrous silica, corn starch, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

Questions or comments?

1-800-426-9391

Principal Display Panel

Walgreens

NDC 0363-0587-32

Compare to the active ingredient
in Bayer® Low Dose Aspirin††

Dye-Free Aspirin 81

81 mg / PAIN RELIEVER (**NSAID**)

Low Dose

Enteric-Coated

120 TABLETS

ACTUAL SIZE

****Talk to your doctor or other healthcare
provider before using this product for your heart.**

DISTRIBUTED BY: **WALGREEN CO.**
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2021 Walgreen Co.

50844 ORG012258732

†Our pharmacists recommend the
Walgreens brand. We invite you to
compare to national brands.

††This product is not manufactured or
distributed by Bayer AG, owner of
the registered trademark Bayer®
Low Dose Aspirin.

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY
SEAL UNDER CAP IS BROKEN OR MISSING**

W30RG1221-F
REV0222

ITEM 500645 W00000-0000-0
11917110729 5

B-2201-587A-32-HR
ORG012258732

Walgreens

Compare to the active ingredient
in Bayer® Low Dose Aspirin^{††}

Dye-Free Aspirin 81
81 mg / PAIN RELIEVER (NSAID)

Low Dose Enteric-Coated

120 TABLETS

HEART HEALTH

ACTUAL SIZE

**Talk to your doctor or other healthcare provider before using this product for your heart.

NDC 0363-0587-32

Drug Facts KEEP OUT OF REACH OF CHILDREN. READ THE ENTIRE CONTENTS OF THE PRODUCT INFORMATION.

Active ingredient (in each tablet) Aspirin 81 mg (NSAID) *Pain reliever, nonsteroidal anti-inflammatory drug

Purpose For the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. Allergy alert: Aspirin may cause a severe allergic reaction, which may include:
■ hives ■ facial swelling ■ shock ■ asthma (wheezing)
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you take a blood thinning (anticoagulant) or steroid drug ■ are age 60 or older ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Drug Facts (continued)

■ have had stomach ulcers or bleeding problems
■ take more or for a longer time than directed
■ have 3 or more alcoholic drinks every day while using this product

Do not use ■ If you are allergic to aspirin or any other pain reliever/fever reducer
■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if ■ you have asthma ■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you are taking a diuretic ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug for
■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if
■ an allergic reaction occurs. Seek medical help right away.
■ you experience any of the following signs of stomach bleeding: ■ vomit blood ■ have bloody or black stools ■ feel faint ■ have stomach pain that does not get better ■ ringing in the ears or a loss of hearing occurs ■ pain gets worse or lasts more than 10 days ■ ever gets worse or lasts more than 3 days ■ redness or swelling is present
These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ do not take more than directed
■ drink a full glass of water with each dose
■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 40 tablets in 24 hours unless directed by a doctor
■ children under 12 years: do not use unless directed by a doctor

Other information
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
■ see end flap for expiration date and lot number

Inactive ingredients colloidal anhydrous silica, corn starch, hypromellose, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, stearic wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, talc, titanium dioxide, triacetin, triethyl citrate

(Questions or comments?) 1-800-426-9391

how2recycle.info

PAPER PLASTIC BOX BOTTLE

Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.
†† This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Low Dose Aspirin.

50844 ORG012258732
DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2021 Walgreen Co.

Walgreens 44-587A

ASPIRIN

aspirin tablet, delayed release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0587
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)		ASPIRIN	81 mg	
Inactive Ingredients				
Ingredient Name			Strength	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
STARCH, CORN (UNII: O8232NY3SJ)				
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)				
METHACRYLIC ACID (UNII: 1CS02G8656)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
POLYDEXTROSE (UNII: VH2XOU12IE)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
SHELLAC (UNII: 46N107B71O)				
DIMETHICONE (UNII: 92RU3N3Y1O)				
WATER (UNII: 059QF0KO0R)				
SODIUM BICARBONATE (UNII: 8MDF5V39QO)				
SODIUM LAURYL SULFATE (UNII: 368GB5141J)				
TALC (UNII: 7SEV7J4R1U)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
TRIACETIN (UNII: XHX3C3X673)				
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)				
Product Characteristics				
Color	white	Score	no score	
Shape	ROUND	Size	6mm	
Flavor		Imprint Code	L	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0587-32	1 in 1 CARTON	03/18/2009	
1		120 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:0363-0587-14	500 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/18/2009	
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M013	03/18/2009	

Labeler - Walgreen Company (008965063)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(0363-0587)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(0363-0587)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(0363-0587)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	manufacture(0363-0587)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(0363-0587)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(0363-0587)

Revised: 4/2023

Walgreen Company