

SOOTHE- bismuth subsalicylate liquid

Walgreens

Drug Facts

Active ingredient (in each 30 mL)

Bismuth subsalicylate 525 mg

Purpose

Upset stomach reliever/Antidiarrheal

Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
 - heartburn
 - indigestion
 - nausea
 - gas
 - fullness
 - belching

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use

if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are

taking any drug for

- diabetes
- gout
- arthritis
- anticoagulation (thinning the blood)

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- mL = milliliter
- shake well before using
- measure only with dosing cup provided. Do not use any other dosing device
- keep dosing cup with product
- adult and children 12 years and over:
 - 30 mL (1 dose) every 1/2 to 1 hour as needed
 - do not exceed 8 doses (240 mL) in 24 hour
 - use until diarrhea stops but not more than 2 days
- children under 12 years of age: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- **each 30 mL contains:** sodium 10 mg
- **each 30 mL contains:** salicylate 248 mg
- low sodium
- keep tightly closed
- protect from freezing
- avoid excessive heat (over 104°F or 40°C)

Inactive ingredients

benzoic acid, D&C red #22, D&C red #28, flavor, magnesium aluminum silicate, methylcellulose, purified water, saccharin sodium, salicylic acid, sodium salicylate, sorbic acid, sucralose

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to Pepto-Bismol® active ingredient††

Soothe®

BISMUTH SUBSALICYLATE 525 mg /

UPSET STOMACH RELIEVER / ANTIDIARRHEAL

SUGAR FREE

ALCOHOL FREE

- Provides protective coating action to relieve heartburn, indigestion, nausea, upset stomach & diarrhea

CHERRY FLAVOR

FL OZ (mL)

††This product is not manufactured or distributed by Procter & Gamble Company. Pepto-Bismol® is a registered trademark of th Procter & Gamble Company

TAMPER EVIDENT; DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD, DEERFIELD.IL 60015

walgreens.com

Package Label

Drug Facts (continued)

Directions ■ mL = milliliter ■ shake well before using

- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- adults and children 12 years and over
 - 30 mL (1 dose) every ½ to 1 hour as needed
 - do not exceed 8 doses (240 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years of age: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information ■ each 30 mL contains: sodium 10 mg

- each 30 mL contains: salicylate 248 mg
- low sodium ■ keep tightly closed ■ protect from freezing
- avoid excessive heat (over 104°F or 40°C)

Inactive ingredients benzoic acid, D&C red #22, D&C red #28, flavor, magnesium aluminum silicate, methylcellulose, purified water, saccharin sodium, salicylic acid, sodium salicylate, sorbic acid, sucralose

Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Walgreens Pharmacist Recommended
Walgreens Pharmacist Survey

This product is not manufactured or distributed by The Procter & Gamble Company. Pepto-Bismol® is a registered trademark of The Procter & Gamble Company.

PLD-D384C
LB006430

AUTO-REORDER & SAVE
Learn more at Walgreens.com/AutoReorder

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2018 Walgreen Co.
ORG0119-F

ITEM 561661 W10023-0119-F

3 11917 08392 6

Walgreens

Compare to Pepto-Bismol® active ingredient!†

NDC 0363-0025-16

Soothe®

BISMUTH SUBSALICYLATE 525 mg /
UPSET STOMACH RELIEVER /
ANTIDIARRHEAL

SUGAR FREE

ALCOHOL FREE

- Provides protective coating action to relieve heartburn, nausea, indigestion, upset stomach & diarrhea

16 FL OZ (1 PT) (473 mL) CHERRY FLAVOR



Drug Facts **TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING**

Active ingredient (in each 30 mL) Bismuth subsalicylate 525 mg **Purpose** Upset stomach reliever/Antidiarrheal

Uses relieves ■ diarrhea ■ travelers' diarrhea

- upset stomach due to overindulgence in food and drink, including
- heartburn ■ gas ■ indigestion
- nausea ■ fullness ■ belching

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. **Allergy alert: Contains salicylate. Do not take if you are:**

- allergic to salicylates (including aspirin) ■ taking other salicylate products

Do not use if you have

- an ulcer ■ a bleeding problem ■ bloody or black stool

Ask a doctor before use if you have

- fever ■ mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for

- diabetes ■ gout ■ arthritis ■ anticoagulation (thinning the blood)

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

WALGREENS Soothe

SOOTHE

bismuth subsalicylate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-0025
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE, SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
BENZOIC ACID (UNII: 8SKN0B0MIM)	
D&C RED NO. 22 (UNII: 1678RKX8RT)	
D&C RED NO. 28 (UNII: 767IP0Y5NH)	
MAGNESIUM ALUMINUM SILICATE (UNII: 6M3P64V0NC)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SALICYLIC ACID (UNII: O414PZ4LPZ)	
SODIUM SALICYLATE (UNII: WMQ1H85SYP)	
SORBIC ACID (UNII: X045WJ989B)	
METHYLCELLULOSE (1500 CPS) (UNII: PONTE48364)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-0025-16	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/31/2015	
2	NDC:0363-0025-08	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/31/2015	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	01/31/2015	

Labeler - Walgreens (008965063)

Revised: 10/2023

Walgreens