

## **UP AND UP CHILDRENS IBUPROFEN- ibuprofen suspension**

**Target Corporation**

-----

### **Target Corporation Children's Ibuprofen Oral Suspension Drug Facts**

#### **Active ingredient (in each 5 mL = 1 teaspoonful)**

Ibuprofen 100 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

#### **Purposes**

Pain reliever/fever reducer

#### **Uses**

temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

#### **Warnings**

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

#### **Do not use**

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### **Ask a doctor before use if**

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

### **Ask a doctor or pharmacist before use if the child is**

- under a doctor's care for any serious condition
- taking any other drug

### **When using this product**

- give with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

### **Stop use and ask a doctor if**

- child experiences any of the following signs of stomach bleeding
- feels faint
- vomits blood
- has bloody or black stools
- has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### **Directions**

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter; tsp = teaspoonful
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.

- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance
- wash dosage cup after each use

Dosing Chart		
Weight (lb)	Age (yr)	Dose (mL or tsp)**
under 24 lbs	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL (1 tsp)
36-47 lbs	4-5 years	7.5 mL (1½ tsp)
48-59 lbs	6-8 years	10 mL (2 tsp)
60-71 lbs	9-10 years	12.5 mL (2½ tsp)
72-95 lbs	11 years	15 mL (3 tsp)

\*\*or as directed by a doctor

### Other information

- each 5 mL (1 tsp) contains: sodium 2 mg
- do not use if printed neckband is broken or missing
- store at 20-25°C (68-77°F)
- do not freeze
- see bottom panel for lot number and expiration date

### Inactive ingredients

anhydrous citric acid, D&C red #33, FD&C blue #1, FD&C red #40, glycerin, high fructose corn syrup, hypromellose, natural and artificial grape flavor, polysorbate 80, purified water, sodium benzoate, sorbitol solution, xanthan gum

### Questions?

Call 1-888-547-7400

### Principal Display Panel

Compare to active ingredient in Children's Motrin® Grape Flavor

children's ibuprofen

oral suspension 100 mg per 5 mL

pain reliever/fever reducer (NSAID)

alcohol free

LASTS UP TO 8 HOURS

GRAPE FLAVOR

AGES 2 TO 11 YEARS

4 FL OZ (120 mL)



NDC 11673-660-26

Compare to active ingredient in Children's Motrin® Grape Flavor\*\*\*

# children's ibuprofen

oral suspension  
100 mg per 5 mL  
pain reliever/fever reducer (NSAID)

alcohol free



4 FL OZ (120 mL)



Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

### Drug Facts

**Active ingredient** (in each 5 mL = 1 teaspoonful)  
Ibuprofen 100 mg (NSAID) .....Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses** temporarily:  
■ relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache  
■ reduces fever

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- asthma (wheezing)
- skin reddening
- facial swelling
- shock
- rash

■ blisters  
If an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

■ takes more or for a longer time than directed  
**Sore throat warning:** Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

**Do not use**  
■ if the child has ever had an allergic reaction to any other pain reliever/fever reducer  
■ right before or after heart surgery

**Ask a doctor before use if**  
■ stomach bleeding warning applies to your child  
■ child has a history of stomach problems, such as heartburn



oral suspension  
100 mg per 5 mL  
pain reliever/fever reducer (NSAID)

# children's ibuprofen

### Drug Facts (continued)

- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

**Ask a doctor or pharmacist before use if the child is**  
■ under a doctor's care for any serious condition  
■ taking any other drug

**When using this product**  
■ give with food or milk if stomach upset occurs  
■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

**Stop use and ask a doctor if**  
■ child experiences any of the following signs of stomach bleeding  
■ feels faint ■ vomits blood  
■ has bloody or black stools  
■ has stomach pain that does not get better  
■ the child does not get any relief within first day (24 hours) of treatment  
■ fever or pain gets worse or lasts more than 3 days  
■ redness or swelling is present in the painful area  
■ any new symptoms appear

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- mL = milliliter; tsp = teaspoonful
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance
- wash dosage cup after each use

### Drug Facts (continued)

#### Dosing Chart

Weight (lb)	Age (yr)	Dose (mL or tsp)**
under 24 lbs	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL (1 tsp)
36-47 lbs	4-5 years	7.5 mL (1½ tsp)
48-59 lbs	6-8 years	10 mL (2 tsp)
60-71 lbs	9-10 years	12.5 mL (2½ tsp)
72-95 lbs	11 years	15 mL (3 tsp)

\*\*or as directed by a doctor

### Other information

- each 5 mL (1 tsp) contains: sodium 2 mg
- do not use if printed neckband is broken or missing
- store at 20-25°C (68-77°F)
- do not freeze
- see bottom panel for lot number and expiration date

**Inactive ingredients** anhydrous citric acid, D&C red #33, FD&C blue #1, FD&C red #40, glycerin, high fructose corn syrup, hypromellose, natural and artificial grape flavor, polysorbate 80, purified water, sodium benzoate, sorbitol solution, xanthan gum

**Questions? Call 1-888-547-7400**

\*\*\*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Motrin® Grape Flavor.

**GLUTEN FREE**

100% satisfaction guaranteed or your money back.

We welcome any questions you may have at [Target.com/comments](http://Target.com/comments) or 1-800-910-6874

094 01 0245 R02 10225432  
Distributed by Target Corporation  
Minneapolis, MN 55403  
© & ©2015 Target Brands, Inc.  
[Shop.Target.com](http://Shop.Target.com)



LOT NO.

EXP.

: 66026 UV CL

## UP AND UP CHILDRENS IBUPROFEN

ibuprofen suspension

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11673-660
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	100 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C BLUE NO. 1 (UNII: HBR47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
WATER (UNII: 059QF0K00R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
XANTHAN GUM (UNII: TTV12P4NEE)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	

### Product Characteristics

Color	PURPLE (opaque)	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-660-26	1 in 1 CARTON		
1		120 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:11673-660-28	1 in 1 CARTON		
2		148 mL in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074937	07/08/2009	

**Labeler** - Target Corporation (006961700)