

**OXY 500 CALF BOLUS- oxytetracycline hydrochloride tablet**  
**Boehringer Ingelheim Animal Health USA Inc.**

-----  
**Oxy 500 Calf Bolus**  
**(oxytetracycline hydrochloride)**

Approved by FDA under NADA # 141-002

**For the control and treatment of bacterial enteritis (scours, diarrhea) and bacterial pneumonia in beef and dairy calves.**

Read entire leaflet carefully before using this product.

**Each bolus contains:** 500 mg oxytetracycline hydrochloride.

**Description:**

OXY 500 CALF BOLUS is an oral formulation containing oxytetracycline, a broad-spectrum antibiotic that possesses a high order of stability and potent antimicrobial activity, for use in beef and dairy calves.

**Indications:**

OXY 500 CALF BOLUS is recommended for oral administration for the control and treatment of the following diseases of beef and dairy calves caused by organisms sensitive to oxytetracycline: bacterial enteritis (scours, diarrhea) caused by *Salmonella typhimurium* and *Escherichia coli* (colibacillosis); bacterial pneumonia (shipping fever complex, pasteurellosis) caused by *Pasteurella multocida*.

**Dosage:**

For the **control** of bacterial enteritis (scours, diarrhea) and bacterial pneumonia in beef and dairy calves: administer orally half a bolus (250 mg oxytetracycline hydrochloride) per 100 lb of body weight every 12 hours (5 mg/lb body weight in divided doses) for up to four consecutive days.

For the **treatment** of bacterial enteritis (scours, diarrhea) and bacterial pneumonia in beef and dairy calves: administer orally one bolus (500 mg oxytetracycline hydrochloride) per 100 lb of body weight every 12 hours (10 mg/lb body weight in divided doses) for up to four consecutive days.

Dosage should continue until the animal returns to normal and for 24 to 48 hours after symptoms have subsided. Treatment should not exceed four consecutive days.

**Residue Warning:**

There is no pre-slaughter withdrawal period when used at the recommended dosage level. Not for use in lactating dairy cattle. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

**Precautions:**

Exceeding the recommended dosage level of one bolus (500 mg oxytetracycline hydrochloride) per 100 lb body weight every 12 hours, or administration at the recommended level for more than four consecutive days, may result in antibiotic residues beyond the withdrawal time.

Organisms may vary in their degree of susceptibility to any chemotherapy. If no improvement is observed after recommended treatment, diagnosis and susceptibility should be reexamined.

Rarely do side reactions or allergic manifestations occur in calves treated with OXY 500 CALF BOLUS. If any unusual reactions are noted, discontinue use of the drug immediately and call a veterinarian.

Since bacteriostatic drugs may interfere with the bactericidal action of penicillin, it is advisable to avoid giving OXY 500 CALF BOLUS in conjunction with penicillin.

To report suspected adverse events, for technical assistance, or to obtain a copy of the Safety Data Sheet, contact Boehringer Ingelheim Animal Health USA Inc. at 1-888-637-4251.

For additional information about adverse drug experience reporting for animal drugs, contact FDA by telephone at 1-888-FDA-VETS or online at [www.fda.gov/reportanimalae](http://www.fda.gov/reportanimalae).

**Care of Sick Animals:**

The use of antibiotics in the management of disease is based on an accurate diagnosis and an adequate course of treatment. When properly used in the treatment of disease caused by oxytetracycline susceptible organisms, most animals treated with OXY 500 CALF BOLUS show a noticeable improvement within 24 to 48 hours. It is recommended that the diagnosis and treatment of animal diseases be carried out by a veterinarian. Since many diseases look alike but require different types of treatment, the use of professional veterinary and laboratory services can reduce treatment time, costs and needless losses. Good housing, sanitation and nutrition are important in the maintenance of healthy animals and are essential in the treatment of disease.

**Storage:**

Store at 20° to 25°C (68° to 77°F), excursions permitted between 15° and 30°C (between 59° and 86°F).

**How Supplied**

OXY 500 CALF BOLUS is available in 25 and 100 bolus containers.

NDC 0010-6834-01 - 25 boluses

NDC 0010-6834-02 - 100 boluses

**Marketed by:**

Boehringer Ingelheim Animal Health USA Inc.

Duluth, GA 30096

683401-04

87018350

Rev: 08/19

## Principal Display Panel - 25 bolus bottle label

NDC 0010-6834-01

### Oxy 500 Calf Bolus

(oxytetracycline HCl), 500 mg

**For the control and treatment of bacterial enteritis (scours, diarrhea) and bacterial pneumonia in beef and dairy calves.**

**Net Contents:** 25 boluses

Approved by FDA under NADA # 141-002

**Each bolus contains:** 500 mg oxytetracycline hydrochloride.

**Indications:** OXY 500 CALF BOLUS is recommended for oral administration for the control and treatment of the following diseases of beef and dairy calves caused by organisms sensitive to oxytetracycline: bacterial enteritis (scours, diarrhea) caused by *Salmonella typhimurium* and *Escherichia coli* (colibacillosis); bacterial pneumonia (shipping fever complex, pasteurellosis) caused by *Pasteurella multocida*.

**Dosage:** For the **control** of bacterial enteritis (scours, diarrhea) and bacterial pneumonia in beef and dairy calves: administer orally half a bolus (250 mg oxytetracycline hydrochloride) per 100 lb of body weight every 12 hours (5 mg/lb body weight in divided doses) for up to four consecutive days.

For the **treatment** of bacterial enteritis (scours, diarrhea) and bacterial pneumonia in beef and dairy calves: administer orally one bolus (500 mg oxytetracycline hydrochloride) per 100 lb of body weight every 12 hours (10 mg/lb body weight in divided doses) for up to four consecutive days. Dosage should continue until the animal returns to normal and for 24 to 48 hours after symptoms have subsided. Treatment should not exceed four consecutive days.

**Residue Warning:** There is no pre-slaughter withdrawal period when used at the recommended dosage level. Not for use in lactating dairy cattle. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

Refer to folded package insert attached to this container for complete product information.

**Storage:** Store at 20° to 25°C (68° to 77°F), excursions permitted between 15° and 30°C (between 59° and 86°F).

For use in animals only.

**Marketed by:**  
Boehringer Ingelheim Animal Health USA Inc.  
Duluth, GA 30096  
683401-04 87018350

## OXY 500 CALF BOLUS

oxytetracycline hydrochloride tablet

### Product Information

<b>Product Type</b>	OTC ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:0010-6834
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>OXYTETRACYCLINE HYDROCHLORIDE</b> (UNII: 4U7K4N52ZM) (OXYTETRACYCLINE ANHYDROUS - UNII:SLF0D9077S)	OXYTETRACYCLINE HYDROCHLORIDE	500 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>DEXTRATES</b> (UNII: G263MI44RU)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	

**Product Characteristics**

<b>Color</b>	YELLOW	<b>Score</b>	2 pieces
<b>Shape</b>	OVAL	<b>Size</b>	21mm
<b>Flavor</b>		<b>Imprint Code</b>	Oxy500
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0010-6834-01	25 in 1 BOTTLE, PLASTIC		
2	NDC:0010-6834-02	100 in 1 BOTTLE, PLASTIC		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA141002	05/24/2000	

**Labeler** - Boehringer Ingelheim Animal Health USA Inc. (007134091)

Revised: 12/2021

Boehringer Ingelheim Animal Health USA Inc.