


## **NYSTATIN- nystatin ointment Bryant Ranch Prepack**

### **Nystatin Ointment, USP**

#### **DESCRIPTION**

Nystatin is a polyene antifungal antibiotic drug obtained from *Streptomyces noursei*.

Structural formula:



Nystatin Ointment is for dermatologic use. Nystatin Ointment USP, for topical use only, contains 100,000 USP Nystatin Units per gram, in a white petrolatum and light mineral oil base.

#### **CLINICAL PHARMACOLOGY**

##### **Pharmacokinetics**

Nystatin is not absorbed from intact skin or mucous membrane.

##### **Microbiology**

Nystatin is an antibiotic which is both fungistatic and fungicidal *in vitro* against a wide variety of yeasts and yeast-like fungi, including *Candida albicans*, *C. parapsilosis*, *C. tropicalis*, *C. guilliermondii*, *C. pseudotropicalis*, *C. krusei*, *Torulopsis glabrata*, *Trichophyton rubrum*, *T. mentagrophytes*.

Nystatin acts by binding to sterols in the cell membrane of susceptible species resulting in a change in membrane permeability and the subsequent leakage of intracellular components. On repeated subculturing with increasing concentrations of nystatin,

*Candida albicans* does not develop resistance to nystatin. Generally, resistance to nystatin does not develop during therapy. However, other species of *Candida* (*C. tropicalis*, *C. guilliermondi*, *C. krusei*, and *C. stellatoidea*) become quite resistant on treatment with nystatin and simultaneously become cross resistant to amphotericin as well. This resistance is lost when the antibiotic is removed.

Nystatin exhibits no appreciable activity against bacteria, protozoa, or viruses.

## **INDICATIONS AND USAGE**

Nystatin Ointment is indicated in the treatment of cutaneous or mucocutaneous mycotic infections caused by *Candida albicans* and other susceptible *Candida* species.

**Nystatin Ointment is not indicated for systemic, oral, intravaginal or ophthalmic use.**

## **CONTRAINDICATIONS**

Nystatin Ointment is contraindicated in patients with a history of hypersensitivity to any of their components.

## **PRECAUTIONS**

### **General**

**Nystatin Ointment should not be used for the treatment of systemic, oral, intravaginal or ophthalmic infections.**

If irritation or sensitization develops, treatment should be discontinued and appropriate measures taken as indicated. It is recommended that KOH smears, cultures, or other diagnostic methods be used to confirm the diagnosis of cutaneous or mucocutaneous candidiasis and to rule out infection caused by other pathogens.

## **INFORMATION FOR PATIENTS**

**Patients using these medications should receive the following information and instructions:**

1. The patient should be instructed to use these medications as directed (including the replacement of missed doses). These medications are not for any disorder other than that for which they are prescribed.
2. Even if symptomatic relief occurs within the first few days of treatment, the patient should be advised not to interrupt or discontinue therapy until the prescribed course of treatment is completed.
3. If symptoms of irritation develop, the patient should be advised to notify the physician promptly

### **Laboratory Tests**

If there is a lack of therapeutic response, KOH smears, cultures, or other diagnostic methods should be repeated.

## **Carcinogenesis, Mutagenesis, Impairment of Fertility**

No long-term animal studies have been performed to evaluate the carcinogenic potential of nystatin. No studies have been performed to determine the mutagenicity of nystatin or its effects on male or female fertility.

## **Pregnancy: Teratogenic Effects**

Category C.

Animal reproduction studies have not been conducted with any nystatin topical preparation. It also is not known whether these preparations can cause fetal harm when used by a pregnant woman or can affect reproductive capacity. Nystatin topical preparations should be prescribed for a pregnant woman only if the potential benefit to the mother outweighs the potential risk to the fetus.

## **Nursing Mothers**

It is not known whether nystatin is excreted in human milk. Caution should be exercised when nystatin is prescribed for a nursing woman.

## **Pediatric Use**

Safety and effectiveness have been established in the pediatric population from birth to 16 years.

See **DOSAGE AND ADMINISTRATION**

## **ADVERSE REACTIONS**

The frequency of adverse events reported in patients using Nystatin Ointment preparations is less than 0.1%. The more common events that were reported include allergic reactions, burning, itching, rash, eczema, and pain on application.

(See **PRECAUTIONS: General.**)

## **DOSAGE AND ADMINISTRATION**

### **NYSTATIN Ointment**

#### **Adults and Pediatric Patients (Neonates and Older):**

Apply liberally to affected areas twice daily or as indicated until healing is complete.

## **HOW SUPPLIED**

Nystatin ointment (100,000 USP Nystatin Units per gram) is a yellow colored ointment is supplied in 30g tubes.

NDC: 63629-2515-1: in tube of 30 gm

## **STORAGE**

NYSTATIN Ointment: Store at room temperature.

Call your doctor for medical advice about side effects. You may report side effects to Viona Pharmaceuticals Inc. at 1-888-304-5011 or FDA at 1-800-FDA-1088.

**Manufactured by:**

Cadila Healthcare Limited  
Changodar, Ahmedabad, India.

**Distributed by:**

**Viona Pharmaceuticals Inc.**

Cranford, NJ 07016

Rev.: 11/19

**Nystatin 100,000u/g Oint, USP #30**



Each gram contains: 100,000 USP Nystatin units in an ointment base of light mineral oil and white petrolatum.

Keep this and all medication out of the reach of children.

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

For Topical Use Only. Not for Ophthalmic Use.

**NDC 63629-2515-1**

**Nystatin Ointment, USP**

**100,000 units/gm**



Relabeled by:  
Bryant Ranch Prepack, Inc.  
Burbank, CA 91504 USA

Rx only  
**30 grams**



**NYSTATIN**

nystatin ointment

**Product Information**

|                                |                         |                           |                               |
|--------------------------------|-------------------------|---------------------------|-------------------------------|
| <b>Product Type</b>            | HUMAN PRESCRIPTION DRUG | <b>Item Code (Source)</b> | NDC:63629-2515(NDC:72578-089) |
| <b>Route of Administration</b> | TOPICAL                 |                           |                               |

**Active Ingredient/Active Moiety**

| <b>Ingredient Name</b>                                   | <b>Basis of Strength</b> | <b>Strength</b> |
|--|--------------------------|-----------------|
| NYSTATIN (UNII: BDF1O1C72E) (NYSTATIN - UNII:BDF1O1C72E) | NYSTATIN                 | 100000 U in 1 g |

**Inactive Ingredients**

| Ingredient Name                |  | Strength  |                      |                    |
|--------------------------------|--|---|----------------------|--------------------|
| MINERAL OIL (UNII: T5L8T28FGP) |  |   |                      |                    |
| PETROLATUM (UNII: 4T6H12BN9U)  |  |   |                      |                    |
| <b>Product Characteristics</b> |  |   |                      |                    |
| Color                          | YELLOW (YELLOW)                          | Score   |                      |                    |
| Shape                          |  | Size  |                      |                    |
| Flavor                         |  | Imprint Code                                      |                      |                    |
| Contains                       |  |   |                      |                    |
| <b>Packaging</b>               |  |   |                      |                    |
| #                              | Item Code                                | Package Description                               | Marketing Start Date | Marketing End Date |
| 1                              | NDC:63629-2515-1                         | 30 g in 1 TUBE; Type 0: Not a Combination Product | 05/18/2021           |                    |
| <b>Marketing Information</b>   |  |   |                      |                    |
| Marketing Category             | Application Number or Monograph Citation | Marketing Start Date                              | Marketing End Date   |                    |
| ANDA                           | ANDA207767                               | 06/24/2020  |                      |                    |

**Labeler** - Bryant Ranch Prepack (171714327)

**Registrant** - Bryant Ranch Prepack (171714327)

### Establishment

| Name                 | Address | ID/FEI    | Business Operations                      |
|----------------------|---------|-----------|--|
| Bryant Ranch Prepack |         | 171714327 | REPACK(63629-2515) , RELABEL(63629-2515) |

Revised: 3/2023

Bryant Ranch Prepack